

Original Research Article

In vivo Evaluation of a Cosmetic Emulsion Containing Soybean Extract for Anti-Aging

MK Waqas¹, N Akhtar¹, A Rasul², SU Rashid³, R Mustafa¹, BA Khan⁴ and G Murtaza^{5*}

¹Department of Pharmacy, Faculty of Pharmacy and Alternative Medicines, The Islamia University of Bahawalpur, Bahawalpur, ²College of Pharmacy, Government College University Faisalabad, Faisalabad, ³Faculty of Pharmacy, University of Sargodha, Sargodha, ⁴Faculty of Pharmacy, Gomal University, Dera Ismail Khan, ⁵Department of Pharmacy, COMSATS Institute of Information Technology, Abbottabad, 22060 Pakistan

*For correspondence: **Email:** gmdogar356@gmail.com; **Tel:** 0092-314-2082826; **Fax:** 0092-922-383441

Received: 10 May 2014

Revised accepted: 6 August 2014

Abstract

Purpose: To develop and assess the anti-aging potential of a cosmetic W/O emulsion containing an extract of soybean, *Glycine max* (L.) Merr. Fabaceae.

Methods: This single-blind placebo-controlled study was performed in 11 healthy male human volunteers. A formulation comprising of 4 % of concentrated extract of soybean was prepared by loading the extract in the internal aqueous phase of the emulsion. A control (base), consisting of the emulsion without the extract, was also prepared. Both formulations were applied to the cheeks of all volunteers for 12 weeks and their effect on different skin parameters, i.e., moisture contents, elasticity and surface evaluation of living skin (SELS) were assessed.

Results: The formulation containing 4 % soybean extract showed significant ($p \leq 0.05$) effects on skin elasticity and moisture contents but the base showed insignificant effect ($p > 0.05$). There was significant ($p \leq 0.05$) decline in SELS, i.e., SEsc (skin scaliness, from 1.73 ± 0.05 to 1.66 ± 0.06), SEw (skin wrinkles, from 71.74 ± 1.52 to 68.51 ± 1.64), SEsm (skin smoothness, from 109.01 ± 4.77 to 102.03 ± 4.23), and SEr (skin roughness, from 4.04 ± 0.09 to 3.82 ± 0.08) parameters after application of the extract formulation for 12 weeks.

Conclusion: Topical application of the cosmetic emulsion containing soybean extract exerts potential skin anti-aging effects.

Keywords: *Glycine max*, Soybean, Anti-aging, Skin elasticity, Cosmetic emulsion, Surface evaluation of living skin (SELS), Skin moisture content

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Skin aging is a complicated procedure promoted by persistent exposure to UV radiation. UV radiations produce the ROS (reactive oxygen species) assisting in collagen deficit and ultimately skin wrinkling. Skin aging is depicted by increased wrinkling, decline in elasticity, uneven pigmentation, increased roughness and

dryness [1]. Oil-in-water or water-in-oil emulsions are among the most frequently used colloidal systems, largely present in many fields including cosmetics, pharmaceuticals, food, paints, and petrochemicals [2,3]. A wide variety of cosmetic emulsions are used as bases for skincare products for healthy and diseased skin [4].

Botanical extracts that support the health, texture, and integrity of the skin, hair, and nails are widely used in cosmetic formulations. They form the largest category of cosmeceutical additives found in the marketplace today due to the rising consumer interest and demand for natural products [5]. Soybean seeds extract has positive research support for its antioxidant, antiproliferative, and anticarcinogenic activities. This extract is used to treat hyperhidrosis in Asian medicine [6]. Topical application of soy has been used to reduce hyperpigmentation, enhance skin elasticity, control oil production, moisturize the skin, and delay hair regrowth. Soy also has the potential to decrease photoaging of the skin and prevent skin cancers through the antioxidant effects of its metabolites [7]. The major components of soy are phospholipids (45 - 60 %), such as phosphatidyl choline, and essential fatty oils (30 - 35 %). The minor components include the most active compounds, such as isoflavones, saponins, essential amino acids, phytosterols, calcium, potassium, iron, and the proteases soybean trypsin inhibitor and Bowman-Birk inhibitor.

The purpose of the present work is to develop a stable formulation of W/O following inclusion of botanical extract containing soybean seeds extract and its effects were analyzed on various parameters related to skin aging.

EXPERIMENTAL

Materials

Cetyl dimethicone copolyol (ABIL EM 90) was purchased from the Franken Chemicals, Germany, n-hexane and paraffin oil were purchased from Merck KGaA Darmstadt Germany and ethanol (95 %) was purchased from the BDH England. Distilled water was prepared in the cosmetics laboratory of the Department of Pharmacy, The Islamia University of Bahawalpur, Pakistan. Soy bean seeds were obtained in May, 2012 from a Hamid Pensaar, Bahawalpur, Pakistan and authenticated by Prof. Mukhtar Ahmad, Cholistan Institute of Desert Plants Studies (CIDS), The Islamia University of Bahawalpur, Pakistan. For future reference, a voucher specimen collected and identified by the CIDS, and specimen was deposited by the Herbarium, The Islamia University of Bahawalpur (voucher no. GM-SD-7-16-27). Rotary evaporator (Eyela, Japan) was used for solvent evaporation. Mexameter MPA 5 (Courage + Khazaka, Germany) used for the determination of skin melanin and skin erythema.

Preparation of extract

Plant material (1 g per sample) was ground to a fine powder in a mill and extracted for 20 min with ethanol: hexane 1:1 (50 ml) solvent mixture under sonication in an ultrasonic bath at ambient temperature. The extract residue was removed by filtration through layers of muslin cloth. The extract was concentrated at rotary evaporator. Concentrated extract was filtered through Whatman No. 1 filter paper and was stored extract in refrigerator.

Evaluation of antioxidant activity of plant extract

1,1-Diphenyl-2-picrylhydrazyl (DPPH) was prepared by taking 1 mg of DPPH in 25 ml methanol to make 100 μ l solution. The stable DPPH radical was used for the determination of antioxidant activity. Different concentrations of extract in respective solvents were added at an equal volume of 10 μ l to 90 μ l of 100 μ M methanolic DPPH solution in a total volume of 100 μ l in 96 well plates. The contents were mixed and incubated at 37 °C for 30 min. Ascorbic acid was used as standard antioxidant. The reduction in absorbance was measured at 517 nm by using microplate reader; in comparison with the control solution (maximum absorption). The decrease in absorbance indicates increased scavenging activity. The DPPH radical scavenging activity is mentioned as percentage [7].

Preparation of cosmetic emulsions

Soybean seeds (1 g per sample) were ground to a fine powder in a mill and extracted for 48 h with ethanol: hexane 1:1 (50 ml) solvent mixture under sonication in an ultrasonic bath at ambient temperature. The extract residue was removed by filtration through layers of muslin cloth. The extract was concentrated at rotary evaporator. Concentrated extract was filtered through Whatman No. 1 filter paper and was stored in refrigerator. For the formulation oily phase composed of paraffin oil (16 %) and emulsifier Abil-EM90 (3 %) were heated together up to 75 \pm 1 °C. At the same time distilled water (quantity sufficient to make 100 %) was heated at the same temperature and then 4 % soy bean extract was added in it. Next, aqueous phase was added to the oil phase drop by drop. Stirring was continued at 2000 rpm by the mechanical mixer for about 15 min until complete aqueous phase was added, 2 to 3 drops of rose oil were added during this stirring time to give good fragrance to the formulation.

Thereafter, the speed of the mixer was reduced to 1000 rpm for homogenization, for a period of 5 min, and then the speed of the mixer was further reduced to 500 rpm for 5 min for complete homogenization; until the emulsion cooled to room temperature. The base was also prepared by the same method and with same ingredients but without the soy bean seeds extract. Both base and the test formulation were found stable after evaluating pH, centrifugation, electrical conductivity, phase separation, organoleptic and physical characteristics (color, creaming and liquefaction) and the temperature stability tests at storage conditions of 8 ± 0.5 , 25 ± 0.5 , 40 ± 0.5 , and also at 40 ± 0.5 °C/70 % RH (relative humidity) for a period of 3 months.

Ethical approval

This work was authorized by the Board of Advanced Studies and Research (BASR), and its Ethical Committee for *in vivo* Studies (reference no. 4710/Acad.), The Islamia University of Bahawalpur, Pakistan and was directed according to the Principles of Good Clinical Practice [8] and the International Guidelines of Helsinki Declaration [9].

Patch test

To determine the skin irritation assessment, on the first day of study Patch tests were performed on forearms of each human volunteer. An area of 5 cm × 4 cm was marked on both the forearms of all the volunteers. Basic values for erythema and melanin were measured with the help of Mexameter. 1.0 g of base and formulation each were applied to the 5 cm × 4 cm marked regions separately on each forearm. Surgical dressing was used to cover the marked area of right and left forearm. After 48 hours application of emulsion, it was removed and the forearms were washed with physiological saline and were observed for any skin redness/irritation by a dermatologist. The measurements of erythema and melanin were repeated on both forearms.

Study protocol

Eleven volunteers were selected whose ages were in between 25 and 35 years. Male volunteers were included in this work and consent forms were taken. Prior to the tests, a dermatologist for any serious skin disease or damage especially on cheeks and forearms examined the volunteers. Each volunteer was provided with the two emulsions. One emulsion was the base (B) with no extract and other was the active formulation (F) containing soybean extract. The study was designed single blinded

placebo-controlled for the comparisons of two emulsions. The investigations were achieved on cheeks of volunteers. They were also given instructions regarding way of application. The volunteers were advised to use an amount which can be absorbed easily on the skin. Each volunteer applied emulsion for the period of 12 weeks. Values for different parameters were taken in controlled room temperature 25 ± 1 °C and 45 ± 2 % relative humidity. The site of measurement was the whole cheek. Every volunteer was instructed to come for measurement on 2nd, 4th, 6th, 8th, 10th and 12th week. Values for each parameter was measured three times (n=3) and average of the values has been noted.

Anti-aging study

The evaluation of skin microrelief parameters was done by using Visio Scan® VC98 (Courage and Khazaka, Germany). The skin elasticity was measured by an Elastometer (Skin Elastometer®EM 25, Courage and Khazaka, Germany). A skin capacitance meter (Corneometer® MPA 5, Courage and Khazaka, Germany) was used for the determination of skin moisture contents.

Data analysis

The evaluation of data was done by using SPSS version 17.0 according to two-way ANOVA for assessing variation between different time intervals defining 5 % level of significance and Paired samples t-test for determining variation between the two preparations. Standard error of means (SEM) was calculated for every mean value.

Percent change in individual values for each parameter, taken weekly, were computed as in Eq 1.

$$\% \text{ Change} = \{(A - B) / B\} * 100 \dots\dots\dots (1)$$

where A = individual value of any parameter on 2nd, 4th, 6th, 8th, 10th and 12th week, and B = the value of the parameter at 0 h, i.e., at commencement of the test.

RESULTS

For eleven volunteers, changes in skin moisture contents are shown in Figure 1. In this work base enhances the moisture contents of skin to some degree but there was consistent increase in skin moisture contents when formulation was applied throughout the study duration. Insignificant effects ($p > 0.05$) were observed by the base while the formulation presented significant ($p \leq$




Figure 1: Change in (a) skin moisture and (b) elastic content after application of base (◆) and soybean extract formulation (□)

Table 1: SELS values (mean \pm SD) of formulations

Parameter	Emulsion	0 h	1 month	2 months	3 months
SEr	Base	4.05 \pm 0.09	4.04 \pm 0.07	4.04 \pm 0.09	4.04 \pm 0.09
	Formulation	4.04 \pm 0.09	4.00 \pm 0.08	3.95 \pm 0.08	3.82 \pm 0.08
SEsc	Base	1.74 \pm 0.05	1.74 \pm 0.05	1.73 \pm 0.05	1.73 \pm 0.05
	Formulation	1.73 \pm 0.05	1.70 \pm 0.05	1.68 \pm 0.05	1.66 \pm 0.06
SEsm	Base	108.99 \pm 4.91	109.98 \pm 4.91	108.64 \pm 4.89	107.82 \pm 4.92
	Formulation	109.01 \pm 4.77	108.72 \pm 4.81	106.72 \pm 4.66	102.03 \pm 4.23
SEw	Base	71.84 \pm 1.54	72.83 \pm 1.55	71.84 \pm 1.54	70.93 \pm 2.00
	Formulation	71.74 \pm 1.52	71.93 \pm 1.53	69.96 \pm 1.65	68.51 \pm 1.64

Note: SEr = skin roughness; SEw = skin wrinkles; SEsm = skin smoothness; SEsc = skin scaliness

0.05) effects on skin moisture contents regarding with time. Throughout the study duration significant differences in moisture values were evaluated after application of formulation. The irregular pattern in the values of skin elasticity was observed by the base but there was regular increase in skin elasticity after the application of formulation throughout the study duration shown in Figure 2. Test formulation showed significant ($p < 0.05$) while the base had insignificant ($p > 0.05$) effects on skin elasticity. Significant ($p < 0.05$) differences in skin elasticity values were noticed after application of the formulation throughout the study duration. Data about SELS parameters SEw, SEsc, SEsm and SEr before the application of emulsions and then at 1st, 2nd and 3rd months of the study duration are given in Table 1. Insignificant ($p > 0.05$) effect was produced by the base on the SEr parameter of skin and the statistically insignificant ($p > 0.05$) effect was produced by formulation at all reading intervals. Significant ($p > 0.05$) effects were observed for formulation when paired sample t-test was performed. It was observed that active formulation showed a consistent decrease in

values of smoothness, scaliness, roughness and wrinkles.

DISCUSSION

Moisturizing remedy involves renovating the skin barrier, restoring the water contents, ability to retain and redistribute water and sustaining the integrity and skin complexion [10]. Significant differences in moisture values were evaluated after application of formulation throughout the study duration. Enhancement of skin moisture contents after the application of formulation may be because of soy bean extract which contain flavonoids. Soybean flavonoids enhance the skin moisture contents due to swelling of corneocytes on skin surface [11]. Consequently, it may be used in moisturizing cosmetic formulations and also as a complement in the treatment of dry skin.

The elasticity of the skin is mainly due to the collagen fibers which is a main constituent of the dermis of the skin. It constitutes about 70 % of the total dry weight of the skin. Vitamin C stimulates the collagen production in the skin fibroblasts and its deficiency will result in harmful

effects on the skin [12]. The collagen is vulnerable to the UV radiations of the sunlight which causes structural changes in them resulting in photo aging. Photo aging is due to the oxidative effects of these radiations on the collagen of the skin [13]. The improvement in skin elasticity after application of formulation is due to the presence phytoestrogens [14]. Soybean extracts was found to instigate the synthesis of collagen and elastin, and to boost the correct assembly of new elastin fibers, providing a complete protection and restoration to the dermal extracellular matrix [15]. Soybeans extract also protected elastic fibers produced by cultured fibroblasts from degradation by exogenously-added elastases [16].

Skin surface is illustrated by several parameters like smoothness (SEsm), wrinkling (SEw), roughness (SEr) and scaliness (SEsc). Skin surface is assessed both qualitatively and quantitatively by the use of SELS parameters [17]. The proportion of dark pixels is calculated by roughness parameter SEr. A higher SEr value reflects more roughness while a lower value SEr represents more smoothness. SEsm is the smoothness parameter which measures the mean depth and width of wrinkles. SEsc shows the level of dryness of the skin. Stratum corneum dehydration is evaluated by SEsc. The smaller value of SEsc indicates more hydration level with less scaliness [18]. The width and number of skin wrinkles are assessed by SEw which is evaluated by proportion of vertical and horizontal wrinkles. A higher value of SEw proves more and wider wrinkles [17]. It was observed that active formulation showed a consistent decrease in values of smoothness, scaliness, roughness and wrinkles.

The decrease in the average values of skin smoothness in contrast to skin roughness represents that formulation acquires the antiaging effects. Lower value of SEsc designates that there was high skin moisture content. This indicates that formulation has the capability to increase the skin moisture. The smaller values of SEw shown by the extract formulation represent a reduction in fine wrinkles due to the enhancement of skin complexion by stimulation of elastin and collagen synthesis and promotion of production of crosslinked, stable elastin fibers by soybean extract. Augmentation of skin surface parameters can be ascribed to the presence of a rich content of antioxidant flavonoids, known as genistein and daidzein, in soybeans. These substances have been classified as phytoestrogens, since they are plant derivatives with chemical structure similar to human estrogen. Topical estrogens have been

shown to function as antiaging cosmeceuticals by increasing skin thickness, promoting the collagen synthesis and prevention of skin from oxidative damage and photoaging [18].

CONCLUSION

The findings indicate that topical application of cosmetic water-in-oil emulsion incorporating 4 % soybean extract exerts potential antiaging effects. Furthermore, the formulation has no detrimental effects and thus can be used safely. It is also a suitable alternative to most commercial formulations that utilize synthetic bases. The non-invasive technique used in this study is a suitable approach for the evaluation of antiaging properties of skin topical agents.

REFERENCES

1. Gilchrest BA. *Skin aging and photoaging: an overview.* *J Am Acad Dermatol* 1989; 21: 610-613.
2. Lin CH, Sheu JY, Wu HL, Huang YL. *Determination of hydroquinone in cosmetic emulsion using microdialysis sampling coupled with high-performance liquid chromatography.* *J Pharm Biomed Anal* 2005; 38: 414-419.
3. Saikia AP, Ryakala VK, Sharma P, Goswami P, Bora U. *Ethnobotany of medicinal plants used by Assamese people for various skin ailments and cosmetics.* *J Ethnopharmacol* 2006; 106: 149-157.
4. Hall WL, Vafeiadou K, Hallund J, Bügel S, Koebnick C, Reimann M, Ferrari M, Branca F, Talbot D, Dadd T. *Soy-isoflavone-enriched foods and inflammatory biomarkers of cardiovascular disease risk in postmenopausal women: interactions with genotype and equol production.* *Am J Clin Nutr* 2005; 82: 1260-1268.
5. Cassidy A, Albertazzi P, Nielsen IL, Hall W, Williamson G, Tetens I, Atkins S, Cross H, Manios Y, Wolk A. *Critical review of health effects of soybean phyto-oestrogens in post-menopausal women.* *P Nutr Soc* 2006; 65: 76-92.
6. Akiyama T, Ishida J, Nakagawa S, Ogawara H, Watanabe S-i, Itoh N, Shibuya M, Fukami Y. *Genistein, a specific inhibitor of tyrosine-specific protein kinases.* *J Biol Chem* 1987; 262: 5592-5595.
7. Waqas MK, Shah PA, Akhtar N, Murtaza G. *Screening of Various Botanical Extracts for Antioxidant Activity Using DPPH Free Radical Method.* *Afr J Tradit Complement Altern Med* 2013; 10: 452-455.
8. *The European Agency for the Evaluation of Medicinal Products, Human Medicines Evaluation Unit, International Conference on Harmonisation: Guideline for Good Clinical Practice [EMEAWeb site], <http://www.emea.eu.int>.*
9. *World Medical Association Declaration of Helsinki: Ethical principles of medical research involving human*

- subjects. <http://www.wma.net/en/30publications/10policies/b3/index.html>.
10. Grahame R, Holt P. The influence of ageing on the in vivo elasticity of human skin. *Gerontol* 1969; 15: 121-139.
 11. Farage M, Miller K, Elsner P, Maibach H. Intrinsic and extrinsic factors in skin ageing: a review. *Int J Cosmetic Sci* 2008; 30: 87-95.
 12. Quella SK, Loprinzi CL, Barton DL, Knost JA, Sloan JA, LaVasseur BI, Swan D, Krupp KR, Miller KD, Novotny PJ. Evaluation of soy phytoestrogens for the treatment of hot flashes in breast cancer survivors: A North Central Cancer Treatment Group Trial. *J Clin Oncol* 2000; 18: 1068-1068.
 13. Draelos ZD. Topical and oral estrogens revisited for antiaging purposes. *Fertil Steril*. 2005; 84: 291-292.
 14. Zhao R, Bruning E, Rossetti D, Starcher B, Seiberg M, Iotsova-Stone V. Extracts from *Glycine max* (soybean) induce elastin synthesis and inhibit elastase activity. *Exp dermatol* 2009; 18: 883-886.
 15. Heinrich U, Tronnier H, Stahl W, Bejot M, Maurette JM. Antioxidant supplements improve parameters related to skin structure in humans. *Skin Pharmacol Phys* 2006; 19: 224-231.
 16. Primavera G, Berardesca E. Clinical and instrumental evaluation of a food supplement in improving skin hydration. *Int J Cosmetic Sci* 2005; 27: 199-204.
 17. Udompataikul M, Sripiroj P, Palungwachira P. An oral nutraceutical containing antioxidants, minerals and glycosaminoglycans improves skin roughness and fine wrinkles. *Int J Cosmetic Sci* 2009; 31: 427-435.
 18. Kim SY, Kim SJ, Lee JY, Kim WG, Park WS, Sim YC, Lee SJ. Protective effects of dietary soy isoflavones against UV-induced skin-aging in hairless mouse model. *J Am Coll Nutr* 2004; 23: 157-162.