

African Research Review

An International Multi-Disciplinary Journal, Ethiopia

Vol. 3 (3), April, 2009

ISSN 1994-9057 (Print)

ISSN 2070-0083 (Online)

Efficiency in Linear Model with AR (1) and Correlated Error-Regressor (Pp. 46-61)

Olaomi, John Olutunji - Department of Applied Mathematics and Computer Science, Faculty of Applied Sciences, University for Development Studies, Ghana olaomijo@yahoo.com

Adedayo, A. Adepoju - Department of Statistics, University of Ibadan, Nigeria. pojoday@yahoo.com

Abstract

In this study, we conduct several Monte-Carlo experiments to examine the sensitivity of the efficiency of FGLS estimators relative to OLS using the Variance and RMSE criteria, in the presence of first order autocorrelated error terms which are also correlated with geometric regressor. We examine the sensitivity of the efficiency to ρ , α , as well as, its asymptotic behaviour, N , when the above two assumptions are violated. We observe that CORC and HILU give similar result, same for ML and MLGRID. OLS is more efficient than CORC and HILU while ML and MLGRID dominate OLS. In the scenarios, efficiency does not increase with increase in autocorrelation level, only ML and MLGRID at $\alpha = 0.05$ show that efficiency increases with increase in autocorrelation level. All estimators show that efficiency reduces as significant level increases only when the autocorrelation value and sample size are small ($\rho = 0.4$, $N = 20$). There is more efficiency gain when N and ρ are large at all significant correlation levels. Asymptotically, the efficiency of FGLS estimators increase with increasing autocorrelation but it is indifferent to the correlation levels. The asymptotic ranking is CORC and HILU followed by MLGRID and ML.

Keywords: Efficiency, Monte-Carlo Experiment, Feasible Generalised Least Squares, Ordinary Least Squares, Autocorrelation, Significant Correlation.

Introduction

To assess the quality and appropriateness of econometric estimators, we are always interested in their statistical properties. For most estimators, these can only be derived in a "large sample" context, (*asymptotic properties*). One estimation procedure may, for example, be selected over another because it is known to provide consistent and asymptotically efficient parameter estimates under certain stochastic environments. Such a heavy reliance on asymptotic theory can and does lead to serious problems of bias and low levels of inferential accuracy when sample sizes are small and asymptotic formulae poorly represent sampling behaviour. This has been acknowledged in mathematical statistics since the seminar work of R. A. Fisher, who recognised very early the limitations of asymptotic machinery, when he wrote; *"Little experience is sufficient to show that the traditional machinery of statistical processes is wholly unsuited to the needs of practical research. Not only does it take a cannon to shoot a sparrow, but it misses the sparrow! The elaborate mechanism built on the theory of infinitely large samples is not accurate enough for simple laboratory data. Only by systematically tackling small sample problems on their merits does it seem possible to apply accurate tests to practical data"*. [1]

Statisticians are often interested in the relative efficiency of different estimators when the underlying assumptions of least squares breakdown. [2]. Assumptions in the classical normal linear regression model include that of lack of autocorrelation of the error terms and the zero covariance between the independent variable and the error terms.

In this follow up study to the estimation of the parameters of a linear model when the above two least squares assumptions are violated ([3], [4], [5]), we are interested in the relative efficiency of FGLS to OLS in the presence of autocorrelated errors and significant correlation between the independent variable and the error terms. Specifically, we investigate, in a Monte Carlo experiment, the sensitivity of the efficiency of OLS and FGLS estimators in linear model to autocorrelation levels (ρ), significant correlation levels (α) between the autocorrelated error terms and the regressor, as well as, the asymptotic behaviour of efficiency using the Variance and RMSE criteria. It

is known that in linear model with autocorrelated error terms which is independent from the regressor, the feasible generalized least squares (FGLS) estimators usually outshine its ordinary least squares (OLS) counterpart in terms of efficiency. ([6], [7], [8]).

Ordinary regression analysis is based on several statistical assumptions. One key assumption is that the errors are independent of each other. However, with time series data, the ordinary regression residuals usually are correlated over time. (This is known as autocorrelation). It is not desirable to use ordinary regression analysis for time series data since the assumptions on which the classical linear regression model is based will usually be violated.

These violations, seen in widespread applications in operations research, like in queuing theory and econometrics, where the usual assumption of independent error terms may not be plausible in most cases. Also, when using time-series data on a number of micro-economic units, such as households and service oriented channels, where the stochastic disturbance terms in part reflect variables which are not included explicitly in the model and which may change slowly over time. [7].

Violation of the independent errors assumption has three important consequences for ordinary regression. First, statistical tests of the significance of the parameters and the confidence limits for the predicted values are not correct. Second, the estimates of the regression coefficients are not as efficient as they would be if the autocorrelation were taken into account. Third, since the ordinary regression residuals are not independent, they contain information that can be used to improve the prediction of future values. [9] Examples of situations generating dependency between errors and regressors include: Errors in Variables (Stochastic regressors), Lagged dependent variables and autocorrelation, and Simultaneous equation bias. It is known that in economics, measurement errors may be correlated both with themselves and with the regressors. [10] have shown that the error terms in most current formulations of economic relations are highly positively autocorrelated. [6] have shown that there is much to gain and little to lose by considering alternatives to the independent error assumption of the classical linear regression model.

Many models with autocorrelated error terms and dependency between regressors and error terms have been discussed in the literature. These

include [10], [11], [6], [12], [13,14], [15], [16], [7], [17], [18], [19], [8], [5], [4], and [2]. Tests for detecting the presence of autocorrelation and alternative consistent methods of estimating linear models with autocorrelated disturbance terms and significant correlation between regressors and autocorrelated errors have been proposed. For instance, [12] derived a “full” maximum likelihood method approach to estimation of relationships with autocorrelated disturbances. They had a Monte Carlo study of their maximum likelihood estimator and the Cochrane-Orcutt procedure. The model used is:

$$\begin{aligned} Y_t &= \beta_1 + \beta_2 X_t + U_t \\ U_t &= \rho U_{t-1} + e_t \\ e_t &\sim NID(0, 0.0036) \end{aligned}$$

and the independent variables were chosen to contain a large trend component, as realization of

$$X_t = \exp(0.04t) + w_t; \quad w_t \sim NID(0, 0.0009)$$

They varied their sample sizes from 20 to 50 in 200 replications each and three different values of ρ , which are 0.6, 0.8 and 0.99. On each replication, both the conventional and full maximum likelihood estimates were computed for a given realization of the e 's, using the Cochrane-Orcutt procedure and the full maximum likelihood estimates procedure. Their findings using Root Mean Square Error (RMSE) is that the full maximum likelihood estimator is very much better than the Cochrane-Orcutt in estimating β_2 and they are often dramatic in estimating β_1 when the X 's are trended as well as for the ρ , the gains are quite small. It was also found out that the full maximum likelihood estimates of β_1 and β_2 always does better than the conventional ones.

However, in spite of these tests and estimation methods, a number of questions in connection with the estimation of the classical linear model with autocorrelation error terms and non-zero covariance between the independent variable and the error terms remained unanswered. These include the most appropriate estimation method and their efficiencies in the above named specification of the independent variable, the effect of the degree of correlation of the disturbance term, the effect of the degree of correlation of

independent variable and the error terms, the asymptotic effect and the sampling properties of the various estimation methods. [3] has shown that the replication only gives stability to the parameter estimates. The answers to most of these questions would allow for correct inferences to be made in linear models plagued by the scenario depicted above. It would also relieve the empirical worker from the reliance placed on asymptotic theory in estimation and inference.

The rest of this paper discusses the model and the experimental framework in section 2, Section 3 presents the simulation results, and section 4 presents the discussions, while we conclude in section 5.

The Model

We assume a simple linear regression model:

$$Y_t = \beta_0 + \beta_1 X_t + U_t \quad \text{-----} \quad (1)$$

$$U_t = \rho U_{t-1} + \varepsilon_t, \quad |\rho, \lambda| < 1,$$

$$X_t = \lambda X_{t-1}, \lambda=0.8, X_0 = 515 \quad U_t \rightarrow N\left(0, \frac{\sigma^2}{1-\rho^2}\right),$$

$$X_t \rightarrow N\left(0, \frac{\sigma^2}{1-\lambda^2}\right), \quad t = 1, 2, \dots N., \quad \square = (1,1)$$

where Y_t is the dependent variable and the first order autoregressive X_t is the independent variable with U_t also autoregressive of order one. ε_t is normally distributed with zero mean and variance σ^2 . ρ and λ are stationarity parameters while the model parameters are assumed to be unity. This independent variable specification had been used by [7], [5], and [20]. It is chosen to allow for comparison of results.

Experimental Framework

We used the Monte-Carlo approach for the investigation due to the fact that when the covariance between the independent variable and the autocorrelated error terms is non-zero, the problem is near intractable by analytical procedure. Small sample investigations are also usually made using the Monte Carlo method. Also the properties of FGLS estimators vary depending

on the form of the variance – covariance matrix, and often the quality of this variance – covariance matrix can not be neatly summarized. Many estimation methods of our model have been developed over the years. Because of the least squares violations in the model, the FGLS estimators are considered relative to the OLS estimator. Some of the FGLS estimators in literatures include the Beach and MacKinnon Maximum Likelihood, Maximum Likelihood Grid, Cochran Orcutt, Durbin, Prais Winstein and Hildreth Lu. The various methods of parameter estimation in linear models with autocorrelated disturbances have known asymptotic properties. [16] while their sampling properties are yet to be well investigated and understood. This corroborates [21] when he asserts that “*The elaborate mechanism built on the theory of infinitely large samples is not accurate enough for simple laboratory data. Only by systematically tackling small sample problems on their merits does it seem possible to apply accurate tests to practical data*”.

Most of the existing estimation methods possess desirable properties; however, the autocorrelation and the significant dependency of independent variable and the error terms, in addition to the specification of the independent variable might affect these properties. Since Monte-Carlo experiments provide a means of modelling small sample properties of estimators, it is used here to study these properties.

The following four FGLS estimators: Cochrane and Orcutt (CORC), Hildreth and Lu (HILU), Maximum Likelihood (ML) and Maximum Likelihood Grid (MLGRID) and Ordinary Least Squares (OLS) estimation methods, choosing in the light of the previous works, are used. These estimators are equivalent with identical asymptotic properties. ([17], [18], [19]). But in small samples, such as in this study, [22] have argued that those that use the T transformation matrix (ML, MLGRID) are generally more efficient than those that use T* transformation matrix (CORC, HILU). (See [5])

The degree of autocorrelation affects the efficiency of the estimators [7]. Consequently, we investigated the sensitivity of the estimators to the degree of autocorrelation by varying $\hat{\rho}$ from 0.4, to 0.8 and 0.9. We also found out the effect of the correlation of the independent variable and the error terms at significant level 1%, 2% and 5% on the estimators. The effects of sample size was also investigated by varying the sample size (N) from 20, 40

to 60 each replicated 50 times. Evaluation of the estimators was done using the Relative Efficiency based on Variance and the RMSE criteria.

A total of 27 data sets spread over three sample sizes were used in generating the data for this study. Using model (1), a value U_o (for specified sample size) was generated by drawing a random value \square_o from $N(0,1)$ and dividing by $\sqrt{(1-\rho^2)}$.

Successive values of \square_t drawn from $N(0,1)$ were used to calculate U_t . X_t was similarly generated. The correlation coefficient between U_t and X_t was then computed and its absolute value tested for significance at, say 1%. If this value is significant, it is chosen; otherwise it is discarded. This procedure is repeated as many times as are necessary (for all \square , α and N) to obtain fifty replications for a desired sample size. Y_t is thus computed for the chosen U_t and fixed geometric trended X_t using the model. The data generations are made using the Excel package while estimations are done via the AR procedure of [23]. Estimation result for this scenario of the independent variable is presented in [5].

The finite sampling properties of estimators used include the Variance (VAR) and the Root Mean Squared Error (RMSE). Additionally, we calculated the Sum of Variances (SVAR) and the Sum of Root Mean Squared Error (SRMSE). These are further used to compute the Relative efficiency. The relative efficiency of the FGLS estimators relative to OLS is:

$$\frac{\text{Var } \hat{\beta}_{(OLS)}}{\text{Var } \hat{\beta}_{(GLS)}} \text{ or } \frac{\text{MSE } \hat{\beta}_{(OLS)}}{\text{MSE } \hat{\beta}_{(GLS)}}$$

Then to get the Total Gain or Loss (G/L), we subtract 1 (original estimate) from the efficiency of each coefficient and add our results. That is, efficiency

gain or loss is $(\hat{\beta}_o - \beta_o) + (\hat{\beta}_1 - \beta_1)$, where $\hat{\beta}(\cdot)$ represents the efficiency of $\hat{\beta}$. If the relative efficiency is negative, then OLS is more efficient. The results for each scenario, using both Variance and RMSE criteria, are summarised in Tables 1 and 2 for RMSE and VAR criteria respectively.

Simulation Results

Perusing Tables 1 and 2, it is observed that in all the scenarios considered in the experiment, CORC and HILU efficiencies are similar and same for ML and MLGRID. OLS is more efficient than CORC and HILU in many of the scenarios while ML and MLGRID are more efficient than OLS. In majority of the scenarios, efficiency does not increase with increase in autocorrelation level, only ML and MLGRID at $\alpha = 0.05$ show that efficiency increases with increase in autocorrelation level. All estimators show that efficiency reduces as significant level increases only when the autocorrelation value and sample size are very small (that is $\rho = 0.4$, $N = 20$). There is more efficiency gain when N and ρ are large at all significant correlation levels. Table 3 summarises Tables 1 and 2, where we found the best estimator (estimator with the largest efficiency under each of the variance and RMSE criteria). Holding N , α and ρ constant, the ML estimator has the largest efficiency in 44.5% of the scenarios, followed by MLGRID (29.6%), CORC (11.1%), HILU (7.4%) and OLS (7.4%).

In order to bring out the most information from this research, we charted the efficiency levels recorded in Tables 1 and 2 in Table 4 showing the asymptotic, autocorrelation and significant level effects. Table 4 gives the frequency distribution of N – chart over \square and \square for both variance and RMSE-based efficiency measures for all estimators (measuring the asymptotic effect). The chart symbols include (\wedge) indicating, minimum efficiency when $N = 60$, intermediate when $N = 40$ and maximum when $N = 20$. (\vee) Efficiency is a minimum when $N = 40$, and maximum when $N = 20$ or 60. (\square) Efficiency is a maximum when $N = 40$, and minimum when $N = 20$ or 60, and ($/$), means Efficiency is a maximum when $N = 60$, intermediate when $N = 40$ and minimum when $N = 20$.

Table 4.1 shows that the trend ' $/$ ' is the most frequent. This implies that the efficiency is highest when $N = 60$ followed by those at $N = 40$ and is smallest at $N = 20$. This most frequent trend occurs with highest frequency when $\square = 0.9$ (19). Table 4.2 also gives additional interesting information that the significant level does not matter for asymptotic efficiency as the most frequent trend ' $/$ ' appear equally among the significant levels. The last column of Table 4.3 contains a summary of the two ranks of each of the four estimators. These estimators rank as follows in decreasing order of conformity with the observed asymptotic behaviour of efficiencies of variance and RMSE: CORC (3), HILU (3), MLGRID (7), and ML (7). In

conclusion, the above results show that, using our criteria, the efficiency of the FGLS estimators increase asymptotically and the optimum combination of ρ , λ and N is: all ρ , $\lambda = 0.9$, and $N = 60$.

Discussion of the Results

We note that the efficiency of ML and MLGRID have very similar behavioural pattern, the same for CORC and HILU as observed in the finite sampling properties of Variance and the RMSE. ML and MLGRID are better than both CORC and HILU as also observed by [24], [7], and [8].

Asymptotically, the estimators increase asymptotically and the optimum combination of ρ , λ and N is: all ρ , $\lambda = 0.9$, and $N = 60$. This implies that, the efficiency of the FGLS estimators, relative to the OLS estimator, increases asymptotically and with increasing autocorrelation. This is similar to the results obtained by [13] and [17] when the regressor and error terms are independent. The estimators rank as follows in decreasing order in conformity with the observed asymptotic behaviour of efficiency: CORC, HILU, MLGRID, and ML. This also indicates that truly, the nature of the regressor affects the efficiency of FGLS estimators. As if we compare this result with that of [8], there is a disparity as a result of the nature of the regressor. Our results have also shown that there is a definite gain to be obtained from using some of the feasible GLS as they are more efficient than OLS. This also conform to the earlier result by [6] where they show that FGLS are better for given values of $|\rho| \geq 0.3$ when there is independence between the error terms and the regressor.

Conclusion

We have investigated the sensitivity of the significant correlation between the error terms and the geometric regressor in a single linear regression model to the efficiency of the various FGLS estimators relative to that of the OLS estimator. It could be concluded that empirically, the OLS estimator is more efficient than the FGLS estimators CORC and HILU as OLS dominated them almost uniformly. Maximum likelihood estimation methods of MLGRID and ML still perform better than other FGLS estimators in terms of efficiency. All estimators show that efficiency reduces as significant level increases only when the autocorrelation value and sample size are very small (that is $\rho = 0.4$, $N = 20$). Asymptotically, the efficiency of FGLS estimators increase asymptotically with increasing autocorrelation but it is indifferent to the

significant correlation levels between the error terms and the geometric regressor. The asymptotic ranking is CORC and HILU followed by MLGRID and ML.

References

- Philips P. C. B (1982): *Small Sample Distribution Theory In Econometric Models Of Simulateneous Equations*. Cowles Foundation Discussion Paper No 617. Yales University.
- Samir Khaled Safi (2008): *Explicit Equations to Determine the Variances of Regression Coefficients of OLS and GLS Estimators In An AutoCorrelated Regression Models*. The Islamic University Journal (Series of Natural Studies and Engineering) Vol.16, No. 1, pp 65-74
<http://www.iugaza.edu.ps/ara/research/>
- Olaomi, J. O. (2004): *Estimation of Parameters of Linear Regression Models with Autocorrelated Error terms which are also correlated with the regressor*. Unpublished Ph.D. Thesis. University of Ibadan, Nigeria.
- Olaomi, J. O. and Ifederu Adetoro, (2008): “Understanding Estimators of Linear Regression model with AR(1) Error which are correlated with Exponential Regressor” – *Asian Journal of Mathematics and Statistics*, AJMS 1 (1):14-23.
- Olaomi, J. O. (2008): “Understanding Estimators of Linear Regression model with AR(1) Error which are correlated with Geometric Regressor” – *European Journal of Science Research (EJSR)* Vol 20: No 1. pp 213 - 223 www.eurojournals.com/ejsr.html
- Rao, P. and Griliches, Z. (1969) “Small Sample Properties of Several Two-stage Regression Methods in the Context of Autocorrelated Errors”. *Journal of the American Statistical Association*, 64, 251-272.
- Nwabueze, J. C (2000): *Estimation of Parameters of Linear Regression Models with Autocorrelated Error terms*. Unpublished Ph.D. Thesis. University of Ibadan, Nigeria.

- Olaomi, J. O. and Iyaniwura J. O. (2006): Efficiency of GLS estimators in linear regression model with autocorrelated error terms which are also correlated with the regressor. *SCIENCE FOCUS, International Journal of Biological and Physical Sciences, Vol. 11*, pg 129-133.
- SAS (Ibid): Autoreg Procedure, SAS OnlineDoc 7-1, www.okstate.edu/sas/v7/saspdf/ets/chap8.pdf
- Cochrane, D. and Orcutt, G.H. (1949) “Application of Least Regression to Relationships Containing Autocorrelated Error Terms”. *Journal of the American Statistical Association*, 44, 32-61.
- Durbin, J. and Watson, G.S. (1971) “Test for Serial Correlation in Least Squares Regression III”, *Biometrika*, 58, 1-42.
- Beach, C. M. and Mackinnon, J. S. (1978) “A Maximum Likelihood Procedure for Regression with Autocorrelated Errors”. *Econometrica*, 46, No. 1, 51-57.
- Kramer, W. (1980): “Finite Sample Efficiency of Ordinary Least Squares in the Linear Regression Model with Autocorrelated Errors”. *Journal of the American Statistical Association*, 75, 1065-1067.
- Kramer, W. (1998): “Asymptotic Equivalence of Ordinary Least Squares and Generalized Least Squares with Trending Regressors and Stationary Autoregressive Disturbances”. In Galata/Kutchenhoff (eds.): *Econometrics in Theory and Practice (Festschrift for Hans Schneeweis β)*, 137-142.
- Busse, R., Jeske, R. and Kramer, W. (1994): “Efficiency of Least-Squares Estimation of Polynomial Trend when Residuals are Autocorrelated”. *Economics Letters* 45, 267-271.
- Kramer, W. and Hassler, U. (1998): “Limiting Efficiency of OLS Vs. GLS when Regressors are Fractionally Integrated”. *Economic Letters* 60, 285-290.

- Kleiber, Christian (2001): “Finite Sample Efficiency of OLS in Linear Regression Models with Long-Memory Disturbances”. *Economic Letters* 72, 131-136.
- Kramer, W. and Marmol F. (2002): “OLS-based Asymptotic Inference in Linear Regression Models with Trending Regressors and AR(P) Disturbances”. *Communications in Statistics – Theory and Methods*, 31, 2, 2002, 261-270.
- Butte Gotu (2002): “The Equality of OLS and GLS Estimators in the Linear Regression Model when the Disturbances are Spatially Correlated”. *Statistical Papers*. Vol. 42 Issue 2, 253-263.
- Ifederu (2006): Estimation of the Parameters of Single Linear Regression Model with Autocorrelated Error Terms which are also correlated with the trended regressor. Unpublished M. Sc. Thesis. University of Ibadan, Nigeria.
- Fisher, R. A. (1925): *Statistical methods for research workers*, Edinburgh; Oliver and Boyd. (Sited in Philips P. C. B, 1982, *Small Sample Distribution Theory In Econometric Models Of Simultaneous Equations*. Cowles Foundation Discussion Paper No 617. Yales University)
- Park, R.E. and Mitchell, B.M. (1980) “Estimating the Autocorrelated Error Model with Trended Data”. *Journal of Econometrics*, 13, 185-201.
- TSP (2005) *Users Guide and Reference Manual*. Time Series Processor. New York.

Table 1: Efficiency of FGLS to OLS using RMSE

		N=20			N=40			N=60		
Signif icant level	Estim ator		$\rho =$ 0.8	$\rho =$ 0.9	$\rho =$ 0.4	$\rho =$ 0.8	$\rho =$ 0.9	$\rho =$ 0.4	$\rho =$ 0.8	$\rho =$ 0.9
0.01	OLS	0	0	0	0	0	0	0	0	0
	COR C	- 0.111 24	- 0.142 1	- 0.007 48	0.014 264	- 0.062 31	- 0.588 18	0.016 629	0.388 47	0.082 889
	HIL U	- 0.111 79	- 16.34 885	- 0.006 81	0.013 81	- 0.057 44	- 0.006 466	0.019 838	0.380 271	0.097 558
	ML	0.059 012	0.071 676	0.000 41	0.013 227	0.211 346	0.290 823	0.055 108	0.024 961	0.513 185
	MLG RID	0.045 324	0.073 073	0.000 131	0.013 162	0.211 593	0.282 707	0.049 374	0.024 905	0.374 133
0.02	OLS	0	0	0	0	0	0	0	0	0
	COR C	- 0.067 4	- 0.965 71	- 0.044 81	- 0.003 61	0.230 127	0.022 946	- 0.003 57	1.633 73	- 0.074 16
	HIL U	- -0.08	- 0.227 47	- 0.044 59	- 0.003 11	0.234 435	- 0.049 35	- 0.003 15	1.364 32	- 0.066 881
	ML	0.032 743	0.090 743	0.013 34	0.003 276	0.405 858	0.254 624	0.003 706	1.445 87	0.301 885
	MLG RID	0.031 84	0.088 679	0.014 16	0.003 349	0.411 975	0.253 686	0.003 599	1.416 06	0.294 989
0.05	OLS	0	0	0	0	0	0	0	0	0
	COR C	2.627 277	0.805 58	0.662 91	0.131 762	0.081 07	0.467 19	0.003 35	0.903 795	0.082 348
	HIL U	- 0.003 26	- 0.947 66	- 0.788 49	0.134 575	- 0.083 13	- 0.333 54	- 0.003 35	0.897 048	- 0.107 49
	ML	0.020 736	0.128 359	0.183 662	0.129 727	0.150 683	0.258 348	0.009 321	0.052 556	0.572 738
	MLG RID	0.019 87	0.093 119	0.161 984	0.128 458	0.155 675	0.257 41	0.010 177	0.049 336	0.578 2

Table 2: Efficiency of FGLS to OLS using Variance

		N = 20			N = 40			N = 60		
Significant level	Estimator	$\rho = 0.4$	$\rho = 0.8$	$\rho = 0.9$	$\rho = 0.4$	$\rho = 0.8$	$\rho = 0.9$	$\rho = 0.4$	$\rho = 0.8$	$\rho = 0.9$
	OLS	0	0	0	0	0	0	0	0	0
	CORC	0.227 74	0.305 89	0.021 63	0.069 42	0.180 1	0.897 43	0.131 691	0.080 06	0.194 709
	HILU	0.228 74	308.3 316	0.018 58	0.068 406	0.162 39	0.007 34	0.139 32	0.274 79	0.254 439
	ML	0.113 447	0.141 86	0.001 844	0.060 568	0.466 597	0.659 519	0.255 485	0.054 86	0.838 65
	MLGRID	0.081 615	0.142 836	0.004 615	0.060 463	0.464 71	0.622 874	0.230 28	0.030 35	0.812 434
	OLS	0	0	0	0	0	0	0	0	0
	CORC	0.124 4	0.998 74	0.086 29	0.005 53	0.496 41	0.043 627	0.001 59	6.855 209	0.060 34
	HILU	0.150 02	0.374 9	0.085 86	0.004 57	0.513 565	0.025 825	0.000 89	7.258 733	0.066 334
	ML	0.066 228	0.165 031	0.026 52	0.014 949	0.793 466	0.703 581	0.019 858	11.79 51	0.860 854
	MLGRID	0.064 277	0.160 319	0.026 82	0.015 043	0.826 917	0.720 795	0.019 75	16.47 161	0.840 649
	OLS	0	0	0	0	0	0	0	0	0
	CORC	12.17 045	1.291 44	1.043 79	0.094 013	0.229 36	0.802 76	0.007	0.070 414	0.058 36
	HILU	0.007 03	1.438 48	1.193 97	0.105 632	0.233 06	0.566 1	7.31E -05	0.103 32	0.160 05
	ML	0.041 434	0.237 734	0.335 683	0.082 023	0.322 055	0.613 454	0.046 454	0.220 231	0.980 49
	MLGRID	0.039 596	0.219 958	0.344 411	0.076 997	0.354 306	0.602 212	0.048 038	0.185 336	1.015 634

Table 3: Best Estimator of our Model for Each Scenario

Criteria	(a) ₁	N = 20			N = 40			N = 60		
		$\rho=0.4$	$\rho=0.8$	$\rho=0.9$	$\rho=0.4$	$\rho=0.8$	$\rho=0.9$	$\rho=0.4$	$\rho=0.8$	$\rho=0.9$
	0.01	ML	HILU	MLG	CORC	ML	ML	ML	OLS	ML
	0.02	ML	ML	OLS	MLG	MLG	MLG	ML	MLG	ML
	0.05	CORC	ML	MLG	HILU	MLG	ML	MLG	ML	MLG
	0.01	ML	HILU	MLG	CORC	MLG	ML	ML	CORC	ML
	0.02	ML	ML	OLS	MLG	MLG	ML	ML	OLS	ML
	0.05	CORC	ML	ML	HILU	MLG	ML	MLG	CORC	MLG

Table 4: Asymptotic Behaviour of Rmse and Variance of Estimators When Sig. Level (α) Is Constant

α	0.01			0.02			0.05								
α	0.4	0.8	0.9	0.4	0.8	0.9	0.4	0.8	0.9	\	V	α	/		
ESTIMATOR	RMSE														
CORC	/	/	V	/	α	α	\	/	/			1	1	2	5
HILU	/	V	/	α	α	V	α	/	/			0	2	3	4
ML	V	α	/	V	α	/	α	α	/			0	2	4	3
MLGRID	V	α	/	V	α	/	α	α	/			0	2	4	3
														(15)	
ESTIMATOR	VAR														
CORC	/	/	V	/	/	α	\	/	/			1	1	1	6
HILU	/	\	/	/	/	/	α	/	/			1	0	1	7
ML	V	α	/	V	/	/	α	α	/			0	2	3	4
MLGRID	V	α	/	V	/	/	α	α	/			0	2	3	4
														(21)	
C – Summary	0.4	0.8	0.9	0.4	0.8	0.9	0.4	0.8	0.9						
\	0	1	0	0	0	0	0	2	0	0					
V	4	1	2	4	0	1	0	0	0						
α	0	4	0	1	4	2	6	4	0						
/	4	2	6	3	4	5	0	4	8						

Table 4.1 And - Estimator Based Summaries of Table 4

α	$\alpha = 0.4$	$\alpha = 0.8$	$\alpha = 0.9$	TOTAL		CORC	HILU	ML	MLGRID
\	2	1	0	3		2	1	0	0
V	8	1	3	12		2	2	4	4
α	7	12	2	21		3	4	7	7
/	7	10	19	36		11	11	7	7

Table 4.2 □ - And - Estimator Based Summaries of Table 4

	□ = 0.01	□ =0.02	□ = 0.05	TOTAL		CORC	HILU	ML	MLGRID
\	1	0	2	3		2	1	0	0
V	7	5	0	12		2	2	4	4
□	4	7	10	21		3	4	7	7
/	12	12	12	36		11	11	7	7

Table 4.3: Summary of the Ranking of Estimators

	Optimum Trend		
	VAR (I)	RMSE (I)	
CORC	(6) 2	(5) 1	3
HILU	(7) 1	(4) 2	3
ML	(4) 3.5	(3) 3.5	7
MLGRID	(4) 3.5	(3) 3.5	7

LEGEND

- \: Efficiency is a minimum when N = 60, intermediate when N = 40 and maximum when N = 20.
- V: Efficiency is a minimum when N = 40, and maximum when N = 20 or 60
- : Efficiency is a maximum when N = 40, and minimum when N = 20 or 60
- /: Efficiency is a maximum when N = 60, intermediate when N = 40 and minimum when N = 20.