

The scourge of head injury among commercial motorcycle riders in Kampala; a preventable clinical and public health menace

Louis H Kamulegeya¹, Mark Kizito¹, Rosemary Nassali², Shiela Bagayana³, Alex E Elobu¹

1. Department of Surgery, Makerere University College of Health Sciences
2. Department of Nursing, Makerere University College of Health Sciences
3. Departement of Biomedical Engineering, Makerere University College of Health Sciences

Abstract

Background: Trauma is an increasingly important cause of disease globally. Half of this trauma is from road traffic injuries with motorcycles contributing 21-58%. Low protective gear use, lack of regulation and weak traffic law enforcement contribute to unsafe nature of commercial motorcycles also known as “boda boda” in Uganda.

Objectives: To determine the prevalence of protective gear use, the occurrence of head injury and the relationship between the two among commercial motorcycle riders in Kampala.

Methods: Following ethical approval we recruited consecutive consenting participants to this analytical cross-sectional study. Data was collected using pretested interviewer administered questionnaires, double entered in Epidata and analyzed with STATA. Proportions and means were used to summarize data. Odds ratios were calculated for association between wearing helmets and occurrence and severity of head injury.

Results: All 328 participants recruited were male. Of these, 18.6% used Protective gear and 71.1 % sustained head injury. Helmets protected users from head injury (OR 0.43, 95% CI, 0.23-0.8) and significantly reduced its severity when it occurred.

Conclusion: Protective gear use was low, with high occurrence of head injury among commercial motorcycle riders in Uganda. More effective strategies are needed to promote protective gear use among Uganda’s commercial motorcycle riders.

Keywords: Boda-boda, okada, commercial motorcycle, head injury, protective gear, helmet, Mulago hospital.

DOI: <http://dx.doi.org/10.4314/ahs.v15i3.41>

Cite as: Kamulegeya LH, Kizito M, Nassali R, Bagayana S, Elobu AE. The scourge of head injury among commercial motorcycle riders in Kampala; a preventable clinical and public health menace. *Afri Health Sci.* 2015;15(3):1016-22. doi: <http://dx.doi.org/10.4314/ahs.v15i3.41>

Introduction

Road traffic injuries are the leading cause of injury-related deaths and disability worldwide but more so in low and middle income countries¹⁻³. Injuries are increasingly becoming an important contributor to disability adjusted life years worldwide^{2,4}

Trauma is the fifth or sixth leading cause of disease burden in Uganda⁵ with road traffic crashes topping the causes of injury in Uganda at 48%⁶.

In East Africa, motorcycle related crashes contribute

21- 58.8% of the road traffic crashes with head injury being common⁷⁻¹⁰. Most of the crash victims require surgical intervention and survivors tend to have chronic disabilities⁸.

The commercial motorcycles, also known as “Boda-boda” in East Africa or “Okada” in West Africa as previously described are popular in both urban and rural setting as a quick means of transport⁸⁻¹⁰. With over 100,000 commercial motorcycles in Kampala alone⁸, this lucrative industry generates more than US\$ 400,000 daily. Attempts by institutions e.g. the Uganda Police Force, Kampala Capital City Authority to regulate the industry have been received with mixed feelings, resistance by operators, political interference and have not yet been successful^{8,11}. Observed crash helmet use is low at 30.5 % and 1% among riders and passengers respectively¹². Efforts to promote road safety especially protective gear use have been curtailed by underfunding, weak implementation and erratic traffic law enforcement. The Uganda Police Force which is constitutionally mandated to enforce such law reports lack of human and non-human resources and inadequate police training to meet current and future challenges of traffic policing.^{7-8,10,13}

Corresponding author:

Alex E Elobu
Makerere University
College of Health Sciences,
Department of Surgery,
P.O.BOX 7072, Kampala,
Fax: +256-414532591,
Tel: +256-782714892,
Email: elobuemmy@yahoo.co.uk

Despite the burden of the problem in Uganda, motorcycle injuries have not received the attention they deserve. There is need for more literature on these injuries that will inform stakeholders on the magnitude of the problem¹⁴. Since the majority of motorcycle injuries are preventable, a clearer understanding of the magnitude, contributing factors and injury patterns is essential for establishment of prevention strategies as well as treatment protocols^{7,14}.

This study documents the prevalence of head injury and use of protective gear among motorcycle accident victims seen at Mulago hospital and makes a case for the use of protective gear to prevent motorcycle related injuries or mitigate their severity.

Methods

Following ethical approval from Mulago hospital Institutional Review Board an analytical cross sectional study was conducted in 2013 at the Accidents and Emergency Department of Mulago Hospital. Mulago hospital is a 1,500 bed national referral hospital and the teaching hospital for Makerere University. On average, 10 to 20 motorcycle crush victims are attended to daily. About 65% of Mulago Hospital trauma victims come from Kampala city and surrounding areas¹⁴.

Consecutive consenting participants were prospectively enrolled. Participants for whom it was impossible to determine if they were wearing protective gear at the time of the crush were excluded. The researchers participated in the initial evaluation and resuscitation of the patients. Participant biodata, injury history and neurological examination findings were collected using pre-tested interviewer administered questionnaires, entered into Epidata Entry and analyzed with STATA 12. Head injury severity was measured using the Glasgow Comma Scale. Data were tabulated. Proportions and means were used to summarize the data. Chi square and Bartlett's tests were used to compare the head injury and no head injury groups. Odds ratio was calculated for the relationship between protective gear use and occurrence and severity of head injury.

Results

Of the 338 participants recruited, questionnaires were completed for 328 (97%). Ten participants were excluded because it was impossible to determine if they were wearing any protective gear at the time of injury.

Table 1: Socio-demographic characteristics of the study population

Characteristic	Head Injury present	No head Injury	Total(328) n (%)	Test statistic	p-value
Sex					
Male	233 (71)	95 (29)	328(100%)		
Age in years (Mean, SD [95%CI])	30.29 (9.6) 28.32-37.27	28.28 (8.9) 27.33-29.63	29.01(9.15) 28.01-30	0.963*	0.326
Ethnicity					
Bantu	212 (71.1)	86(28.9)	298(90.9%)	3.250 [†]	0.355
Nilotics	17 (73.9)	6(26.1)	23(7.0%)		
Nilo-Hamites	2(100)	0(0)	2(0.6%)		
Sudanese	2 (40)	3 (60)	5(1.5%)		
Level of Formal Education					
None	39(81.3)	9(18.8)	48(14.6%)	11.112 [†]	0.025
Primary	104(75.4)	34(24.6)	138(42.1%)		
Secondary	72 (62.1)	44(37.9)	116(35.4%)		
Vocational training	5(100)	0	5(1.5%)		
Tertiary	13(61.9)	8(38.1)	21(6.4%)		

* Bartlett's test for homogeneity of variance

[†] Chi square test

The mean age was 29.01 years (SD=9.15), and all participants were male. The majority (90%) were of Bantu ethnicity and slightly over half (56.7%) had primary level or no formal education at all.

Table 2: Comparative analysis of the study arms

Characteristic	Head Injury	No head Injury	Total(328) n (%)	Test statistic	p- value
Mechanism of Injury					
Motor-cycle Vs Motor-cycle	56(82.4)	12(7.6)	68(20.7%)	5.499 [‡]	0.139
Motor-cycle Vs other vehicle	123 (68)	58(32)	181(55.2%)		
Motor-cycle Vs pedestrian	15(65.2)	8(4.8)	23(7.0%)		
Other collisions	39 (69.6)	17 (30.4)	56(17.1%)		
No of People on the Boda-boda					
1	113(77.9)	32(22.1)	145(44.2%)	10.295 [‡]	0.036
2	97 (68.3)	45 (31.7)	142(43.3%)		
3	20(54.1)	17 (45.9)	37(11.3%)		
4	2(100)	0	2(0.6%)		
5	1(50)	1 (50.1)	2(0.6%)		
Pre hospital transport					
Private vehicle	70(64.2)	39(35.8)	109 (33.2)	30.306 [‡]	<0.001
Police vehicle*	108(81.2)	25(8.8)	133(40.5)		
Police ambulance	4(100)	0	4(1.2)		
Ambulance	30(85.7)	5(4.3)	35(10.7)		
Motorcycle	21(44.7)	26(55.3)	47(14.3)		
Time from Injury to hospital in minutes	619.2 (1338.01)	317.42 (841.43)	531.66 (1221.66)	24.637 [†]	<0.001
(Mean, SD [95%CI])		446.3-791.72	398.96-664.36		

* These are non-ambulance vehicles for other police duties e.g. highway patrol pickup trucks

[‡] Chi square test

[†] Bartlett's test for homogeneity of variance

In half (55.2%) of the times, crashes were between a motor cycle and a vehicle. On most occasions (87.5%) there were one or two people on the motorcycle. Nearly three quarters (73.7%) of the crash victim reached

hospital by means of a private car or police vehicle (not ambulance). Notably, only 11.9% were brought by ambulance. On average it took up to 531 minutes (8 hours and 55 minutes) for injured victims to reach the hospital.

Table 3: Prevalence of Head injury with wearing of protective gear

Protective gear	Head injury present		
	YES n (%)	NO n (%)	TOTAL n (%)
YES	34(55.7%)	27 (44.3%)	61 (18.6%)
NO	199(74.4%)	68 (25.5%)	267 (81.4%)
TOTAL	233 (71%)	95 (29%)	328 (100%)

Of the 328 participants recruited, only 61 (18.6%), wore protective gear and 233 (71%) sustained head injury. Wearing of protective gear was protective of the pre-

venting head injury with a statistically significant odds ratio of 0.43 (p= 95% p value= 0.004, CI 0.23-0.8). Conversely, not wearing head gear was associated with a 2.3 fold increase in odds of sustaining head injury.

Table 4: Relating the severity of head injury and wearing of protective gear

Protective gear	Severity of Head Injury			
	Severe N (%)	Moderate N (%)	Mild N (%)	Total
NO	23(92%)	95(92.2%)	81(77.1%)	199
YES	2 (8%)	8(7.77%)	24(22.9%)	34
TOTAL	25(10.7%)	103(44.2%)	105(45.1%)	233

Only 14.6% of the participants who sustained head injury reported wearing any protective gear at the time of the crash. Also, the proportion of the participants with severe head injury who were not wearing protective gear (92%) was statistically significantly (p value = 0.005) higher than those with mild head injury who also were not wearing protective gear (77.1%) This shows

that wearing of protective gear reduced the severity of head injury among crash victims.

Discussion

This study revealed a very low prevalence of protective gear use and an alarmingly high occurrence of head injury, findings that are of great clinical and public health importance.

All the study participants were male, with average age of 29 years. The majority (90.9%) was of Bantu ethnicity and over three quarters (77%) had no formal or primary level education. This participant population generally compares to that of other studies in other low and middle income countries^{7,15-16}.

In this study only 18.6% of the participants reported wearing crash helmets at the time of the road crash. This is somewhat higher than the 13% reported in 2011 in the same hospital¹². It compares to the other reported prevalence rates of 17.1-24% in Uganda and Tanzania⁷⁻⁸ but is higher than that reported in Nigeria at only 8%¹⁷. This low prevalence arises from the observed low rate of protective gear use of 30.5 % among riders in Kampala¹².

Even if over estimated, the 18.1% rate of protective gear use is unacceptably low indicating that our current efforts to promote the use of protective gear are largely inadequate and have had very limited positive impact. There was an extremely high proportion of head injury at 71 % of all motorcyclists. The reported proportion of head injury was 10.1% in 2001⁹, 35.5% in 2009⁸ and 71.1% in 2013 in this study. This alarming development could be explained by the persistent low use of protective gear and increased rider recklessness at a time when the commercial motorcycles being used are more powerful¹⁸ and capable of higher energy transfers leading to more severe injuries.

This study was restricted to motorcycle riders. Pedestrians may not undergo similar energy transfers as riders and may be less likely to sustain head injury. This may explain the rather high head injury proportion in this study that was limited to riders.

In this study, helmet use was protective of head injury with an Odds ratio of 0.43 (95% CI 0.23-0.8). Conversely, not wearing protective gear increased the odds of sustaining head injury by 2.3 fold. This compares to previous studies in which the OR was from 2.5 to 4.4¹⁹⁻²⁰. Also there was a trend towards more severe forms of head injury for motorcyclists who were not wearing protective gear.

Several systematic reviews recommend helmets for preventing crash fatalities and head injury²¹⁻²². At a provincial hospital in Kenya, all trauma fatalities had head injury among other injuries¹⁵. In Tanzania, none of the crash victims who wore helmets sustained head injury. Therefore, all stakeholders should promote protective gear use among motorcyclists and their passengers.

Non-use of protective gear is a major clinical problem because the resultant head injury is associated with high mortality and morbidity, prolonged hospital stay and chronic disability among the victims^{8,21-22}. It is also a significant public health problem partly because trauma related treatment is very costly at about US \$369 per victim of commercial motorcycle related injuries⁸. Motorcycle related injuries consume 4.2% of Mulago hospital annual budget²³. Trauma alone takes up nearly 2% of Uganda's Gross Domestic Product¹⁴. For a country that struggles with under funding of healthcare²⁴, these are colossal sums of money to spend on a single preventable disease condition. On the other hand, as seen in this study and many others, most victims are young men in their more productive years^{6,8,10,15-16}. The resultant reduced productivity and chronic disability that many of these crash victims suffer makes them a financial and social burden to their families and the community at large. With the right strategies, these injuries can be reduced or their severity mitigated significantly. This would potentially reduce the associated huge expenses and social misery. This study attests to the inadequacy of the efforts to promote helmet use in Uganda.

In slightly over half of the cases in this study as in several others^{9,15}, the crashes were between a motorcycle and other vehicles. Could this be due to low visibility of the motorcycles to other drivers²⁵ or the previously reported recklessness?¹⁸ Rider conspicuity to other road users can be significantly improved by use of reflective clothing and "always on" head lamp policy as show from previous work²⁵. The occurrence of head injury had no significant relationship with number of people on the motorcycle at the time of crash. Being that in most instances (87.5%) the number of people was within the legal limit of two, there is need for additional preventive measures. Nearly two thirds of victims reached the hospital by way of private vehicle and police vehicle. Hopefully the current efforts by police to increase the emergency response capacity¹³ will lead to more people arriving in safer conditions by ambulance.

The Uganda Police Force recognizes that enforcing traffic laws reduces road traffic injuries but reportedly lacks the capacity to do so¹³. Traffic law enforcement has significantly reduced road traffic injuries in Rwanda and Tanzania and needs to be supported in Uganda as well²⁶⁻²⁷.

Limitations

We may not have captured all motorcycle crash victims

as many victims including those who sustained minor or fatal injuries may not have been seen at Mulago hospital. Also, because not wearing helmets is unsafe and illegal, motorcyclists could have claimed to have been wearing protective gear thus introducing reporting bias leading to a spuriously higher prevalence. The findings of this study should thus be interpreted with caution.

Conclusion

This study revealed that the vast majority of motorcycle crash victims seen at Mulago hospital sustained head injury. Only a minority of these victims were using protective gear at the time of the crash. Also, the use of protective gear significantly reduced the occurrence and severity of head injury among crash victims.

Therefore, more efforts need to be invested in increasing the use of protective gear to reduce the occurrence and severity of head injury among commercial motorcycle riders in Uganda.

Competing interest

The authors declare no conflict of interest.

Acknowledgements

This research project was funded by a research grant from the Global Partners of Anesthesia and Surgery (GPAS) through the department of surgery at Makerere University College of Health Sciences. The funders had no influence on the design and conduct of the study or its data analysis and manuscript writing.

References

1. Lagarde E. Road Traffic Injury Is an Escalating Burden in Africa and Deserves Proportionate Research Efforts. *PLoS Med.* [doi:10.1371/journal.pmed.0040170]. 2007;4(6):170.
2. Nantulya VM, Reich MR. The neglected epidemic: road traffic injuries in developing countries. *BMJ.* 2002 May 11;324(7346):1139-1141.
3. Peden M, Scurfield R, Sleat D, Mohan D, Hyder AA, Jarawan E, Mathers. World Report on Road Traffic Injury Prevention Geneva. World Health Organisation; 2004.
4. Laxminarayan R, Mills AJ, Breman JG, Measham AR, Alleyne G, Claeson M, et al. Advancement of global health: key messages from the Disease Control Priorities Project. *Lancet.* 2006 Apr 8;367(9517):1193-1208.
5. Burden of disease study. Ministry of Health, Uganda. 1996.
6. Center-Uganda IC. Final report for the Hospital and Trauma Surveillance Program in Uganda. 2010:8.
7. Chalya P, Mabula J, Dass R, Mbelenge N, Ngayomela I, Chandika A, et al. Injury characteristics and outcome of road traffic crash victims at Bugando Medical Centre in Northwestern Tanzania. *Journal of Trauma Management & Outcomes.* 2012;6(1):1 PubMed .
8. Galukande M, Jombwe J, Fualal J, Gakwaya A. Boda-boda Injuries a Health Problem and a Burden of Disease in Uganda: a Tertiary Hospital Survey. *East and Central African Journal of Surgery* 2009;14(2):33 PubMed -37
9. Naddumba E. A cross sectional retrospective study of boda boda injuries at Mulago Hospital in Kampala. Uganda. *East Cent Afr J Surg.* 2004;9:44 - 47.
10. Kitara DL. Boda Boda Injuries in Gulu Regional Hospital, Northern Uganda. *East and Central African Journal of Surgery.* 2011;16(2):97-101.
11. Tumwebaze F. Ministerial Statement on Motorcycle registration. [cited 2015 07 Feb]; Available from: <http://www.officeofthepresident.go.ug/index.php/tenders/item/127-ministerial-statement-on-motorcycle-registration>.
12. <http://www.ipifa.net/downloads/presentations/Boniface%20MutatinaUganda%20-%20Helmet%20Surveys%20abstract.pdf>. accessed 31/10/13 4:50 PM.
13. Force TUP. Annual Crime and Traffic/ Road Safety Report. 2012:31.
14. Hsia RY, Ozgediz D, Mutto M, Jayaraman S, Kyamanywa P, Kobusingye OC. Epidemiology of injuries presenting to the national hospital in Kampala, Uganda: implications for research and policy. *Int J Emerg Med.* 2010;3(3):165 PubMed -172.
15. Khanbhai M, Lutomia MBL. Motorcycle Accident injuries seen at Kakamega Provincial Hospital in Kenya. *East and Central African Journal of Surgery.* 2012;17(1):43-46.
16. Chalya P, Mabula J, Ngayomela I, Kanumba E, Chandika A, Giiti G, et al. Motorcycle injuries as an emerging public health problem in Mwanza City, north-western Tanzania. *Tanzan J Health Res.* 2010;12:214 - 221.
17. Johnson OE. Prevalence and pattern of road traffic accidents among commercial motorcyclists in a city in Southern Nigeria. *Educational Research.* 2012;3(6):537-542.
18. Kigera J W M, Naddumba EK. Patterns Of Injuries After Road Traffic Crashes Involving Bodabodas The Annals of African Surgery. 2010;5:12-15 PubMed.
19. LaTorre G, Bertazzoni G, Zotta D, van Beeck E, Ricciardi G. Epidemiology of accidents among users of two-wheeled motor vehicles. A surveillance study in two Italian cities. *Eur J Public Health.* 2002 Jun;12(2):99-103.

20. Conrad P, Bradshaw YS, Lamsudin R, Kasniyah N, Costello C. Helmets, injuries and cultural definitions: motorcycle injury in urban Indonesia. *Accident Analysis and Prevention* 1996 Mar;28(2):193-200.
21. MacLeod JB, Digiacomio JC, Tinkoff G. An evidence-based review: helmet efficacy to reduce head injury and mortality in motorcycle crashes: EAST practice management guidelines. *J Trauma*. 2010 Nov;69(5):1101-1111.
22. Liu B, Ivers R, Norton R, Blows S, Lo SK. Helmets for preventing injury in motorcycle riders. *Cochrane Database Syst Rev*. 2004(2):CD004333.
23. Kigera J NL, Naddumba E K. The Impact of Boda-boda Motor Crashes on the Budget for Clinical Services at Mulago Hospital, Kampala. *East and Central African Journal of Surgery*. 2010;15(1):57-61.
24. Zikusooka CM, Kyomuhang R, Orem JN, Tumwine M. Is health care financing in Uganda equitable? *Afr Health Sci*. 2009 Oct;9 Suppl 2:S52-58.
25. Wells S, Mullin B, Norton R, Langley J, Connor J, Jackson R, et al. Motorcycle rider conspicuity and crash related injury: case-control study. *BMJ*. 2004 2004-04-08 PubMed 00:00:00;328(7444):857.
26. Twagirayezu E, Teteli R, Bonane A, Rugwizangoga E. Road traffic injuries at Kigali University Central Teaching Hospital, Rwanda. *East Cent Afr J Surg*. 2008;13:73 - 76.
27. Mcharo B. Motorcycle crash: injuries pattern and associated factors among patients treated at Muhimbili Orthopaedic Institute. Dissertation for Master of Medicine (Orthopaedic and Traumatology) of Muhimbili University of Health and Allied Sciences. 2012.