

Prevalence and associated factors of TB/HIV co-infection among HIV Infected patients in Amhara region, Ethiopia.

Aweke Abebaw Mitku, Zelalem Getahun Dessie, Essey Kebede Muluneh, Demeke Lakew Workie

Department of Statistics, Bahir Dar University, Bahir Dar, Ethiopia.

Abstract

Background: Tuberculosis is one of the world's most common causes of death in the era of Human immunodeficiency virus. The purpose of this study was to determine the prevalence and associated factors of TB/HIV co-infection.

Methods: Hospital based retrospective studies were conducted among adult HIV-positive patients. Logistic regression method and Chi square test were applied.

Results: A total of 571 HIV positive study participants were enrolled. Of these, 158 (27.7%) were found to have pulmonary tuberculosis. Lower baseline CD4 count <200 cell/ μ l, patients who drunk alcohol, patients who were ambulatory at the initiation of ART, patients whose marital status was single were significant predictors for increased risk of tuberculosis in PLWHIV ($P < 0.05$). Non smoker patients, patients in WHO clinical stage I, patients in WHO clinical stage II and ownership of the house had significant protective benefit against risk of TB ($P < 0.05$).

Conclusion: The prevalence of TB/HIV co-infection in adults on ART in our study was moderately high. Having advanced clinical status and presence of risk factors were found to be the predicting factors for co-infection. The health office should open TB/HIV co-infection units in the hospitals and health workers should be cautious when a patient has an advanced disease.

Keywords: Tuberculosis, co-infection, HIV/AIDS, risk factors

DOI: <http://dx.doi.org/10.4314/ahs.v16i2.29>

Cite as: Mitku AA, Dessie ZG, Muluneh EK, Workie DL. Prevalence and associated factors of TB/HIV co-infection among HIV Infected patients in Amhara region, Ethiopia. *Afri Health Sci* 2016;16(2): 588-595. <http://dx.doi.org/10.4314/ahs.v16i2.29>

Background

Tuberculosis (TB) is one of the world's most common causes of death in the era of Human Immunodeficiency Virus (HIV). It is among the leading causes of death for people living with HIV (PLWHIV) which shares about twenty-five percent of all causes of the deaths¹. HIV promotes progression of latent or recent infections of Mycobacterium tuberculosis to active disease and also increases the rate of occurrence of TB². PLWHIV may also be more susceptible to TB infection³. HIV is the first and TB is the second leading cause of death from infectious disease worldwide⁴. Globally, there were an estimated 9 million incident cases of TB in 2013 with 13% co-

infected with HIV. There were an estimated 1.5 million TB deaths (1.1 million among HIV-negative people and 0.4 million among HIV-positive people)⁵. Hospital data indicated that TB is the leading cause of morbidity and the third cause of hospital admissions in the country⁶. According to World Health Organization report Ethiopia ranks 7th among the 22 high burden countries with TB and HIV infection in the world⁷.

TB/HIV co-infection is associated with special diagnostic and therapeutic challenges and constitutes an immense burden on healthcare systems of heavily infected countries like Ethiopia⁸. Studies indicated that certain HIV-infected people develop TB, while others do not. This phenomenon iterates that being HIV positive is not the only factor for being infected with TB, and there are various determinant factors that contribute to the TB/HIV co-infection⁹⁻¹¹. Understanding the predictors of TB/HIV co-infections in the local context is critical for Ethiopia to improve TB/HIV co-infected patients' co-management. Therefore, the main goal of this study was

Corresponding author:

Zelalem Getahun Dessie,
Department of Statistics,
Bahir Dar University, Bahir Dar, Ethiopia.
Email: zelalem_getahune@yahoo.com

to determine prevalence and associated factors of TB co-infection among HIV patients in the HIV clinics in Amhara Region, Ethiopia. It is anticipated that findings from this study will contribute to the body of knowledge that informs TB/HIV program planners, decision makers, and project implementers by providing predictors of TB/HIV co-infection.

Methods

Study design and area

Hospital based retrospective studies were conducted among adult HIV-positive patients attending HIV clinic between June 2006 and January 2014 in Amhara Region, Ethiopia. The Amhara Region is located in the North-Western and North Central parts of Ethiopia and lies within 9° and 23°45'N and 36° and 40°30'E. Elevation ranges from 700m in the Eastern parts to over 4620m in the NorthWest. It has a total area of 170,000km², which is divided in to 11 administrative Zones and 105 Weredas. In this region, there are 5 referral hospitals. The sources of data for this study were Felege Hiwot, Gonder University, Dessie, Debere Berihan and Debere Markos referral Hospitals. The sampling frame consisted of 65000 HIV/AIDS patients who have visited these referral hospitals since the initiation of ART.

Sample size and sampling procedure

Sample size was calculated using single population proportion formula using 95% confidence level, 4% degree of precision and 33% proportion of TB/HIV co-infection¹⁷. The sample size was 520 which after adding 10% for non-response gave an overall sample size of 572. A stratified random sampling method was adopted for selecting a representative sample. The sample size was allocated proportionally based on the number of patients in each referral hospital. Patients were selected randomly using their ART unique identification number. The study considered all HIV infected patients on ART, whose age was >15 years regardless of their treatment category during the study period in the referral hospitals.

Measurements

The response variable for this study was the occurrence of TB/HIV co-infection. The predictor variables included socio-demographic and economic characteristics (age, sex, income, employment status, residence, educational status, and marital status) and clinical related characteristics

(WHO clinical stage, baseline CD4 count, month on ART, functional status, and smoking status).

Functional status was measured at baseline, and a person was categorized as working if “he/she was able to perform usual work in or out of the house”; ambulatory if “he/she was able to perform activities of daily living” and bedridden if “he/she was not able to perform activities of daily living”. Alcohol use was categorized as “yes” if a person was currently or previously an alcoholic beverages consumer and “no” if he/she had never been an alcoholic beverages consumer.

Data collection

Nurses who worked in ART clinics were selected to collect data from January, 2013 to January, 2014. A structured questionnaire was used to collect data on socio demographic, behavioral and economic factors. All records of HIV/AIDS patients between June 2006 and September 2014 were considered for data collection. Lab requests, follow-up forms, anti-TB record forms, ART intake forms, and patient cards were reviewed. Laboratory results recorded before starting ART were used as base line values. If there was no pre-treatment laboratory test, however, results obtained within one month of ART initiation were considered as baseline values. Data quality was assured by using a pre-tested data collection tool and trained data collectors. Two professionals were engaged in continuous supervision and monitoring. Completeness and consistency of data was checked by supervisors, data clerks and investigators before and after data entry.

Ethics statement

This investigation was conducted according to the principles expressed in the Declaration of Bahir Dar University, Ethiopia. It was approved by the research ethics committee at Bahir Dar University and all participants who agreed to participate in this study signed a consent form.

Data analysis

The data was coded and entered to Epi Info 3.14 for Windows. Statistical package for social science (SPSS) version 20 and STATA version 12.0 were used for analysis. Descriptive statistics were used to assess normality, outliers and identify missing values. Chi square test and independent t-test were used to test the relationship between each covariate and TB/HIV co-infection status. Multi covariate logistic regression was done by entering all variables

with *P* value less than 0.25 in the bivariate analysis. Finally, logistic regression with forward likelihood selection method with *P* value less than 0.05 was used to identify the determinant factors associated with TB/HIV co-infection. The crude and adjusted odds ratio (OR) and its 95% confidence interval (CI) were estimated.

Results

Descriptive statistics for socio-demographic and economic factors

There were 571 respondents in the study. Of these, 413 (72.3%) were not found to have TB/HIV-co infection while 158(27.7%) had TB/HIV co-infection. The proportion of female respondents who had TB/HIV-co infection accounted for a larger proportion in the sample 107(69%) compared to male respondents 48(31%). Patients with primary education accounted for the larger proportion 64(40%) of having TB/HIV co-infection compared to those with no education 39(24.8%), secondary education 33(21%), and certificate and above 21(13.4%).

In addition, majority of TB/HIV patients 130(82.8%) were urban residents while 27(17.2%) were residing in rural areas. Among TB/HIV co-infected patients, 149(96.8%) were non smokers 123(79.9%) of TB/HIV co-infected patients were nonalcoholics. The majority of participants who had TB/HIV co-infection, 104(67.5%) were in WHO clinical stage of III followed by WHO clinical stage IV 23(14.9%) and WHO clinical stage II 19(12.3%). Only 8(5.2%) study participants who have TB/HIV-co infection were found to be in WHO clinical stage I.

Chi-square test shows that the TB/HIV co infection is significantly associated with marital status, alcohol intake, baseline CD4 count, baseline WHO clinical stage, baseline functional status, TB Smear type (*p*-value < 0.05) (Table 1).

The mean monthly income of HIV infected participants was 948.99±800 birr and TB/HIV co-infected patients had mean monthly income of 900+707.11 birr. The mean age of HIV mono infected participants was

33.00+2.83 and TB/HIV co-infected patients had mean age of 35.83+8.4 (Table 1).

Multivariate analysis for predicting TB/HIV co-infection One problem of single covariate approach is that it ignores the possibility that a collection of variables, each of which is weakly associated with the outcome, can become an important predictor of the outcome when taken together. It is therefore important to reduce the possibility of excluding variables at the univariable analysis stage. It is for this reason that a univariate test *p*-value of 0.25 or less was used for selection of variables for the multi covariate analysis from single covariate findings. Based on this marital status, alcohol intake, smoking status, baseline CD4 count, baseline functional status, home ownership and baseline WHO stage were selected with forward likelihood logistic regression method.

After adjusting other covariates, a patient whose marital status was single was 2.17 times more likely to develop TB than those patients whose marital status was divorced and widowed (Adjusted OR=2.17, 95% CI=1.19-3.96). A non smoker was about 69% less likely to develop TB than those who were smokers (Adjusted OR=0.31, 95% CI= 0.10-0.96).

Low CD4 T-lymphocytes count was the strongest clinical predictor of TB/HIV co-infection. Patients with CD4 T-lymphocytes count less than 200 cells/ μ l were 1.71 times more likely to develop TB as compared to patients with CD4 T-lymphocytes count of more than 200 cells/ μ l (Adjusted OR=1.71, 95%CI=1.09-2.68). Patients in WHO clinical stage I were 78% less likely to develop TB than that of the patient in WHO clinical stage IV (Adjusted OR=0.22, 95%CI=0.08-0.58). Similarly, patients in WHO clinical stage II were about 70% less likely to develop TB than those patient in WHO clinical stage IV (Adjusted OR=0.30, 95%CI=0.14-0.66). Patients with ambulatory baseline functional status (Adjusted OR=1.73, 95%CI=1.10-2.78) had an increased risk to develop TB. A patient who had a home was about 41% less likely to develop TB as compared to those patients who did not have their own home (Adjusted OR=0.59, 95%CI=0.39-0.92). Furthermore, patients who drunk alcohol were 2.26 times more likely to develop TB than those who did not drink (adjusted OR=2.26, 95%CI=1.29-4.02) (Table 2).

Table 1 Summary result of TB/HIV co-infection vs. socio-demographic, economic and clinical, and risk variables in Amhara Region, Ethiopia.

Variables	TB/HIV Co infection		Chi-square p-value
	No, n(%)	Yes, n(%)	
Sex			
Male	112(27.8)	48(31)	0.457
Female	291(72.2)	107(69)	
Age			
Mean±SD	33.00+2.83	35.83+8.4	0.894
Monthly Income			
Mean±SD	948.99±800	900+707.11	0.058
Educational Status			
No Education	106(26.2)	39(24.8)	0.961
Primary	168(41.6)	64(40.8)	
Secondary	81(20)	33(21)	
Certificate and above	49(12.1)	21(13.4)	
Marital Status			
Single	48(11.8)	34(21.7)	0.017*
Married	204(50.1)	67(42.7)	
Divorced/separated	90(22)	37(23.6)	
Widowed	65(16)	19(12.1)	
Employment status			
Employed	209(51.9)	70(44.9)	0.295
Unemployed	94(23.3)	39(25)	
Retired	100(24.8)	47(30.1)	
Residence			
Urban	350(86.8)	130(82.8)	0.219
Rural	53(13.2)	27(17.2)	
Smoking Status			
Smoker	19(4.7)	5(3.2)	0.758
Non smoker	388(95.3)	149(96.8)	
Alcohol Intake			
Yes	49(12.3)	31(20.1)	0.005*
No	351(87.8)	123(79.9)	
Baseline CD4 count			
≤200	254(61.5)	121(76.6)	0.001*
>200	159(38.5)	37(23.4)	
Baseline WHO stage			
Stage I	53(13.1)	8(5.2)	0.000*
Stage II	94(23.2)	19(12.3)	
Stage III	226(55.7)	104(67.5)	
Stage IV	33(8.1)	23(14.9)	
Missed to take HIV medication			
Yes	3(18.8)	6(3.9)	0.012*
No	13(81.2)	147(96.1)	
Baseline functional status			
Bed driven	25(6.3)	16(10.5)	0.009*
Ambulatory	85(21.4)	47(30.7)	
Working	287(72.3)	90(58.8)	
TB Smear type			
Positive	5(9.3)	21(44.7)	0.000*
Negative	48(88.9)	25(53.2)	

*The relationship is a significant at $\alpha=0.05$

Table 2 Multi-Covariate and Bivariate Analysis result for different socio-economic, demographic, clinical and risk variables that affect TB/HIV Co-infection, Amahra region, Ethiopia.

Variables	COR(95%CI)	COR value	P-	AOR(95% CI)	AOR value	P-
Marital Status						
Single	1.96(1.15-3.35)	0.014*		2.17(1.19-3.96)	0.012*	
Married	0.91(0.60-1.37)	0.650		1.01(0.64-1.59)	0.971	
Divorce and others	1			1		
Smoking status						
Non smoker	0.69(0.25-1.87)	0.247		0.31(0.10-0.96)	0.042*	
Smoker	1			1		
Alcohol Intake						
Yes	1.81(1.10-2.96)	0.005*		2.26(1.29-4.02)	0.005*	
No	1			1		
Baseline CD4 Count						
≤200 cells/μl	2.05(1.35-3.11)	0.001*		1.71(1.09-2.68)	0.019*	
>200 cells/μl	1			1		
Baseline WHO Stage						
Stage I	0.22(0.09-0.54)	0.002*		0.22(0.08-0.58)	0.002*	
Stage II	0.29(0.14-0.59)	0.001*		0.30(0.14-0.66)	0.003*	
Stage III	0.66(0.37-1.18)	0.108		0.63(0.33-1.18)	0.147	
Stage IV	1			1		
Baseline Functional Status						
Bed driven	2.04(1.04-3.99)	0.037*		1.69(0.83-3.49)	0.150	
Ambulatory	1.76(1.15-2.71)	0.009*		1.73(1.10-2.78)	0.020*	
Working	1			1		
Home Ownership						
Yes	0.60(0.41-0.89)	0.012*		0.59(0.39-0.92)	0.018*	
No	1			1		
Residence						
Urban	0.73(0.44-1.21)	0.220				
Rural	1					
Missed to take HIV Medication						
Yes	0.18(0.04-0.79)	0.023*				
No	1					

Keys: COR (Crude odds ratio)

AOR (Adjusted Odds Ratio)

CI (Confidence Interval)

¹ (Reference Category)

*Odds Ratio is a significant at $\alpha=0.05$

Discussion

The prevalence of TB/HIV co-infection in adults on ART in our study was moderately high at 27.7%. This is in line with other studies¹²⁻¹⁵. It was slightly lower than 32.8% reported from Nigeria¹⁶ and 33% reported from Ethiopia¹⁷. However, the findings of this study were higher compared to studies conducted in Ethiopia (7.5%), Nigeria

(7.8%) and Tanzania (8.5%)^{2,19-20}. These wide variations in the co-infection rates of TB/HIV across the globe, as reported, can partly be accounted for by the following reasons: coverage level of highly active antiretroviral treatment (HAART), under-reporting, diagnostic procedures used, difference in TB diagnosis, epidemiology of TB in different countries and study methodology applied.

Our study also revealed the determinants of TB/HIV co-infection among HIV-positive adults attending clinical care in Amhara region, Ethiopia. Baseline functional status, CD4 count, WHO clinical stage, alcohol use, smoking status and home ownership status were independent predictors of TB/HIV co-infection among HIV-positive adults.

As CD4+ lymphocyte count decreased the body defence mechanism will be overwhelmed by various opportunistic infections. The results showed that, patients with CD4 T- lymphocytes count less than 200 cells/ μ l were about 2 times more likely to develop TB as compared to CD4 T-lymphocytes count more than 200 cells/ μ l. A study conducted in Nigeria revealed similar finding where lower CD4+ lymphocyte count was observed in co-infected patients than mono infected patients²¹. Substance use such as cigarette smoking and alcohol consumption were the predictors of TB/HIV co-infection in HIV positive in the current study. A study conducted in West Africa revealed similar finding where cigarette smoking and alcohol consumption were the risk factors for the development of TB²². In contrast, a study from the Gambia found that cigarette smoking and alcohol consumption were not associated with TB²³.

In our study we found that marital status was significantly associated with TB. Divorced or widowed Patients were less likely to develop TB compared to unmarried (single), which is consistent with other reports in West Africa and Ethiopia²⁴. It might be explained as unmarried (single) persons are younger than married persons and have a different lifestyle, especially males, who often migrate to towns in search of a job where they live alone or with friends. Similarly, education was not the predictor of TB/HIV co-infection in HIV clinic. This result is consistent with Kiberet et al.¹⁷ and Wondimeneh et al.²⁰ in Ethiopia. In contrast, studies from south west Ethiopia^{9,25} have showed that low level of education was associated with TB. This could be due to the high prevalence of literates in our study population.

The other important finding identified was association of patient's WHO clinical stage with TB-co infection. Those patients in first and second WHO clinical stages were about 78% and 70% less likely to develop TB compared with those in WHO clinical stages IV, respectively. Congruent findings have been reported from in-country

and outside of the country^{10,26,27}. This could be explained as once the patients get into late stages, the immunity protective capacity will be minimal which would make them prone to tuberculosis infection. A worth to mention as well is that TB is one of the AIDS defining criteria to categorize the patients in to the late WHO clinical staging which was also used as a criteria in HIV/AIDS clinics in Ethiopia.

Patients who were ambulatory at the initiation of ART were about 2 times more likely to develop TB than those who were working at the initiation of ART. This is consistent with the retrospective cohort study in Ethiopia¹⁷⁻¹⁸. Furthermore, ownership of the house by the TB patient's family was a predictor of TB/HIV co-infection among HIV positive. A study conducted in West Africa revealed similar finding where ownership of the house by the TB patient's family was associated with lower risk for developing TB²².

Conclusion and recommendations

The study identified marital status, functional status, WHO clinical staging, baseline CD4 count, smoking status, alcohol intake, and home ownership were associated with TB/HIV co-infection among adult HIV positives. In general, advanced clinical disease, smoking and alcohol consumption were found to be the main predictors for TB/HIV co-infection. All PLWHIV should be screened for TB, but for substance abusers and patients with advanced disease (WHO clinical stage IV, being ambulatory and CD4 count < 200 cell/ μ l) intensified screening is highly recommended during treatment follow up. Since the results of the study underlined risk factors (alcohol use and smoking status) as the predictors of TB/HIV co-infection in HIV positive, physicians are expected to work hard to bring about behavioural changes towards substance use. According to the results of this study, the main factors associated with TB/HIV co-infection are clinical variables. So, health workers should be cautious when a patient has lower CD4 counts, is ambulatory at the initiation of ART and in WHO clinical stage IV. Furthermore, the health office should open TB/HIV co-infection units in the Hospitals.

Competing interest

The authors declare that they have no competing interests.

Authors contributions

ZG and AA Designed the study, reviewed the questionnaires, collected the data, analyzed data and wrote the article. EK designed the study, reviewed the questionnaires and critically edited the manuscript. DL designed the study, collected the data and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The study was funded by Bahir Dar University, Ethiopia. Our special gratitude goes to Dr Yewulsew Kassie (Regional coordinator MSH/Heal TB, USAID) for his immense and invaluable advice and guidance that contributed to the successful realization of this study. Our appreciation also goes to all ART members and clinic staff of the Amahara region Referral Hospitals, Ethiopia for their cooperation during data collection.

References

1. World Health Organization: Tuberculosis. Fourth Edition, The Essentials, Edited By Dr Mario C. Raviglione, World Health Organization, Geneva Switzerland, 2010, http://www.who.int/tb/features_archive/the_essentials/en/.
2. Ilyasu Z, Babashani M: Prevalence and prediction of TB co-infection among HIV seropositive patients attending Aminu Kano Teaching Hospital, Northern Nigeria. *J. Epid.* 2009, 19: 81-87 PubMed ,doi.org/10.2188/ica.J20080026.
3. Stop TB Partnership: Talking Points-TB/HIV, [updated 2007 June; cited 2014 28 March]. Available from: www.stoptb.org/ng/tbhiv/assesseddocuments/tbhivtalkingpoints.pdf.
4. World Health Organization: Global tuberculosis report 2012. [http://apps.who.int/iris/bitstream/10665/75938/1/9789241564502_eng.pdf].
5. World Health Organization: Global Tuberculosis Report 2014, Geneva, Switzerland, 2014: World Health Organization.
6. Federal Ministry of Health Ethiopia: Tuberculosis, Leprosy and TB/HIV Prevention and Control Programme Manual, Fourth Edition, MOH, Addis Ababa, Ethiopia, 2008.
7. World Health Organization: WHO Country Cooperation Strategy 2012–2015, Ethiopia, WHO Regional Office for Africa, Republic of Congo, 2013, <http://www.afro.who.int/>.
8. World Health Organization: Treatment of tuberculosis guidelines. 4th edition. Geneva: World Health Organization; 2010.
9. Taha M, Deribew A, Tessema F, Assegid S, Duchateau L, and Colebunders R: “Risk factors of active tuberculosis in people living with HIV/AIDS in southwest Ethiopia: a case control study, in Jimma Hospital and Karl Hospital in Southwest Ethiopia,” *Ethiopian Journal of HealthDevelopment* 2009, 219: 131–139,doi.org/10.4314/ejhs.v21i2.69053
10. Akanbi MO, Achenbach CJ, Feinglass J, Taiwo B, Onu A, et al.: Tuberculosis after one year of combination antiretroviral therapy in Nigeria: a retrospective cohort study. *AIDS Res Hum Retroviruses* 2013, 29: 931-937, DOI: 10.1089/aid.2012.0231.
11. Molaeipoor L, Poorolajal J, Mohraz M, Esmailnasab N: Predictors of tuberculosis and human immunodeficiency virus co-infection: a case-control study. *Epidemiology and Health*, 2014:36, Article ID: e2014024, DOI: 10.4178/epih/e2014024.
12. Omair MA, Al-Ghamdi AA, Alrajhi AA: Incidence of tuberculosis in people living with the human immunodeficiency virus in Saudi Arabia. *Int J Tuberc Lung Dis.* 2010, 14:600–3.
13. Jam S, Sabzvari D, SeyedAlinaghi S, Fattahi F, Jabbari H, Mohraz M: Frequency of Mycobacterium tuberculosis infection among Iranian patients with HIV/AIDS by PPD test. *Acta Med Iran.* 2010, 48:67–71.
14. Ifebunandu NA, Ukwaja KN, Obi SN: Treatment outcome of HIV-associated tuberculosis in a resource-poor setting. *Trop Doct.* 2012, 42:74–6, DOI: 10.1258/td.2011.110421.
15. Rajasekaran S, Mahilmaran A, Annadurai S, Kumar S, Raja K: Manifestation of tuberculosis in patients with human immunodeficiency virus: A large Indian study. *Ann Thorac Med* 2007, 2:58–60.
16. Awoyemi OB, Ige M, Onadeko BO: Prevalence of active pulmonary tuberculosis in human immunodeficiency virus seropositive adult patients in University College Hospital, Ibadan, Nigeria. *Afr J Med* 2002, 31:329–PubMed ;332.
17. Kibret KT, Yalew AW, Belaineh BG, Asres MM: Determinant Factors Associated with Occurrence of Tuberculosis among Adult People Living with HIV after Antiretroviral Treatment Initiation in Addis Ababa, Ethiopia: A Case Control Study. *PLoS One* 2013 8:5: e64488. PubMed

- doi:10.1371/journal.pone.0064488,DOI: 10.1371/journal.pone.0064488.
18. Fekadu S, Teshome W, Alemu G: Prevalence and determinants of Tuberculosis among HIV infected patients in south Ethiopia, *J Infect Dev Ctries* 2015; 9(8):898-904, doi:10.3855/jidc.5667.
 19. Ngowi BJ, Mfinanga SG, Bruun JN, Morkve O: Pulmonary tuberculosis among people living with HIV/AIDS attending care and treatment in rural northern Tanzania. *BMC Publ Health* 2008, 8:341, DOI: 10.1186/1471-2458-8-341.
 20. Wondimeneh Y, Muluye D and Belyhun Y: Prevalence of Pulmonary tuberculosis and immunological profile of HIV co-infected patients in Northwest Ethiopia. *BMC Research Notes* 2012 5:331, DOI:10.1186/1756-0500-5-331.
 21. Agbaji O, Ebonyi AO, Meloni ST, Anejo-Okopi JA, Akanbi MO, et al.: Factors Associated With Pulmonary Tuberculosis-HIV Co-Infection in Treatment-Naive Adults in Jos, North Central Nigeria. *J AIDS Clin Res* 2013, 4: 222, doi: 10.4172/2155-6113.1000222.
 22. Lienhardt C, Fielding K, Sillah JS, Bah B, Gustafson P, et al: Investigation of the risk factors for tuberculosis: a case-control study in three countries in West Africa. *Int J of Epidemiol* 2005, 34: 914– PubMed ;923, doi: 10.1093/ije/dyi100.
 23. Teck R, Ascurra O, Gomani P et al.: “WHO clinical staging of HIV infection and disease, tuberculosis and eligibility for antiretroviral treatment relationship to CD4 lymphocyte counts,” *International Journal of Tuberculosis and Lung Disease* 2005, 9: 258–262.
 24. Melkamu H, Seyoum B, and Dessie Y: Determinants of Tuberculosis Infection among Adult HIV Positives Attending Clinical Care in Western Ethiopia: A Case-Control Study. *AIDS Research and Treatment* 2013, Article ID 279876, 7 pages <http://dx.doi.org/10.1155/2013/279876>.
 25. Kapata N, Chanda-Kapata P, Michelo C: The social determinants of tuberculosis and their association with TB/HIV co-infection in Lusaka, Zambia. *Medical Journal of Zambia* 2013, 40: 2.
 26. Lawn SD, Badri M, and Wood R: “Tuberculosis among HIV infected patients receiving HAART: long term incidence and risk factors in a South African cohort,” *AIDS*, vol. 19, no. 18, pp. 2109–2116, 2005 PubMed .
 27. Hill Philip C, Jackson-Sillah D, Donkor SA, Out J, Ad-egbola RA, et al: Risk factors for pulmonary tuberculosis: a clinic-based case control study in The Gambia. *BMC Public Health* 2006, 6:156, doi:10.1186/1471-2458-6-156.