

Sero-prevalence and associated factors of *Helicobacter pylori* infection among adult patients with dyspepsia attending the gastroenterology unit in a tertiary hospital in Mwanza, Tanzania.

Hyasinta Jaka¹, Martha F. Mushi², Mariam M Mirambo², Leonard Wilson²,
Jeremiah Seni², Majigo Mtebe³, Stephen E. Mshana²

1. Department of Internal medicine, Weill Bugando School of Medicine, P.O.Box 1464, Mwanza, Tanzania.
2. Department of Microbiology/Immunology, Weill Bugando School of Medicine, P.O.Box 1464, Mwanza, Tanzania.
3. Department of Microbiology and Immunology, Muhimbili university of Health and allied sciences P.O.Box 65001, Dar es Salaam.

Abstract

Background: Prevalence of *H.pylori* infection varies greatly between populations in different countries. This study was conducted to determine the magnitude of *H.pylori* among adult patients with dyspepsia attending the gastroenterology unit at Bugando medical centre.

Methods: A cross sectional study involving 202 dyspeptic patients was conducted between June and July 2014. A Standardized data collection tool was used to collect socio-demographic characteristics. *H.pylori* antibodies were detected using rapid immunochromatographic tests according to manufacturer's instructions.

Results: The median age of study population was 42 (IQR: 33-54). Females 105 (51.9%) formed majority of the population studied. Of 202 participants; 119 (58.9%) were from rural areas. Seroprevalence of *H.pylori* infection was found to be 79/202 (39.1%, 95% CI: 32.3 -45.7). As the age increased the risk of having *H.pylori* infection also increased (OR: 1.02 95% CI: 1-1.04, P=0.02). On multivariate logistic regression analysis untreated drinking water was found to predict *H.pylori* seropositivity (OR: 2.33, CI: 1.09-4.96, p=0.028).

Conclusion: The seroprevalence of *H.pylori* among dyspeptic patients is high in this setting. Therefore the community in Mwanza should be educated on the use of safe drinking water in order to minimize *H. pylori* infections.

Keywords: *H.pylori*, dyspepsia, Tanzania.

DOI: <http://dx.doi.org/10.4314/ahs.v16i3.7>

Cite as: Jaka H, Mushi MF, Mirambo MM, Wilson L, Seni J, Mtebe M, Mshana SE. Sero-prevalence and associated factors of *Helicobacter pylori* infection among adult patients with dyspepsia attending the gastroenterology unit in a tertiary hospital in Mwanza, Tanzania. *Afri Health Sci* 2016;16(3): 684-689. DOI: <http://dx.doi.org/10.4314/ahs.v16i3.7>

Background

Helicobacter pylori is one of the major public health problems worldwide. More than half of the world's population is infected and humans are considered to be the only reservoirs¹. *H.pylori* infections can lead to gastritis, peptic ulcer, gastric carcinoma and mucosal associated lymphoid tissue (MALT) lymphoma². Majority of the infected individuals are asymptomatic; nevertheless

a small percentage tends to develop manifestations of peptic ulcer diseases later in life¹.

H.pylori is commonly transmitted by fecal oral route through contaminated water and food³. Lack of proper sanitation, basic hygiene, poor diets and overcrowding have been found to play a significant role in *H.pylori* infection⁴. *H.pylori* infection is common throughout the world, the seroprevalence as high as >90% has been reported in many developing countries compared to 1.2-48.8% in developed countries⁵. In Africa the seroprevalence has been found to range between 19.26% and 92%^{6,7} while in Tanzania the prevalence range from 65% -79% among dyspeptic patients^{7,8}. A wide variation in the seroprevalence of *H.pylori* infection within and between different populations has been observed. Despite these variations

Corresponding author:

Mariam M Mirambo,
Department of Microbiology/Immunology,
Weill Bugando School of Medicine,
P.O.Box, 1464, Mwanza, Tanzania
Email: mmmirambo@gmail.com

there is no single study in Mwanza to show the magnitude of previous *H. pylori* infections. This study was done for the first time in Mwanza, Tanzania to investigate the magnitude and factors associated with previous *H. pylori* infection.

Materials and methods

Study design and settings

This was a cross-sectional hospital based study conducted at gastroenterology unit at the Bugando Medical Centre (BMC) between June and July 2014 to investigate the magnitude of *H. pylori* infection. The study included all dyspeptic patients who consented to participate in the study. BMC is a tertiary, consultant and teaching hospital serving a population of about 13 million in the Lake zone of Tanzania.

Sample size and sampling

The sample size was determined by using Kish Leslie formula; a proportion of 22.5% was used⁹. The minimum sample size obtained was 188. Dyspeptic adult patients with no active bleeding were serially enrolled until the sample size was reached.

Data collection

A standardized structured data collection tool was used to collect associated factors such as demographic and general characteristics. Dyspepsia was defined as difficult, or disturbed digestion, which may be accompanied by symptoms such as epigastric pain, nausea, vomiting, bloating and heartburn¹⁰. In this study illiterate and primary school level was defined as low education while secondary school and tertiary level were defined as high education. Treated drinking water was defined as water from the city supply or commercial available bottled water.

To determine *H. pylori* specific IgG antibodies about 4-5ml of blood sample was drawn from participants and placed in a plain vacutainer tubes (BD, UK). Specimens were taken to BMC laboratory where sera were extracted. *H. pylori* IgG antibodies were detected using rapid immunochromatographic kits (Flexsure-HP, SmithKline Diag-

nostics, Inc USA). The test has specificity and sensitivity of 95% and 96% respectively¹¹.

Data analysis

Data was entered in the computer using excel software and analysed using STATA version 11 (STATA Corp LP, USA). The main outcome in this study was being *H. pylori* IgG antibody positive. Categorical variables such as sex, residence, source of drinking water, education level, water treatment, and hand washing before eating were summarized as proportions while age was summarized as median with interquartile range (IQR). Univariate and multivariate logistic regression analysis were done to determine factors predicting *H. pylori* infection. Factors with p-values less than 0.05 were considered statistically significant at 95% confidence interval (CI).

Ethical considerations

The study was approved by the joint CUHAS/BMC research ethics and review committee and permission was obtained from gastroenterology unit administration. Written informed consent was sought from each participant prior to enrolment in the study.

Results


Demographic characteristics

A total of 202 patients were recruited at the gastroenterology unit with median age of 42(IQR: 33-54) years. Females 105(51.9%) formed majority of the study population. Of 202 participants; 119(59%) and 83(41%) were residing in rural and urban areas respectively. Regarding occupational status; 16(7.9%) were students, 46(22.7%) were employed and 140(69.3%) were unemployed.

Prevalence and factors associated with *H. pylori* infections

A total of 79(39.1%; 95% CI: 32.3 -45.7) participants were sero-positive for *H. pylori*. As the age increased the risk of having *H. pylori* infection also increased (OR: 1.02 95% CI: 1-1.04, P=0.02) (figure1).

Figure 1: As the age increases the risk of having *H.pylori* infection also increases by 2% while the seroprevalence of *H.pylori* increases by 0.5%



On univariate logistic regression analysis; low education level was significantly associated with *H.pylori* seropositivity (49.5% vs. 28.7%; OR: 2.43, CI 1.36-4.35, P=0.003). Patients who used untreated drinking water had 2.33 times risk of getting *H.pylori* infection compared to

those used treated water (58.7% vs.30%, P<0.001) table1. Only water treatment was found to be an independent predictor of *H.pylori* seropositivity on multivariate logistic regression analysis (OR: 3.28, CI: 1.77-6.10, P=0.028). (Table 1)

Table1: Univariate and multivariate logistic regression analysis of factors associated with *H.pylori* infection among 202 adult patients

| Characteristics | Positive test (N, %) | Univariate | | Multivariate | |
|-----------------------------------|----------------------|---------------------|--------------|------------------------|--------------|
| | | OR(95% CI) | P value | OR(95% CI) | P value |
| Age(years) | 42(IQR:33-54) | 1.02(1-1.04) | 0.024 | 1.01(0.98-1.03) | 0.469 |
| Sex | | | | | |
| Male(97) | 36(37.11%) | 1 | | | |
| Female (105) | 43(40.95%) | 1.17(0.66-2.07) | 0.577 | 1.31(0.69-2.44) | 0.410 |
| Occupation | | | | | |
| Students(16) | 4(25%) | 1 | | | |
| Not employed(141) | 58(41.13%) | 2.09(0.64-6.82) | 0.219 | | |
| Employed(45) | 17(37.78%) | 1.82(0.51-6.56) | 0.359 | 1.93(0.92-4.02) | 0.08 |
| Education level | | | | | |
| High (101) | 29(28.7%) | 1 | | | |
| Low (101) | 50(49.5%) | 2.43(1.36-4.35) | 0.003 | 1.32(0.64-2.67) | 0.448 |
| Source of drinking water | | | | | |
| Tape water(101) | 32(31.68%) | 1 | | | |
| Other sources(101) | 47(46.53%) | 1.87(1.05-3.32) | 0.031 | 0.83(0.39-1.77) | 0.634 |
| Water treatment | | | | | |
| Yes(139) | 42(30.22%) | 1 | | | |
| No(63) | 37(58.73%) | 3.28(1.77-6.10) | <0.001 | 2.33(1.09-4.96) | 0.028 |
| Hand washing before eating | | | | | |
| Yes(184) | 72(39.13%) | 1 | | | |
| No(18) | 7(38.89%) | 1.02(0.37-2.7) | 0.984 | 1.11(0.37-3.30) | 0.854 |
| Residence | | | | | |
| Urban(83) | 27(32.5%) | 1 | | | |
| Rural(119) | 52(43.70%) | 1.69(0.89-2.88) | 0.111 | 1.00(0.49-2.06) | 0.984 |

Discussion

Although many individuals with *H. pylori* gastric colonization are asymptomatic, often those who seek medical attention may present with dyspepsia. Prolonged infection may result in gastric ulcers formation. The current study indicates the true magnitude of *H.pylori* infection among dyspeptic patients in our settings. Based on the natural history of the disease the seropositive individuals in the present study have 10-20% life time risk of developing peptic ulcers¹² with 1-2% of the infected subjects at risk of developing gastric cancer¹³. This study has found

comparable prevalence of *H.pylori* infection among dyspeptic patients as previously reported in Africa^{7,8,14,15}. As compared to other studies in Africa^{6,16} the observed prevalence in the current study is lower and this could be explained by geographical variation as documented previously^{17,18}. Moreover; the sensitivity of the method used may explain such a difference.

However, this prevalence is higher as compared to previous reports from developed countries¹⁹⁻²². This could be explained by poor living conditions which favor *H.pylori* transmission in developing countries.

In this study the odds of having *H.pylori* infection increased as the age increased which coincides with what has been reported earlier^{14,15,23} and contrary to data from Uganda where the prevalence was found to decrease as the age increased²⁴. This could be explained by different study populations whereby in Uganda study the population studied was patients with different levels of malignancies. As suggested earlier gastric changes such as malignancies may cause decline of *H.pylori* specific antibodies and is likely to occur in patients with advanced disease and age; which is not the case in our study²⁵.

On univariate logistic regression analysis, the odds of having *H.pylori* infections were significantly higher in rural areas compared to urban areas. Similarly, the odds of having *H.pylori* infection were higher among people with low education level than those with higher education level. Our findings are inconsistent with the previous studies which reported higher prevalence rates in urban areas and those with good socioeconomic status^{26,27}. In rural settings of Mwanza majority of population have low socioeconomic status whereby the main economic activities are small scale farming and fishing unlike in urban areas where majority of the urban population have high education level and good socioeconomic status.

In the present study, untreated drinking water was independently found to predict *H.pylori* infections. These findings are in agreement with what has been reported earlier^{3,4} whereby unsafe drinking water and poor hygiene were found to be associated with *H.pylori* infection. Moreover; low education level in almost half of these participants and majority being from rural areas may explain why they cannot afford to treat water despite the fact that it is unsafe to drink.

Study limitations

Due to limited resources, endoscopy was not done; this resulted in failure to diagnose active disease because urease/histopathological and breath tests were not done. In addition stool antigen tests and other differentials of dyspepsia were not established²⁸.

Conclusion and recommendations

Drinking untreated water and increasing age have been found to be associated with *H.pylori* infection. More studies should focus on assessing incidence rates in relation to

age. Improving drinking water in our setting will reduce *H.pylori* transmission and its associated complications such as gastritis, gastric ulcers and gastric malignancies. Moreover, community in Mwanza should be educated on the preventive measures as one of the efforts to control gastric diseases associated with *H.pylori* infection.

Conflict of interest

None declared

Authors' contributions

HJ, MMM, MM and SEM participated in the design of the work. HJ MFM and LW participated in the collection of specimens and clinical data. MMM, JS and LW performed serological tests. MMM, MM, JS and SEM analyzed and interpreted the data. MMM, MFM and SEM wrote the first draft of the manuscript. MMM and SEM critically revised the final draft of the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

The authors would like to acknowledge the technical support provided by Mr.Vitus Silago. We thank all staff in gastroenterology unit at Bugando Medical Centre. This study was supported by research grant from CUHAS to HJ

References

1. Logan R, Gummert P, Schaufelberger H, Greaves R, Mendelson G, Walker M, Thomas P, Baron J, Misiewicz J: Eradication of *Helicobacter pylori* with clarithromycin and omeprazole. *Gut* 1994, 35:323-326.
2. Egan BJ, Holmes K, O'Connor HJ, O'Morain CA: *Helicobacter pylori* gastritis, the unifying concept for gastric diseases. *Helicobacter* 2007, 12:39-44.
3. Cave DR: Transmission and epidemiology of *Helicobacter pylori*. *The American journal of medicine* 1996, 100:12S-18S.
4. Hunt R, Xiao S, Megraud F, Leon-Barua R, Bazzoli F, Van der Merwe S, Vaz Coelho L, Fock M, Fedail S, Cohen H: *Helicobacter pylori* in developing countries. World Gastroenterology Organisation global guideline. *J Gastrointest Liver Dis* 2011, 20:299-304.
5. Hazell S: *H. pylori* in developing countries. In *Helicobacter pylori*. Springer; 1994: 85-94
6. Asombang AW, Kelly P: Gastric cancer in Africa: what do we know about incidence and risk factors? *Transac-*

- tions of the Royal Society of Tropical Medicine and Hygiene 2012, 106:69-74.
7. Arun V, Mchembe M: Endoscopic Findings And Prevalence Of Helicobacter Pylori In Dyspeptic Patients Referred For Upper Gastrointestinal Endoscopy In Dar-Esalaam. *PMJUMU| PIONEER MEDICAL JOURNAL UMUAHLA* 2012, 2.
 8. Mbulaiteye SM, Hisada M, El-Omar EM: Helicobacter pylori associated global gastric cancer burden. *Frontiers in bioscience: a journal and virtual library* 2009, 14:1490.
 9. Mujeeer MP: Prevalence and factors associated with helicobacter pylori infection among children aged one month to twelve years in Mulago hospital. Makerere University, 2008.
 10. Ford AC, Marwaha A, Lim A, Moayyedi P: Systematic review and meta-analysis of the prevalence of irritable bowel syndrome in individuals with dyspepsia. *Clinical Gastroenterology and Hepatology* 2010, 8:401-409.
 11. Graham DY, Evans Jr DJ, Peacock J, Baker JT, Schrier WH: Comparison of rapid serological tests (FlexSure HP and QuickVue) with conventional ELISA for detection of Helicobacter pylori infection. *American Journal of Gastroenterology* 1996, 91.
 12. Kusters JG, van Vliet AH, Kuipers EJ: Pathogenesis of Helicobacter pylori infection. *Clinical microbiology reviews* 2006, 19:449-490.
 13. Correa P, Piazuelo MB: Natural history of Helicobacter pylori infection. *Digestive and Liver Disease* 2008, 40:490-496.
 14. Mbulaiteye SM, Gold BD, Pfeiffer RM, Brubaker GR, Shao J, Biggar RJ, Hisada M: H. pylori- infection and antibody immune response in a rural Tanzanian population. *Infect Agent Cancer* 2006, 1.
 15. Tanih N, Okeleye B, Ndip L, Clarke A, Naidoo N, Mkwetshana N, Green E, Ndip R: Helicobacter pylori prevalence in dyspeptic patients in the Eastern Cape province: race and disease status. *SAMJ: South African Medical Journal* 2010, 100:734-737.
 16. Baako B, Darko R: Incidence of Helicobacter pylori infection in Ghanaian patients with dyspeptic symptoms referred for upper gastrointestinal endoscopy. *West African Journal of Medicine* 1995, 15:223-227.
 17. Perez-Perez GI, Rothenbacher D, Brenner H: Epidemiology of Helicobacter pylori infection. *Helicobacter* 2004, 9:1-6.
 18. Feldman RA, Eccersley AJP, Hardie JM: Epidemiology of Helicobacter pylori: acquisition, transmission, population prevalence and disease-to-infection ratio. *British Medical Bulletin* 1998, 54:39-53.
 19. Chong V, Lim K, Rajendran N: Prevalence of active Helicobacter pylori infection among patients referred for endoscopy in Brunei Darussalam. *Singapore Medical Journal* 2008, 49:42-46.
 20. Rahim AA, Lee YY, Majid NA, Choo KE, Raj SM, Derakhshan MH, Graham DY: Helicobacter pylori infection among Aborigines (the Orang Asli) in the northeastern region of Peninsular Malaysia. *The American Journal of Tropical Medicine and Hygiene* 2010, 83:1119-1122.
 21. Roosendaal R, Kuipers EJ, Buitenwerf J, Van Uffelen C, Meuwissen S, Van Kamp GJ, Vandembroucke-Grauls C: Helicobacter pylori and the birth cohort effect: evidence of a continuous decrease of infection rates in childhood. *The American Journal of Gastroenterology* 1997, 92:1480-1482.
 22. Loffeld R, Van Der Putten A: Changes in prevalence of Helicobacter pylori infection in two groups of patients undergoing endoscopy and living in the same region in the Netherlands. *Changes* 2003, 38:938-941.
 23. Bureš J, Kopáčová M, Koupil I, Voříšek V, Rejchrt S, Beránek M, Seifert B, Pozler O, Živný P, Douda T: Epidemiology of Helicobacter pylori infection in the Czech Republic. *Helicobacter* 2006, 11:56-65.
 24. Newton R, Ziegler JL, Casabonne D, Carpenter L, Gold BD, Owens M, Beral V, Mbidde E, Parkin DM, Wabinga H: Helicobacter pylori and cancer among adults in Uganda. *Infect Agent Cancer* 2006, 1.
 25. Forman D, Newell D, Fullerton F, Yarnell J, Stacey A, Wald N, Sitas F: Association between infection with Helicobacter pylori and risk of gastric cancer: evidence from a prospective investigation. *British Medical Journal* 1991, 6788:1302-1305.
 26. Kawasaki M, Kawasaki T, Ogaki T, Kobayashi S, Yoshimizu Y, Aoyagi K, Takahashi S, Sharma S, Acharya GP: Seroprevalence of Helicobacter pylori infection in Nepal: low prevalence in an isolated rural village. *European Journal of Gastroenterology & Hepatology* 1998, 10:47-50.
 27. Torres J, Leal-Herrera Y, Perez-Perez G, Gomez A, Camorlinga-Ponce M, Cedillo-Rivera R, Tapia-Conyer R, Muñoz O: A community-based seroepidemiologic study of Helicobacter pylori infection in Mexico. *Journal of Infectious Diseases* 1998, 178:1089-1094.
 28. Longstreth GF, Lacy BE, Talley NJ, Grover S: Approach to the adult with dyspepsia. UpToDate; 2013.