

# Family communication about HIV/AIDS and sexual behaviour among senior secondary school students in Accra, Ghana

Samuel Adu-Mireku

Department of Social Sciences, Fayetteville State University, 1200 Murchison Road, Fayetteville, NC 28301, USA

## ABSTRACT

**Background:** Sexually active adolescents in Ghana are increasingly at risk of HIV and other sexually transmitted infections. As a primary agent of socialization, the family can exert a strong influence on adolescent sexual behaviour. Therefore, to aid in the design and implementation of effective prevention programmes, it is important to understand the role of the family in influencing sexual behaviour among school-going adolescents.

**Objectives:** To evaluate the relationship between family communications about HIV/AIDS and sexual activity and condom use among school-going adolescents in Accra, Ghana.

**Method:** A sample of 894 students (56.9% girls, 43.1% boys; mean age = 17.4 years, SD = 1.40) at two senior secondary schools in Accra completed a modified version of the Youth Risk Behavior Survey (YRBS) questionnaire, a self-administered instrument developed by the Centers for Disease Control and Prevention. Analytical techniques utilized included logistic regression and chi-square.

**Results:** Twenty-five percent of the participants reported being sexually experienced, and 73.6% had talked about HIV/AIDS with parents or other family members. Of the sexually experienced students, 64.7% initiated first sexual intercourse by age 16; and 55.7% did not use a condom at last sexual intercourse. Bivariate analysis showed significant gender differences in sexual activity, condom use, and family communication about HIV/AIDS. Logistic regression analysis showed that student-family communication about HIV/AIDS was not associated with sexual activity. However, communication about HIV/AIDS between students and parents or other family members increased the odds of using a condom at last sexual intercourse.

**Conclusions:** The findings of this study suggest that prevention programmes that seek to educate Ghanaian school-going adolescents about sexual risk behaviour must strongly encourage communication about HIV/AIDS between students and family members.

**Key words:** Ghana, adolescent health, risky sexual behaviour, condom use, gender differences.

*African Health Sciences 2003; 3(1): 7 - 14*

## INTRODUCTION

As the HIV/AIDS epidemic continues to create serious public health concerns in Ghana and other sub-Saharan African countries, there is an urgent need to better understand the social processes and factors that influence sexual behaviour so that effective prevention programmes can be designed and implemented. Any such effort must begin with a focus on adolescents and young people, whose responsible sexual behaviours are more likely to positively impact the current struggle against the HIV/AIDS epidemic<sup>1</sup>. The factual basis for such a strategy is clear. As UNAIDS data

indicate<sup>1</sup>, today, more than half of new infected HIV cases are among persons between the ages of 15 and 24.

In Ghana, it is estimated that 1.2 million people will be living with HIV/AIDS by the year 2005, with the majority expected to be adolescents and young people<sup>2</sup>. This estimate is not surprising because previous studies on adolescents and youth in Ghana have documented that they are engaging in sexual activities; they are doing so at an early age; and they are not using condoms consistently<sup>3-8</sup>. By engaging in risky sexual behaviours, adolescents and youth in Ghana face a number of serious negative health consequences, including placing themselves at risk of acquiring HIV and other sexually transmitted diseases. In light of the previous findings, research efforts must continue to identify the social factors and processes that are likely to positively impact the sexual behaviours of adolescents in the country.

For many years, because the family plays a primary role in the socialization of children, researchers have focused on how family members influence adolescent sexual behaviour. In the United States, for example, a large

*Correspondence to:*  
Samuel Adu-Mireku  
Department of Social Sciences  
Fayetteville State University  
1200 Murchison Road  
Fayetteville, NC 28301, USA  
Tel: (910) 672-1042  
Fax: (910) 672-1378  
E-mail: sadu-mireku@uncfsu.edu

body of research has examined the relationship between parent-adolescent communication about sexual issues, including HIV/AIDS, and sexual behaviour<sup>9-11</sup>. The findings from the U.S. studies, so far, have yielded inconsistent results. The majority of the studies, however, have reported that adolescents who discussed sexual issues, including HIV/AIDS, with parents were less likely than those who did not to engage in risky sexual behaviours<sup>9, 12-13</sup>.

In Ghana, to date, none of the studies on adolescents and youth has examined this important question of whether or not students are communicating with their parents and other family members about HIV/AIDS, and whether or not such family communication is associated with safer sexual behaviour. An extension of cross-cultural research on this topic to Ghanaian adolescents is important for two main reasons. First, in Ghana, as in the United States, the institution of the family plays a significant role in the life of adolescents. Therefore, research efforts in Ghana should focus on how the family influences sexual behaviours of adolescents. Ultimately, significant findings from such research should help in the effort to design and implement feasible HIV/AIDS prevention programmes. Second, considering the health risks posed to Ghanaian school-age adolescents that are sexually active, the lack of data and research that examine the association between sexual activities, risk behaviours, and family communication, is a major gap in the existing literature on adolescents and youth in Ghana.

This study, therefore, examined the association between student-family communication about HIV/AIDS and sexual activity and condom use among Senior Secondary School (SSS) students in Accra, Ghana. Following previous studies in the U.S.<sup>9-13</sup>, this study also examined whether or not there are socio-demographic differences in student-family communication about HIV/AIDS, sexual activity, and condom use. However, unlike previous studies on adolescents and youth in Ghana, this study focused exclusively on school-going adolescents. By focusing exclusively on secondary school students in Ghana, it might be easier to measure the success or failure of prevention efforts against school-going adolescent mortality and morbidity as well as enhance ongoing health education programmes and policies within the context of schools.

## **METHOD**

### *Data collection procedures and sample*

The setting for this study was Accra. With a population of 1.67 million, the Accra metropolitan area is the most urbanized region in Ghana (14). Data for the study were collected during a five-week period—from July to August 2000. Two schools were randomly selected from a list of senior secondary schools in Accra. The two schools, labelled 'A' and 'B', have three grade levels—SSS1 (10<sup>th</sup> grade), SSS2 (11<sup>th</sup> grade), and SSS3 (12<sup>th</sup> grade). School A enrolled more students than School B (1863 vs. 1301), but the gender distribution at the two schools was comparable (59.9% girls in School A vs. 58.1% girls in School B). Slightly more residential students were enrolled in School A than in School B (24.2% vs. 19.3%). School A is located in one of the best residential communities in Accra, and its buildings and campus are more attractive. Administrators at the two schools indicated that students in School A have higher socio-economic backgrounds. Initially, a sample of 1000 students from both schools was randomly selected based on the number of classes in each grade. With a response rate of 89.4%, the sample size was 894 students from both schools.

### *Instrument*

Because of its established reliability and extensive use in the United States and other countries, a slightly modified version of the 1999 Youth Risk Behavior Survey (YRBS) questionnaire, an instrument which was developed by the Centers for Disease Control and Prevention (CDC), was used to collect data for the study<sup>15, 16</sup>. The survey included questions on tobacco use, alcohol use, marijuana use, sexual activity, condom use, and other risk behaviours. In this article, questions dealing with family communication, sexual activity, and condom use are analysed. Ghana Education Service (GES), a unit of the country's Ministry of Education, reviewed the questionnaire and approved the survey. Administrators at the two selected schools also reviewed the questionnaire and endorsed the Ministry's approval for the survey to be administered.

The survey was administered with the assistance of a female graduate student from a local university. Completing the survey was voluntary, and respondents were advised of their right to decline participation at any time. In addition, respondents were advised that they could refuse to answer or leave blank any question they found uncomfortable. The questionnaires were not precoded to ensure anonymity of students.

### **Measures**

*Sexual activity.* Students were asked whether they have ever had sexual intercourse, with yes = 1, no = 0 response options. Previous studies<sup>17</sup> have distinguished between

sexual experience (a student who has had sexual intercourse at least once in her or his lifetime) and current sexual activity (a student who has had sexual intercourse in the past three months). The present analysis includes only sexual experience.

*Condom use.* Students were asked whether they or their partners used a condom during the last time they had sexual intercourse. The two response options were yes = 1, no = 0. In Ghana, the main mode of transmission of HIV is sexual intercourse<sup>2,18</sup>. Therefore, the rate of condom use among adolescents is a key determinant of exposure to sexual risks and it is included in the present study.

*Student-family communication.* Students were asked whether they have ever talked about HIV/AIDS with their parents or other adults in their family. The response options were dichotomised as yes = 1, no = 0. Other adults are included in family communication because of the role of the extended family in the social structure of Ghana<sup>19</sup>.

*Gender.* The question, 'What is your sex?' measured this variable. The coding categories were girls = 1, boys = 0. Previous studies have noted the need to examine gender differences in sexual activity and sexual risk behaviours, especially among adolescents and youth<sup>20</sup>.

*Age.* Asking students to respond to the question, 'What is your age?' measured this variable. Age is measured as a continuous variable.

*Grade level.* The three grade levels were coded into a dummy variable (SSS1—10<sup>th</sup> grade = 0; SSS2—11<sup>th</sup> grade = 1; SSS3—12<sup>th</sup> grade = 2).

*School.* A dummy variable was created for the two schools to test whether the schools differed significantly in relationship to the outcome variables. The coding categories were School A = 1, School B = 0.

### **Data analysis**

The data were analysed using *SPSS for Windows Version 11.0* software<sup>21,22</sup>. Descriptive analyses were conducted by calculating the frequency distribution of gender and other study variables. To test for statistically significant associations between sexual activity, condom use, and family communication about HIV/AIDS and gender, a series of chi-

square analyses were conducted. A second set of chi-square analyses was done to test the association between the three variables and school of attendance. With two dichotomous dependent variables, logistic regression was used to estimate the effect of family communication and demographic variables on sexual activity and condom use.

The analyses for the two dependent variables were done in three steps. In step 1, the effect of only family communication on the dependent variables was assessed. Next, in step 2, gender and age were added as controls to the first model. These two key demographic variables are important because previous studies have reported differential effects of communication on sexual behaviour among adolescents, depending on their gender and age<sup>9,10,12</sup>. Finally, in step 3, grade level and school of attendance were added to the second model. In steps 2 and 3, the effect of family communication on the outcome variables and controlling for the added variables were examined. The results are presented as odds ratios, which estimate the effect on the odds of the outcome variable for each unit change in the independent variable. For dichotomous independent variables, odds ratios measure the effect of being in one category (girls) compared to another (boys). If the odds ratio is less than 1, it indicates a decrease in the odds of the outcome; odds ratios greater than 1 indicate an increase in the outcome; and odds ratios equal to 1 indicate no effect<sup>23</sup>. Diagnostics tests did not indicate multicollinearity among the variables. The alpha level to determine significance for all statistical tests was set at .05.

## **RESULTS**

### **Descriptive analysis**

Before presenting the results of the multivariate logistic regression models, it is instructive to present descriptive information about the sample and variables. The sample was composed of 56.9% girls and 43.1% boys, which is representative of the gender distribution of the student population at the two schools. The mean age of respondents was 17.4 years (SD = 1.40). Overall, 25% reported that they have ever had sexual intercourse (i.e., they are sexually experienced). Of those who were sexually experienced, 25.7% had their first sexual intercourse at age 11 years or younger; and by 16 years, 64.7% had had their first sexual intercourse. The descriptive analyses also examined condom use among the respondents who are sexually experienced. The finding indicates that 55.7% of the students responded that they or their partners did not use a condom at last sexual intercourse.

In Table 1, the results of chi-square analyses of sexual activity, condom use, and family communication by gender are presented. Table 1 shows that boys were more likely than girls to have ever had sexual intercourse,

that is, to be sexually experienced (odds ratio = 2.44; 95% CI = 2.44-4.74). Table 1 also shows significant gender

differences in whether respondents or their partners used a condom at last sexual intercourse. Boys were less likely than girls to report condom use (odds ratio = 2.2.7; 95% CI = 1.22-4.27). Furthermore, boys were less likely than girls to have ever talked about HIV/AIDS with their parents or other family members (odds ratio = 2.48; 95% CI = 1.75-3.52).

**Table 1 Crosstabulation of Sexual Activity, Condom Use, and Family Communication about HIV/AIDS, by Gender**

Variable	Gender		Odds ratio	95% CI	pValue
	Girls n(%)	Boys n(%)			
Ever had sexual intercourse					
Yes	76(15.1)	143(37.7)	3.40	2.44-4.47	0.0000
No	426(84.9)	236(62.3)			
Condom use*					
Yes	40(57.1)	51(37.0)	2.2.7	1.22-4.27	0.006
No	30(42.9)	87(63.0)			
Talk about HIV/AIDS†					
Yes	346(81.0)	196(63.2)	2.48	1.75-3.52	0.000001
No	81(19.0)	114(36.8)			
Total %	(100.0)	(100.0)			

Note.

CI = Confidence intervals

\* To determine whether or not there is an association between sexual activity, condom use, family communication, and school of attendance, additional chi-square analyses were conducted. The results, presented in Table 2, indicate that only sexual activity show statistically significant association with school of attendance. Students in School B reported being more sexually experienced than were students in School A (Odds ratio = 2.6; 95% CI = 1.87-3.64).

**Table 2 Crosstabulation of Sexual Activity, Condom Use, and Family Communication about HIV/AIDS, by School of Attendance**

Variable	School of Attendance		Odds ration	95% CI	pValue
	School A n(%)	School B n(%)			
<b>Ever had sexual intercourse</b>					
Yes	72(16.3)	150(33.6)	2.61	1.87-3.64	0.0000
No	371(83.7)	296(66.4)			
Condom use*					
Yes	32(46.4)	61(43.3)	1.13	0.61-2.11	0.6700
No	37(53.6)	80(56.7)			
Talk about HIV/AIDS†					
Yes	266(71.1)	281(76.2)	1.30	0.92-1.82	0.1199
No	108(28.9)	88(23.8)			
Total %	(100.0)	(100.0)			

Note.

CI = Confidence intervals

\* Includes only respondents who have ever had sexual intercourse.

† Includes only respondents who have heard about HIV/AIDS.

### Multivariate analysis

Results from the logistic regression analyses of family communication on sexual activity and condom use are presented in Table 3. Columns 1, 2, and 3 show the relative odds that students were sexually experienced. Column 1 shows that family communication about HIV/AIDS by itself has no significant influence on being sexually experienced. Given that prior research has noted the importance of demographic characteristics in sexual activity, in column 2, gender and age are added as controls. The results indicate that girls are less likely than boys to be sexually experienced (OR = .28). That is, the odds of being sexually experienced are 72% less likely for girls than boys. Column 2 also shows that age is associated with being sexually experienced. For a

year's increase in age, the odds of being sexually experienced increase by a factor of 1.37. Controlling for gender and age, however, does not affect the effect of family communication on sexual experience, which remains insignificant. In column 3, grade level and school of attendance are included as additional controls. The results show that being sexually experienced is associated with the school a student attends (OR = .42), but not with grade level. Respondents who attend School A are 58% less likely than students who attend School B to be sexually experienced. Column 3 also shows that, controlling for these factors hardly affected the effect of gender and age on being sexually experienced. Furthermore, controlling for all four factors did not affect the effect of family communication about HIV/AIDS on being sexually experienced, which remains insignificant.

**Table 3 Odds Ratios of Logistic Regression of Lifetime Sexual Activity and Condom Use Among Senior Secondary School Students in Accra, Ghana, by Family Communication and Demographic Characteristics**

Variable	Ever Had Sexual Intercourse			Used Condom at Last Sexual Intercourse <sup>a</sup>		
	(1)	(2)	(3)	(4)	(5)	(6)
Talked about HIV/AIDS						
No (ref.)	—	—	—	—	—	—
Yes	.70	.92	.85	2.21*	2.20*	2.23*
Gender						
Boys (ref.)		—	—		—	—
Girls		.28*	.30*		2.62*	2.72*
Age (in years)		1.37*	1.43*		1.09	1.24
Grade						
10 <sup>th</sup> —SSS1 (ref.)			—			—
11 <sup>th</sup> —SSS2			1.50			2.29
12 <sup>th</sup> —SSS3			1.49			1.46
School of attendance						
School B (ref.)			—			—
School A			.42*			1.19*
(-2) Log Likelihood	789.8	691.6	670.2	212.1	198.0	195.2
Model Chi-Square <sup>†</sup>	3.5	85.2	106.6	5.0	12.8	15.6
d.f.	1	3	6	1	3	6
N	741	728	728	159	155	155

Note. <sup>a</sup>Includes only students who have ever had sexual intercourse.

\*  $p < .05$

<sup>†</sup> All model chi-squares are significant at  $p < .05$

Columns 4, 5, and 6 show the relative odds of using a condom at last sexual intercourse. The results in column 4 indicate that respondents who have ever talked about HIV/AIDS with parents or other family members are significantly more likely than those who have never talked with parents or other family members to report condom use at last sexual intercourse (OR = 2.21). In column 5, gender and age are added as controls. The results indicate that condom use is significantly associated with gender, but not with age. Girls were significantly more likely than boys to report condom use at last sexual intercourse (OR = 2.62). Controlling for the two variables appear to have little effect on the association between family communication and condom use. Finally, in column 6, grade level and school of attendance are added as additional controls to column 5. As indicated in column 6, condom use is significantly associated with school of attendance, but not with grade level. The odds of using a condom at last sexual intercourse is greater for respondents who attend School A than for those who attend School B (OR = 1.19). By comparing columns 5 and 6, the results show that the addition of grade and school of attendance hardly affected the effect of family communication and gender on condom use.

## DISCUSSION

This study examined sexual activity and condom use and the extent to which they are affected by family communication about HIV/AIDS among a sample of senior secondary school students in Accra, Ghana. The findings of the study are noteworthy. A quarter of the students had engaged in sexual intercourse, with boys more likely to do so than girls. Similarly, a quarter of the sexually experienced students initiated sexual intercourse when they were 11 years old or younger, and by the age of 16, over 64 percent had already initiated sexual activity. These findings suggest that information about sexual activity must be conveyed to students early.

The findings also indicated that the sexually experienced students are at great risk of exposure to HIV and other sexually transmitted diseases. It is notable that only 44.3% of the sexually experienced students reported that they or their partners did use a condom at last sexual intercourse. Adih and Alexander (1996) reported a much lower percent of condom use at last sexual intercourse in their study of youth in Ghana<sup>3</sup>. The present

finding suggests the urgent need for students to receive information about condom use before initiating sexual intercourse. In addition, considering that 25.7% of the sexually experienced students had initiated first sexual intercourse at age 11 years or younger, this information must be given at an early age.

The most important finding of this study is the association between family communication about HIV/AIDS and condom use. This finding is consistent with previous studies that have shown the importance of family communication about sexual issues<sup>9</sup>. The present result is encouraging because the increased use of condoms attributable to family communication is likely to reduce sexually related risks among sexually active school-age adolescents in Ghana. It may, however, be necessary in future research to build upon the present finding and examine additional aspects of family communication, especially the role of fathers and mothers. The assessment of each parent's role in family communication is critical in a society such as Ghana, where traditional norms greatly inhibit discussions of sexual issues with parents, especially with fathers.

Another important finding is the significant association between sexual activity, condom use, and school of attendance. Students attending School A, which has better facilities were less likely to be sexually experienced but more likely to use a condom. Differences in socio-economic background of students at the two schools may be one plausible explanation. Another possible explanation may be due to the content and process of family communication, which cannot be determined from the data of the present study. It would be insightful for future research to collect data on these factors in order to better explain the role of school of attendance in sexual behaviours.

### *Implications of the study*

The findings of this study have implications for health educators and health care practitioners seeking to better understand the prevalence of sexual activity and risk behaviours among school-age children in Ghana and perhaps other sub-Saharan African countries. The percentage of sexually experienced students who are not using condoms at sexual intercourse suggests that current educational efforts and strategies are inadequate. The findings of this study suggest that school-based interventions programmes must involve parents because of the strong association between family communication and condom use.

### *Limitations of the study*

The implications of the study, however, need to be

examined within the context of the study's limitations. First, as an exploratory study, it was limited to two senior secondary schools in Accra. Being the capital city of Ghana, life in Accra is different from life in peri-urban and rural areas of the country. At the two selected schools, 77.8% of the students were non-residential. By contrast, schools in peri-urban and rural areas of the country tend to have higher percentages of boarding (residential) students. Therefore, the student population from which the sample was drawn may differ from the population of students in schools located in peri-urban and rural areas of the country in terms of contextual factors. Second, the data analysed for this study are cross-sectional; therefore, no definitive statements regarding the causal relations between family communication and the outcome variables can be made.

## CONCLUSION

These limitations notwithstanding, the findings of this study contribute significantly to the growing literature on adolescents in Ghana. The next logical step for future research is to use the YRBS questionnaire to conduct a nationwide survey of junior and secondary school students in Ghana. The addition of junior secondary schools to the population of schools in this study is critical because of the reported early initiation of sexual intercourse. Furthermore, the inclusion of students from peri-urban and rural schools will address one of the limitations of the present study. In addition, there is a need to examine the content of discussions between students and family members as well as the extent to which outcomes of such discussions are dependent upon the sex of students and family members. In conclusion, the findings of this study should provide baseline data to which future surveys using the YRBS questionnaire may be compared.

## ACKNOWLEDGEMENTS

I would like to thank Vida Benson for her assistance in data collection.

## REFERENCES

1. Young people and HIV/AIDS: Opportunity in crisis. UNAIDS web site: <http://www.unaids.org/barcelona/presskit/youngpeople/index.html>. Accessed 6 September 2002.
2. Ghana: Joint Country Mission Report. UNAIDS (International Partnership against AIDS in Africa) web site: [http://www.unaids.org/hivaidinfo/statistics/fact\\_sheets/pdfs/Ghana\\_en.pdf](http://www.unaids.org/hivaidinfo/statistics/fact_sheets/pdfs/Ghana_en.pdf). Accessed 6 September 2002.
3. Adih WK, Alexander CS. Determinants of condom use to prevent HIV infection among youth in Ghana. *J Adolesc Health* 1999; 24: 63-72.
4. Agyei WKA, Biritwum RB, Ashitey AG, Hill RB. Sexual behaviour and contraception among unmarried adolescents and young adults in Greater Accra and Eastern regions of Ghana. *J Biosoc Sci* 2000; 32: 495-512.
5. Anarfi JK. Vulnerability to sexually transmitted disease: Street children in Accra. *Health Transition Review* 1997; 7(Suppl): 281-306.
6. Ankomah A. Condom use in sexual exchange relationships among young single adults in Ghana. *AIDS Educ Prev* 1998; 10: 303-16.
7. Awusabo-Asare K, Abane AM, Badasu DM, Anarfi JK. 'All die be die': Obstacles to change in the face of HIV infection in Ghana. In Caldwell JC, Caldwell P, Anarfi J, Awusabo-Asare K, Ntozi J, Orubuloye IO et al., editors. Resistances to behavioural change to reduce HIV/AIDS infection in predominantly heterosexual epidemics in third world countries. Canberra: Australia National University; 1999; p.125-32.
8. Wolf CR, Tawfik LA, Bond KC. Peer promotion programs and social networks in Ghana: Methods for monitoring and evaluating AIDS prevention and reproductive health programs among adolescents and young adults. *Journal of Health Communication* 2000; 5: 61-80.
9. Holtzman D, Rubinson R. Parent and peer communication effects on AIDS-related behavior among U.S. high school students. *Fam Plann Perspec* 1995; 27: 235-68.
10. Miller KS, Kotchick BA, Dorsey S, Forehand R, Ham AY. Family communication about sex: What are parents saying and are their adolescents listening? *Fam Plann Perspec* 1998; 30: 218-35.
11. Shoop DM, Davidson PM. AIDS and adolescents: the relation of parent and partner communication to adolescent condom use. *J Adolesc* 1994; 17: 137-48.
12. Pick S, Palos P. Impact of the family on the sex lives of adolescents. *Adolescence* 1995; 30(119): 667-75.
13. Fisher TD. An extension of the findings of Moore, Peterson, and Furstenburg (1986) regarding family sexual communication and adolescent sexual behavior. *Journal of Marriage and the Family* 1989; 51(3): 637-39.
14. Ghana Statistical Service. 2000 Population & Housing Census: Provisional Results. Accra (Ghana): Ghana Statistical Service; 2000.
15. Brener ND, Collins JL, Kann L, Warren CW, Williams BI. Reliability of the Youth Risk Behavior Survey questionnaire. *Am J Epidemiol* 1995; 141(6): 575-80.
16. Centers for Disease Control and Prevention. 1999 National School-Based Youth Risk Behavior Survey (YRBS). Centers for Disease Control and Prevention (CDC), Atlanta, GA: <http://www.cdc.gov/nccdphp/dash/yrbs/data/1999/yrbs1999.pdf>

17. Kann L, Kinchen SA, Williams BI, Ross JG, Lowry R, Grunbaum JA et al. Youth risk behavior surveillance—United States, 1999. *J Sch Health* 2000; 70: 271-85.
18. Neequaye AR, Neequaye JE, Biggar RJ. Factors that could influence the spread of AIDS in Ghana, West Africa: Knowledge of AIDS, sexual behaviour, prostitution, and traditional medical practices. *J Acquir Immune Defic Syndr* 1991; 4: 914-19.
19. Assimeng M. Social structure of Ghana: A study in persistence and change. 2<sup>nd</sup> ed. Tema (Ghana): Ghana Publishing Corporation; 1999.
20. World Health Organization. Preventing HIV/AIDS/STI and related discrimination: An important responsibility of health-promoting schools. WHO Information Series on School Health (WHO/SCHOOL/98.6). World Health Organization Web site: <http://www5.who.int/school-youth-health/main.cfm>? Accessed 5 May 2000.
21. SPSS Inc. SPSS for Windows, Release. 11.0.1. Chicago: SPSS Inc; 2001.
22. Norusis MJ. SPSS 11.0: Guide to data analysis. Upper Saddle River (NJ): Prentice Hall; 2002.
23. DeMaris A. Logit Modeling: Practical Applications. Sage University Paper Series on Quantitative Applications in the Social Sciences, 07-086. Newbury Park (CA): Sage; 1992.