

Exclusive breastfeeding practices in the Coast region, Tanzania.

Method Kazaura

School of Public Health, Epi/Biostat

Abstract

Background: Although breastfeeding in general is common and culturally accepted in many sub-Saharan countries, recommended exclusive breastfeeding infants to 6 months is rare. In rural Tanzania, data on infant feeding practices is rare.

Objective: To examine and describe exclusive breastfeeding practices in rural settings (Coast Region) of Tanzania.

Methods: A cross-sectional study was conducted in Coast Region of Tanzania involving 342 mothers. Only mothers with children aged between 6 and 23 months were interviewed in their residences. Data analyses included descriptive and logistic regression analyses.

Results: The majority, 66%, of mothers reported to have breastfed their new born within the first hour of life. About 30% reported to have breastfed exclusively for up to at least six months. Those who did not practice complete exclusive breastfeeding mentioned insufficient milk as the main reason. Correlates of exclusive breastfeeding included maternal education and attitudes towards exclusive breastfeeding.

Conclusion: The rate of exclusive breastfeeding in rural areas like the Coast Region of Tanzania is still very low. Programs aimed to promote exclusive breastfeeding must take multi-factorial considerations.

Keywords: Exclusive breastfeeding, practice, rural, Tanzania

DOI: <http://dx.doi.org/10.4314/ahs.v16i1.6>

Cite as: Kazaura M. *Exclusive breastfeeding practices in the Coast Region, Tanzania.* *Afri Health Sci.* 2016;16(1): 44-50. <http://dx.doi.org/10.4314/ahs.v16i1.6>

Introduction

Exclusive breastfeeding (EBF) is defined as giving breast milk only and no other liquids, except drops or syrups with vitamins, mineral supplements or medicines. The World Health Organization (WHO) recommends that newborns and infants should be fed only with breast milk within the first hour after birth until 6 months of life; therefore practice exclusive breastfeeding for the first six months of life^{1,2}. Various benefits for practicing EBF that include nutritional, social, economic, environmental and health, nutritional, psychological and developmental have been outlined and discussed extensively elsewhere³⁻⁷.

Although breastfeeding in general is common and culturally accepted in many sub-Saharan countries⁸, recommended EBF of infants to 6 months of life is rare⁹⁻¹¹. Reported rates of EBF especially in developing countries are still low¹². Epidemiological studies show that in most of the low and middle income countries, the rate of EBF for infants less than six months is around

30%¹³. A recent report in Tanzania indicated 17% of infants less than 6 months of age were predominantly breastfed but rare in rural areas¹⁴. The national duration average of EBF is 2.4 months being longer in rural areas than in urban areas (2.5 and 1.9 months respectively)¹⁵.

Practices of feeding infants during the first six months and beyond are diverse based on geographical, economic and cultural settings. However, the main concerns are the time when mothers initiate breastfeeding, the duration of breastfeeding and the age at which infants are weaned¹⁶.

In Tanzania, enough data on infant feeding practices is available either in urban settings or in relation to HIV. The gap is on the situation about breastfeeding practices in rural settings of Tanzania. Therefore, we conducted a study to examine and describe EBF practices in Coast Region which is a rural setting of Tanzania.

Methodology

A cross-sectional study was conducted in a rural setting (Coast Region) of Tanzania. The Region was projected to have a rural total population of 844,643 (74.4% of total region population). Furthermore, infants aged between 0 and 2 years were estimated to be 85,684¹⁷.

The study targeted mothers of children aged at least 6 months but less than 2 years (23 months) at their res-

Corresponding author:

Method Kazaura,
School of Public Health, Epi/Biostat
Email: mrkazaura@muhas.ac.tz

idences. Using sample size calculator for a single proportion (proportion of mothers breastfeeding exclusively was estimated to be 30%), we estimated a sample of 350 mothers to be sufficient for the study. Sampling was based on a random selection of wards (an administrative unit comprising of villages (in rural settings) or streets (in urban settings). Selected wards were Kibiti, Ikwiriri, Chalinze, Bagamoyo, Zinga and Mlingotini. In each of the selected wards, one village was selected at random from which all households with mothers having a living child aged at least 6 months but less than 24 months and present at the time of the survey were interviewed. For a mother with more than one child aged between the desired age ranges, the youngest child was selected. We excluded multiple births, for example twins. Interview schedules were used to capture data. The study tool included demographic information of the mother and the child, mother's attitude towards EBF and feeding practices of the selected child during the first 6 months of life. Recruitment into the study and interviews of all study participants lasted for four working days.

An oral informed consent was acquired from each mother before initiating the interview. Privacy and confidentiality were observed throughout the study so that participants were comfortable when responding to the questions and none was identified by names rather by code numbers necessary during data entry, cleaning and further processing. Permission to implement the study was sought from the district administrative and health authorities.

Reported feeding practices were based on mother's ability to recall their experiences during the past 6 to 23 months. There was no missing information for the main variable (case-wise, outcome variable) but for the

independent, like background variables, no attempts were made to adjust for missing information before cross-tabulations. The analyses were performed using the Statistical Package for Social Sciences (SPSS) software (Version 20).

We assessed attitudes towards exclusive breastfeeding on the basis of 16-items; each rated on a five-point Likert scale (1 = strongly agree; 5 = strongly disagree). Therefore, during data processing, a respondent could potentially score from 16 to 80 points. Total low scores, indicated positive attitudes and total high scores suggested negative attitudes. A respondent scoring below the average was considered to have a positive attitude and those scoring above the average were considered to have a negative attitude. In assessing the overall internal consistency, we used Cronbach's alpha and we found a moderate reliability (Cronbach's alpha=0.54).

In order to assess predictors of EBF, we first run bivariate analysis (with a Chi-square test) between the outcome variable (mothers practicing EBF) and possible independent variables. Variables whose p-values were at most 0.20 in the bivariate analyses were included as explanatory variables in the binary logistic regression models. Using the logistic regression analysis, we estimated the odds of practicing EBF with 95% confidence intervals to assess the strengths of association.

Results

Characteristics of the sample

We recruited 342 (97.7%) mothers out of the original estimated sample size. Their mean age was 27.2 (SD=6.7) years. The majority, 126 (36.8%), either never attended school or had incomplete primary education. Almost half, 169 (49.4%) of these mothers were peasants and two thirds, 228 (66.7%), reported were currently married (Table 1).

Table 1: Distribution of mothers by their background characteristics (n= 342)

| Characteristics | Number (%) |
|-----------------------------------|------------|
| Maternal age group (years) | |
| 15 – 19 | 42 (12.3) |
| 20 – 24 | 90 (26.3) |
| 25 – 29 | 86 (25.1) |
| 30 – 34 | 65 (19.0) |
| 35 + | 59 (17.3) |
| Level of Education | |
| None/Incomplete primary | 126 (36.8) |
| Primary education | 175 (51.2) |
| Above primary education | 41 (12.0) |
| Marital status | |
| Single | 68 (19.9) |
| Married/Cohabiting | 250 (73.1) |
| Divorced/Widow | 24 (7.0) |
| Occupation | |
| Subsistence farming | 169 (49.4) |
| Unemployed | 85 (24.9) |
| Formally employed | 13 (3.8) |
| Petty business | 75 (21.9) |


With their mothers, 172 (50.3%) children were girls. The mean age of all children was 13.0 (SD = 4.8) years. Mothers of nulligravida, gravida 1 or gravida 2 were 98 (28.7%), 94 (27.5%) and 56 (16.1%). The remaining 94 (27.8%) were above gravida 2.

Breast feeding practices

Breastfeeding initiating

The majority, 227 (66.3), reported to have breastfed the newborn within an hour of delivery whereas only 5 (1.5%) mothers did not breastfeed the baby at all (Figure 1). (Table 1)

Figure 1. Time (hours) taken before breastfeeding the newborn after delivery




Exclusive breastfeeding

Of the total mothers interviewed, 100 (29.2%) reported to have breastfed their infants exclusively up to six

months. Only 6% of children aged 24 months were breastfed exclusively at least 23 months. The rate of breastfeeding exclusively decreased dramatically from birth to about 9 months (Figure 2).

Figure 2. Proportion of children under the age of 24 months breastfed exclusively



Reasons for not breastfeeding exclusively

Of the mothers who did not exclusively breastfeed their children or who interrupted exclusive breastfeeding, 158 (68.7%) alleged that breast milk was insufficient, 39 (16.8%) were advised by relatives to immediately begin complementary feeding and 31 (16.6%) reported stopping due to work commitments. The main diet during this period was maize or cassava porridge reported by 225 (94.6%) mothers.

(Figure 1) (Figure 2)

Predictors of exclusive breastfeeding

Table 2 presents results from binary logistic regression

analysis with EBF being the dependent variable and three independent variables (education level, occupation and attitude towards EBF). The proportion of mothers reporting EBF increased with increasing level of maternal education. Mothers who attained above primary education had almost six odds to exclusively breastfeed for six months as compared to those with incomplete or no primary education (AOR=5.5; 95% CI = 2.2, 13.6). Furthermore, mothers with positive attitudes towards EBF had twice the odds, (AOR = 2.1; 95% CI = 1.2, 3.6), to practice EBF up to six months after birth as compared to mothers with negative attitudes. (Table 2)

Table 2. Correlates of Exclusive Breastfeeding

| Characteristic | Number EBF (%) | OR (95% CI)* | |
|-----------------------------|----------------|-----------------|-----------------|
| | | Unadjusted | Adjusted |
| Education level | | | |
| None/some primary | 21 (16.7) | Reference | Reference |
| Completed primary | 58 (33.1) | 2.5 (1.4, 4.4) | 2.0 (1.1, 3.6) |
| Above primary | 23 (56.1) | 6.4 (2.9, 13.9) | 5.5 (2.2, 13.6) |
| Occupation | | | |
| Peasant | 38 (22.5) | Reference | Reference |
| Employed | 3 (23.1) | 2.1 (1.2, 3.7) | 1.8 (1.0, 3.3) |
| Petty business | 29 (38.7) | 2.2 (1.2, 3.9) | 1.2 (0.6, 2.4) |
| Unemployed | 32 (37.6) | 1.0 (0.3, 3.9) | 0.3 (0.1, 1.5) |
| Attitude towards EBF | | | |
| Negative | 30 (19.0) | Reference | Reference |
| Positive | 72 (39.1) | 2.7 (1.7, 4.5) | 2.1 (1.2, 3.6) |

* Odds ratio (95% Confidence Interval)

Discussion

Despite strong initiatives to promote breastfeeding in general and EBF in particular, the proportion of women EBF has remained low¹⁸⁻²⁰. In this study, less than 30% of women reported to breastfeed infants exclusively for up to six months. This level is lower than the 50% reported national average¹⁵. However, similar or lower proportions have been reported from middle- and lower-income countries specifically from the sub-Saharan African countries²¹⁻²⁴. More than two-thirds, (69%), reported insufficient milk to be the main reason for not breastfeeding exclusively; a reason that has been cited earlier elsewhere²⁵⁻²⁶.

In this study, education level of mothers and attitude towards exclusive breastfeeding were identified to be independent predictors of exclusive breastfeeding. Although, to the best of our search, we could not get literature citing independent relationship between EBF in rural sub-Saharan Africa areas with maternal education, a study in Canada indicated such association²⁷. Furthermore, few studies from high-income countries have documented maternal attitudes to be independent predictors of EBF²⁸⁻³⁰.

Findings from this study must be interpreted with caution bearing in mind that this was a cross-sectional study. Therefore, it is difficult to make inference about the causality. Secondly, responses might have been limited to the ability to remember breastfeeding outcomes of 6 to 24 months before the survey. However, as re-

ported in the past, it is optimistic that mothers always remember accurately the duration of breastfeeding their infants³¹. Thirdly, although inference made from may apply to this region (Coast) only, cultural differences and other factors that may influence EBF practices between rural populations in Tanzania may not vary significantly. Nevertheless, we recommend additional qualitative data to explore factors for EBF in rural areas. Last but not least, although we used interviewers with social/medical background, we are not able to assess how these attributes influenced the respondent's opinion in describing the breast-feeding practices and the possibility of respondents offering socially desirable answers causing desirability bias. However, because of intensive training of research assistants and having control questions in the tool, if such biases existed, they were very minimal and random.

Conclusion

The rate of EBF in rural areas like the Coast Region in Tanzania is still low. In order to enhance comprehensive EBF during the first six months of the baby, programs aimed to promote EBF must take multi-factorial considerations.

Conflict of interest

None declared.

Funding

The study was supported by Muhimbili University of Health and Allied Sciences.

References

1. World Health Organization. Global strategy for infant and young child feeding. Geneva: World Health Organization, 2003
2. World Health Organization. Planning guide for national implementation of the global strategy for infant and young child feeding. 2006. http://www.who.int/nutrition/publications/Planning_guide.pdf.
3. Rockville, MD; US Department of Health and Human Services; 2007. Jones G, Steketee R, Black R, Bhutta Z, Morris S, the Bellagio Child Survival Study Group: How many child deaths can we prevent this year? *Lancet* 2003, 362:65-71
4. Bhandari N, Bahl R, Mazumdar S, Martines J, Black RE, Bhan MK: Effect of community-based promotion of exclusive breast-feeding on diarrhoeal illness and growth: a cluster randomized controlled trial. *Lancet* 2003, 361: 1418-1423
5. Arifeen S, Black RE, Antelman G, Baqui A, Caulfield L, Becker S: Exclusive breastfeeding reduces acute respiratory infection and diarrhea deaths among infants in Dhaka slums. *Pediatrics* 2001, 108:E67
6. Andrew N, Harvey K. Infant feeding choices: experience, self-identity and lifestyle. *Matern Child Nutr* 2011;7:48-60
7. American Academy of Pediatrics: Breastfeeding and the Use of Human Milk. Section on Breastfeeding. *Pediatrics* 2005; 115:2 496-506; doi:10.1542/peds.2004-2491
8. Dop MC. Breastfeeding in Africa: will positive trends be challenged by the AIDS epidemic? *Sante*. 2002;12:64-72
9. de Paoli M, Manongi R, Helsing E, Klepp KI. Exclusive breastfeeding in the era of AIDS. *J Hum Lact*. 2001;17:313-20
10. Shirima R, Gebre-Medhin M, Greiner T. Information and socioeconomic factors associated with early breastfeeding practices in rural and urban Morogoro, Tanzania. *Acta Paediatrica*. 2001;90:936-42
11. Agnarsson I, Mpello A, GunnLaugsson G, Hofvander Y, Greiner T. Infant feeding practices during the first six months of life in a rural area in Tanzania. *East Afr Med J*. 2001;78:9-13
12. Arabi M, Frongillo EA, Avula R, Mangasaryan N. Infant and young child feeding in developing countries. *Child Dev*. 2012;83(1):32-45. doi: 10.1111/j.1467-8624.2011.01675.x
13. Bhutta ZA, Ahmed T, Black RE, Cousens S, Dewey K, Giugliani E, Haider BA, Kirkwood B, Morris SS, Sachdev HP, et al. Maternal and Child Undernutrition Study Group. What works? Interventions for maternal and child undernutrition and survival. *Lancet*. 2008;371:417-40
14. Victor R, Baines SK, Agho KE, Dibley MJ. Determinants of breastfeeding indicators among children less than 24 months of age in Tanzania: a secondary analysis of the 2010 Tanzania Demographic and Health Survey. *BMJ Open*. 2013;3(1). doi:pii: e001529. 10.1136/bmjopen-2012-001529
15. National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2010. Tanzania Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro
16. Clayton HB, Li R, Perrine CG, Scanlon KS. Prevalence and reasons for introducing infants early to solid foods: variations by milk feeding type. *Pediatrics*. 2013;13:e1108-14. doi: 10.1542/peds.2012-2265.
17. United Republic of Tanzania, 2006. Regional and District Projections. Vol XII. National Bureau of Statistics, Ministry of Planning, Economy and Empowerment. Dar es Salaam
18. WHO. HIV and infant feeding. Revised principles and recommendations. Rapid Advice. Geneva: WHO; 2009
19. WHO. New data on the Prevention of Mother-to-child Transmission of HIV and their Policy Implications: Conclusions and Recommendations. WHO Technical Consultation on behalf of UNFPA/UNICEF/WHO/UNAIDS Inter-agency Task Team on Mother-to-child Transmission of HIV. Geneva 11-13 October, 2000. Geneva: WHO; 2001. WHO/RHR/01.28
20. United Nations Children's Fund. The State of the World's Children 2011: Adolescence—An Age of Opportunity. UNICEF; New York, NY, USA: 2011
21. Ochola SA, Labadarios D, Nduati RW. Impact of counselling on exclusive breast-feeding practices in a poor urban setting in Kenya: a randomized controlled trial. *Public Health Nutr*. 2012 Oct 8:1-9.[Epub ahead of print]
22. Radwan H. Patterns and determinants of breastfeeding and complementary feeding practices of Emirati Mothers in the United Arab Emirates. *BMC Public Health*. 2013;13:171. doi: 10.1186/1471-2458-13-171
23. Oche MO, Umar AS, Ahmed H. Knowledge and practice of exclusive breastfeeding in Kware, Nigeria. *Afr Health Sci*. 2011;11:518-23
24. Faruque AS, Ahmed AM, Ahmed T, Islam MM, Hossain MI, Roy SK, Alam N, Kabir I, Sack DA. Nutrition: basis for healthy children and mothers in Bangladesh. *J Health Popul Nutr*. 2008; 26:325-39
25. Ho YJ, McGrath JM. Predicting breastfeeding dura-

- tion related to maternal attitudes in a taiwanese sample. *J Perinat Educ.* 2011 Fall;20:188-99. doi: 10.1891/1058-1243.20.4.188
26. Giugliani ER. [Common problems during lactation and their management]. *J Pediatr (Rio J).* 2004;80(5 Suppl):S147-54
27. Jessri M, Farmer AP, Maximova K, Willows ND, Bell RC. Predictors of exclusive breastfeeding: observations from the Alberta pregnancy outcomes and nutrition (APrON) study. *BMC Pediatr.* 2013;13:77. [Epub ahead of print]
28. de Jager E, Skouteris H, Broadbent J, Amir L, Mellor K. Psychosocial correlates of exclusive breastfeeding: a systematic review. *Midwifery.* 2013;29:506-18. doi: 10.1016/j.midw.2012.04.009. 2012 Oct 23. PubMed Epub.
29. de Jager E, Broadbent J, Fuller-Tyszkiewicz M, Skouteris H. The role of psychosocial factors in exclusive breastfeeding to six months postpartum. *Midwifery.* 2013 Aug 7. pii: S0266-6138(13)00213-1. doi: 10.1016/j.midw.2013.07.008. [PubMed Epub ahead of print]
30. Scott JA, Shaker I, Reid M. Parental attitudes toward breastfeeding: their association with feeding outcome at hospital discharge. *Birth.* 2004;31:125-31
31. Eaton-Evans J, Dugdale AE. Recall by mothers of the birth weights and feeding of their children. *Hum Nutr Appl Nutr.* 1986;40:171-5 sedentary.