

## Short Communication

# Proximate analysis of *Lentinus squarrosulus* (Mont.) Singer and *Psathyrella atroumbonata* Pegler

NWANZE PI<sup>1\*</sup>, JATTO W<sup>1</sup>, ORANUSI S<sup>1</sup> and JOSIAH SJ<sup>2</sup>

<sup>1</sup>Department of Biological Sciences, College of Natural and Applied Sciences, Igbinedion University, Okada, P.M.B. 0006, Edo State, Nigeria.

<sup>2</sup>Department of Biochemistry, School of Basic Medical Sciences, College of Health Sciences, Igbinedion University, Okada, P.M.B. 0006, Edo State, Nigeria.

Accepted 9 January, 2006

*Lentinus squarrosulus* and *Psathyrella atroumbonata*, two mushroom species commonly found growing on dead leaves and logs, were collected from the Zaria environ and taken to the laboratory for further studies. Each of the mushroom species was separated into its stipe and pileus and used for proximate analysis. There was a highly significant difference ( $p < 0.01$ ) in the proximate composition of the two species. *P. atroumbonata* had significantly higher crude protein, crude fibre and moisture content than *L. squarrosulus* while the reverse was the case for ash, dry matter, crude fat and soluble carbohydrates. In addition, there was a highly significant difference ( $p < 0.01$ ) in the proximate composition of the different mushroom parts. The pilei contained significantly higher amounts of crude protein, crude fibre, ash, and dry matter than the stipes while the converse was the case for moisture, crude fat and soluble carbohydrates. There was also a highly significant difference ( $p < 0.01$ ) in the interaction of species by parts.

**Key words:** *Lentinus squarrosulus*, *Psathyrella atroumbonata*, proximate analysis.

## INTRODUCTION

Mushrooms are cultured world wide for their taste, nutritional attributes and potential applications in industries (Sunagawa and Magae, 2005; D'Annibale et al., 2005; Mata et al., 2005). In addition, they have many medicinal uses and are good agents of bioremediation (Magingo et al., 2004; Lim et al., 2004; Adenipekun and Fasidi, 2005; Estévez et al., 2005).

We have worked extensively on the culture of *Lentinus squarrosulus* and *Psathyrella atroumbonata* at the hyphal level in submerged liquid cultures as well as at the carpophore level on various different media (Nwanze et al., 2004, 2005a, 2005b). In addition, we have examined the mineral content and amino acid composition of the two species (Nwanze and Adamu, 2004). The present investigations, however, center on the proximate composition of the mushrooms.

## Proximate analysis

The moisture content, dry matter, crude fat, crude protein, crude fibre, ash and soluble carbohydrate content were determined on a dry weight basis as described by Praveena et al. (2001). The percentage of crude protein in the different samples was calculated as %N x 4.32 as described by Crisan and Sands (1978) while the amount of soluble carbohydrate was calculated by difference.

## Statistics

The data obtained was analyzed by ANOVA with factorial treatment structures and interactions (Christie et al., 2001). Difference between means was compared using Duncan's multiple range test (DMRT, Snedecor and Cochran, 1987).

## RESULTS

The proximate analysis of *L. squarrosulus* and *P. atroumbonata* is depicted in Table 1 along with their first order interactions. There was a highly significant difference ( $p < 0.01$ ) in the protein, fibre, ash, moisture, dry

\*Corresponding authors E-mail: stonenwanze@yahoo.com.

**Table 1.** Proximate composition (g/100g dry matter) of the stipe and pileus of *L. squarrosulus* and *P. atroumbonata* pooled.

Treatments	% crude protein	% crude fibre	% ash	% moisture	% dry matter	% crude fat	% soluble carbohydrate
<b>Species</b>							
<i>L. squarrosulus</i>	22.82b	7.64b	7.52b	2.76b	97.25a	6.29a	60.65a
<i>P. atroumbonata</i>	30.20a	9.71a	17.76a	5.46a	94.55b	3.05b	51.59b
Significance	**	**	**	**	**	**	**
SE±	0.22	0.09	0.04	0.05	0.06	0.06	0.27
<b>Parts</b>							
Pileus	29.82a	9.31a	13.06a	4.01b	96.01a	4.53b	52.40b
Stipe	23.19b	8.05b	12.22b	4.21a	95.79b	4.80a	59.84a
Significance	**	**	**	**	**	**	**
SE±	0.22	0.09	0.04	0.05	0.06	0.06	0.27
<b>Interactions</b>							
S X P	**	**	**	**	**	**	**

Means followed by the same letter(s) within a treatment group are not significantly different statistically at 5% level of probability using DMRT.

\* and \*\* = significant at 5% and 1% levels, respectively; NS = not significant.

**Table 2.** The proximate composition (g/100g dry matter) of *L. squarrosulus* and *P. atroumbonata* parts

Mushroom species and parts	% crude protein	% crude fibre	% ash	% moisture	% dry matter	% crude fat	% Soluble carbohydrate
<i>L. squarrosulus</i> stipe	18.32c	6.80d	6.62c	3.83c	96.18b	6.01b	65.07a
<i>L. squarrosulus</i> pileus	27.25b	8.48c	8.42b	1.7d	98.33a	6.56a	56.23b
<i>P. atroumbonata</i> stipe	28.00b	9.30b	17.82a	4.60b	95.4c	3.59c	54.61c
<i>P. atroumbonata</i> pileus	32.40a	10.14a	17.70a	6.33a	93.7d	2.51d	48.56d
SE±	0.31	0.13	0.05	0.07	0.08	0.09	0.38

Means followed by the same letter(s) within a treatment group are not significantly different statistically at 5% level of probability using DMRT.

\* and \*\* = significant at 5% and 1% levels, respectively; NS = not significant.

matter, fat and soluble carbohydrate content of the two mushroom species. Analysis of the data showed that *P. atroumbonata* had significantly higher crude protein, crude fibre and moisture content than *L. squarrosulus* while the reverse was the case for ash, dry matter, crude fat and soluble carbohydrates.

The proximate composition of the stipe and pileus of *L. squarrosulus* and *P. atroumbonata* is shown in Table 2. There was a significant difference ( $p < 0.01$ ) in the means due to the protein, fibre, ash, moisture, dry matter, crude fat and soluble carbohydrate content of the different mushroom parts. The pileus of *P. atroumbonata* contained a significantly larger amount of crude protein than the corresponding stipe as well as the pileus of *L. squarrosulus*, both of which had similar means, but nonetheless contained significantly more crude protein than the stipe of *L. squarrosulus*. The pileus of *P.*

*atroumbonata* contained a significantly larger amount of crude fibre than the corresponding stipe followed by the pileus and stipe of *L. squarrosulus*. The ash content of the pileus and stipe of *P. atroumbonata* were at par, but of greater magnitude than the ash content of the pileus and stipe of *L. squarrosulus*. The pileus of *L. squarrosulus* contained a significantly larger amount of dry matter and crude fat than the corresponding stipes. However, it was the stalk of *P. atroumbonata* that contained a significantly larger amount of dry matter and crude fat than the corresponding pileus. There was a significantly higher amount of soluble carbohydrates in the stipe of *L. squarrosulus* than in its pileus. The soluble carbohydrate content of the stipe of *P. atroumbonata*, though significantly lower compared to the stipe of *L. squarrosulus*, contained a significantly larger amount of soluble carbohydrate than its corresponding pileus.

## DISCUSSION

Of the two species examined in this investigation, *P. atroumbonata* had the higher protein content. The protein content obtained for *P. atroumbonata* agreed with the value obtained by Aletor (1995) while the value for *L. squarrosulus* was slightly higher. However, the crude protein content of *L. squarrosulus* was comparable to that reported for *L. tigrinus* by Adejumo and Awosanya (2005). The pilei of both mushroom species contained higher levels of protein than the stipes. The protein contents obtained in the pileus and stipe of *P. atroumbonata* by Alofe (1985) were higher than the values presently observed while the reverse was true for *L. squarrosulus*. In addition, the protein content of both *L. squarrosulus* and *P. atroumbonata* was higher than the values reported for *Auricularia auricula*, *A. polytricha*, *Tremella fuciformis*, *Ganoderma lucidum*, *Calvatia cyathiformis* and *Poria cocos* by Cheung (1997) and Aletor (1995).

The two mushroom species contained more crude fibre in their pilei than in their stipes. However, Alofe (1985) reported the converse for *P. atroumbonata*. Of the two mushroom species, *P. atroumbonata* contained more fibre than *L. squarrosulus*. The values obtained for the two species were slightly higher than the values reported by Aletor (1995), but lower than the values reported by Alofe (1985). In addition, the crude fibre content reported for *L. squarrosulus* by Fasidi and Kadiri (1991) was much higher than the values presently observed. Nonetheless, the crude fibre content presently reported was in agreement with the values reported for *Poria cocos*, *T. fuciformis*, *A. polytricha* and *L. tigrinus* (Cheung, 1997; Adejumo and Awosanya, 2005). Thus, *L. squarrosulus* and *P. atroumbonata* should be regarded as a good source of dietary fibre for humans.

The pileus and stipe of *P. atroumbonata* contained significantly higher levels of ash than the corresponding parts of *L. squarrosulus*. The ash levels obtained for both parts of *P. atroumbonata* were in agreement with the values obtained by Alofe (1985) and Aletor (1995) for the same species. The value obtained for *L. squarrosulus* agreed with that reported by Fasidi and Kadiri (1991), but was lower than the value reported by Alofe (1985). In general, the lower ash content observed for *L. squarrosulus* was in agreement with those reported for *A. auricula*, *A. polytricha* and *P. cocos* (Cheung, 1997). Both *L. squarrosulus* and *P. atroumbonata*, however, had higher ash content than *Lentinus edodes*, *L. shimeiji*, *Pleurotus sajor-caju* and *V. volvcea* as reported by Cheung (1997).

Both the pileus of *L. squarrosulus* and that of *P. atroumbonata* had higher levels of carbohydrate than their corresponding stipes. However, Alofe (1985) reported the stipes as having higher carbohydrate levels than the corresponding pilei. Of the two species *L. squarrosulus* had the higher carbohydrate level and it

compared favourably with the levels reported in *A. bisporus* and *L. tigrinus* by Cheung (1997) and Adejumo and Awosanya (2005), respectively.

## REFERENCES

- Adejumo TO, Awosanya OB (2005). Proximate and mineral composition of four edible mushroom species from South Western Nigeria. *Afr. J. Biotechnol.* 4(10):1084-1088.
- Adenipekun CO, Fasidi IO (2005). Bioremediation of oil polluted soil by *Lentinus subnudus*, a Nigerian white-rot fungus. *Afr. J. Biotechnol.* 4(8): 796-798.
- Aletor VA (1995). Compositional studies on edible tropical species of mushrooms. *Food Chem.* 54: 265-268.
- Alofe FV (1985). The general characteristics and cultivation of some Nigerian mushrooms. Unpublished PhD Thesis, University of Ife, Nigeria.
- Cheung CK (1997). Dietary fibre content and composition of some edible fungi determined by two methods of analysis. *J. Sci. Food Agric.* 73: 255-260.
- Christie P, Easson DC, Picton JR, Love CP (2001). Agronomic value of alkaline-stabilized sewage biosolids for spring barley. *Agronomy J.* 93(1): 144-151.
- Crisan EV, Sands A (1978). Nutritional value. In: *The Biology and Cultivation of Edible Mushrooms*. (Eds Chang ST, Hayes WA). Academic Press, New York. pp. 137-168.
- D'Annibale A, Ricci M, Leonardi V, Quarantino D, Mincione F, Petruccioli M (2005). Degradation of aromatic hydrocarbons by white-rot fungi in a historically contaminated soil. *Biotechnol. Bioeng.* 90(6): 723-731.
- Estévez E, Veiga MC, Kennes C (2005). Biodegradation of toluene by the new fungal isolates *Paecilomyces varioti* and *Exophiala oligosperma*. *J. Ind. Microbiol. Biotechnol.* 32(1): 33-37.
- Fasidi IO, Kadiri M (1991). Changes in nutrient contents of *Termitomyces robustus* (Beeli) Heim and *Lentinus subnudus* Berk during sporophore development *Acta Bot. Hungarica* 36: 167-172.
- Lim BO, Yamada K, Cho BG, Jeon T, Hwang SG, Kang SA, Park DK (2004). Comparative study on the modulation of IgE and cytokine production by *Phellinus linteus* grown on germinated brown rice, *Phellinus linteus* and germinated brown rice in murine splenocytes. *Biosci. Biotechnol. Biochem.* 68(11): 2391-2394.
- Magingo FS, Oriyo NM, Kivaisi AK, Danell E (2004). Cultivation of *Oudemansiella tanzanica* nom. prov. on agricultural solid wastes in Tanzania. *Mycologia* 96(2): 197-204.
- Mata G, Hernandez DM, Andreu LG (2005). Changes in lignocellulolytic enzyme activities in six *Pleurotus* spp. strains cultivated on coffee pulp in confrontation with *Trichoderma* spp. *World J. Microbiol. Biotechnol.* 21(2): 143-150.
- Nwanze PI, Adamu L (2004). Mineral content and amino acid composition of *Lentinus squarrosulus* and *Psathyrella atroumbonata*. *Know. Rev.* 8(2): 33-38.
- Nwanze PI, Khan AU, Ameh JB, Umoh VJ (2004). The effect of grain, media, different oils at different concentrations and type on carpophore production of *P. atroumbonata*. *Roan* 3(1&2): 85-97.
- Nwanze PI, Khan AU, Ameh JB, Umoh VJ (2005a). The effect of media, oil type and rate on the mycelial wet and dry weights of *Lentinus squarrosulus* (Mont.) Singer and *Psathyrella atroumbonata* Pegler in submerged liquid culture. *Afr. J. Biotechnol.* 4(3): 326-331.
- Nwanze PI, Khan AU, Ameh JB, Umoh VJ (2005b). The effect of spawn grains, culture media, oil types and rates on carpophore production of *Lentinus squarrosulus* (Mont.) Singer. *Afr. J. Biotechnol.* 4(6): 472-477.
- Praveena B, Srinivas CVS, Nagaraj G (2000). Quality of some Indian sunflower genotypes and utilization of cakes in snack foods. *HELIA* 23(33): 121-128.
- Snedecor GW, Cochran WG (1987). *Statistical Methods*. Oxford IBH Publishing Co. Ltd., New Delhi. pp. 20-35.
- Sunagawa M, Magae Y (2005). Isolation of genes differentially expressed during fruit body development of *Pleurotus ostreatus* by differential display of RAPD. *FEMS Microbiol. Lett.* 246(2): 279-284.