

Full Length Research Paper

Comparison of multi-locus enzyme and protein gel electrophoresis in the discrimination of five *Fusarium* species isolated from Egyptian cottons

Ibrahim N. Aly¹, Mohmed A. Abdel-Sattar¹,
Kamel A. Abd-Elsalam^{2,3*}, Mohmed S. Khalil² and Joseph A. Verreet³

¹ Suez Canal University, Faculty of Agriculture, Ismailia, Egypt.

² Agricultural Research Center, Plant Pathology Research Institute, Giza, Egypt.

³ Christian Albrechts Universität zu Kiel, Institut für Phytopathologie, Kiel, Germany.

Accepted 21 May 2003

Electrophoretic studies of multilocus-enzymes (MLEE) and whole-cell protein (SDS-PAGE) were carried out in order to evaluate the parity between different methods for the characterization of five *Fusarium* species recovered from cotton-growing areas in Egypt by numerical taxonomy methods. The obtained data revealed that SDS-PAGE and esterase isozymes are more efficient in grouping isolates in their respective species while peroxidase and malate dehydrogenase isozyme has much limited resolution in organizing all isolates in their respective species-specific clusters. A low correlations was detected between geographical origin of isolates and genetic diversity. Results indicate that the estimated inter-specific variation may be more pronounced with protein markers than with isozymes when the two approaches are applied to the same populations. The level of genetic variability detected within and between *Fusarium* spp. accessions with protein and esterase isozymes analysis suggests that it is a reliable, efficient, and effective marker technology for determining genetic relationships in *Fusarium* genus.

Key words: Cotton, *Fusarium*, Isozymes, polyacrylamide gel electrophoresis.

INTRODUCTION

Numerous biochemical and molecular markers are available for the clarification and delineation of fungal taxa. Isozymes studies have proven useful for the identification of fungal cultures at species (Bonde et al., 1991; Petrunak and Christ 1992) or subspecies level (Julian and Lucas 1990; Damaj et al., 1993).

*Corresponding author; phone: (49 431) 880 2993; fax: (49 431) 880 1583; e-mail: kaabdel salam@msn.com

Abbreviations: MLEE, multilocus-enzymes electrophoresis; SDS-PAGE, sodium dodecyl sulphate polyacrylamide gel electrophoresis; UPGMA, unweighted pair group method using arithmetic.

Electrophoretic analysis of whole cell proteins by one-dimensional protein patterns provides a rough measure of the number and physicochemical properties of gene products. One-dimensional polyacrylamide gel electrophoresis of proteins has been used extensively for identification and classification at the strain and species level (Snider, 1973). However, as seen with other methods based on the analysis of gene products, effects such as the influence of the composition of growth media on the banding pattern and superposition of proteins cause problems in the interpretation of data. Highly standardized conditions and computerized comparison of the electropherograms are necessary to obtain reproducibility of results obtained from different laboratories. Isozymes are also useful tools for

differentiating fungal taxa at the specific and intra-specific level. Bosland and Williams (1987) were able to define isozyme electrophoretic types, which could be used to differentiate *Fusarium* species and four *F. oxysporum* formae specialis. The role of *Fusarium* spp. as a pathogen of cotton seedlings, and other crops is well known. Although Johnson and Doyle (1986) reported that *Fusarium* spp. were not important pathogens in cotton seedling disease complex, they are often the most frequently isolated fungi. Isolates of *Fusarium* spp. have previously been shown to be pathogenic on seedling cotton (Colyer, 1988; Aly et al., 1996; Zhang et al., 1996). The objectives of the present study were to differentiate the five closely related *Fusarium* species (*F. oxysporum*, *F. moniliforme*, *F. solani*, *F. avenaceum* and *F. chlamydosporum*) through two main approaches: isozyme analysis and protein gel electrophoresis.

MATERIALS AND METHODS

Fungal cultures

Forty-six isolates of *Fusarium* spp. from diverse geographical origins were used. Mycelium production was carried out by culturing the fungi in 100 ml of potato dextrose broth in 500-ml Erlenmeyer flasks and incubating them at 28°C for 6 days. Mycelial mats were then filtered under vacuum on a Büchner funnel, rinsed twice with distilled water and blotted dry. Mycelium to be used for isozyme and protein analyses was freeze-dried and stored at -20°C.

Preparation of protein extracts

Soluble proteins were extracted by grinding 100 mg freeze-dried mycelium with pestle and mortar in liquid nitrogen and 5 ml buffer solution (0.1M Tris-HCl, pH 6.8). The mixture was centrifuged for 20 min at 17,000 rpm and the supernatant collected. The protein content in supernatant was estimated according to the method of Bradford (1976) with bovine serum albumin as standard protein. Protein content was adjusted to 2 mg/ml per sample.

Standard conditions for SDS polyacrylamide gel electrophoresis (protein fingerprinting)

SDS-PAGE was performed by the method described by Laemmli (1970), and modified according to Hames (1995). Proteins were analyzed on 1.5-mm thick and 15-cm long gels run in a dual vertical slab unit (Hoefer Scientific Instruments, San Francisco, CA, USA, Model SE 600 Series Hoefer® Pharmacia Biotech). From each sample, 25 µl of extract was loaded on a polyacrylamide gel. The separation gel (10%) and stacking gel (3.5%) were prepared from an acrylamide monomer solution (Roth, Karlsruhe, Germany). Electrophoresis was carried out at a constant current of 35 mA through the stacking gel, and at 90 mA through the separation gel at room temperature. After electrophoresis, the gels were stained by silver nitrate (Rabilloud et al., 1988).

Sample preparation for isozymes electrophoresis

One-hundred milligrams of freeze-dried mycelium were homogenized by grinding in liquid nitrogen with a mortar and pestle, and suspended in 1.5 ml of extraction buffer (10 mM Tris-HCl, 1

mM EDTA, 0.1 mM ascorbic acid, pH 7). Extracts were chilled in an ice bath for 1-3 h and centrifuged at 18,000 rpm for 30 min at 4 °C. The supernatant was collected, divided into 100-µl aliquots, stored at -80 °C, and used within 6 months.

Multilocus-enzymes electrophoresis

Isozymes electrophoresis was performed in vertical polyacrylamide gels with a discontinuous buffer system as described by Iglesias et al. (1974). The following enzyme systems were screened: esterase (EST, 3.1.1.1), malate dehydrogenase (MDH, 1.1.1.37), and peroxidase (POX, EC 1.11.1.7). The staining protocols used are described by Gall et al. (1995).

Computing numerical data

Dendrograms for the different MLEE systems and SDS-PAGE were generated by using the one-dimensional software analysis (Advanced American Biotechnology and Imaging, Fullerton CA 92831, USA). For cluster analysis of protein and isozymes banding patterns, the unweighted pair group method using average linkages (UPGMA) was used (Vauterin and Vauterin, 1992).

RESULTS

SDS-PAGE analysis

Cluster analysis of the protein markers data placed the *Fusarium* species into main groups (Figure 1), according to their previous species assignment. The genetic similarity between *Fusarium* spp. isolates ranged from 21 to 54% for inter-specific and 62 to 97% for intra-specific comparisons. The application of UPGMA clustering produced two large clusters within the population with a branched-off at genetic similarity of 21.46%, each consisting of several subclusters (phenons).

The first main cluster included all *F. oxysporum*, all *F. moniliforme* and *F. chlamydosporum* isolates. Within this cluster, Fa1 and Fa2 have a genetic similarity of GS=95.98%. Isolates Fo24 and Fo25 also showed very high genetic similarity of 99.22%. The second cluster consists of *F. solani* isolates at the genetic similarity of 31.37%. Here, isolates Fs3 and Fs4 showed very high genetic similarity of 97.26%. Clustering analysis indicates that the 46 *Fusarium* spp. isolates can be assigned to six protein types. There was low correlation between clustering in the protein dendrogram and geographic origin. These results suggested that protein profiles data clearly separated *Fusarium* spp. isolates.

Isozymes analysis

The activity and resolution of three enzyme-staining procedures for *Fusarium* spp. were tested. Poor resolution was observed for the enzymes peroxidase and malate dehydrogenase (data not shown), while esterase




Figure 2. Dendrogram derived from the UPGMA linkage of correlation coefficients between esterase isozyme profiles from five *Fusarium* spp. isolates. The levels of linkage representing the Pearson product-moment correlation coefficient (r) are expressed as percentages for convenience. The banding profiles against each branch represent normalized and background-subtracted digitized gel strips processed in ABB program.

isozymes to differentiate *Fusarium* species and four *F. oxysporum* formae specialis, as well as isolates obtained from different crucifers species. On the contrary, no clear-cut results were found for isozymes study of *F. oxysporum* f. sp. *lycopersici* recovered from different sites and different pathogenic races or vegetative compatibility groups (Elias and Schneider, 1992). SDS-PAGE is used because the method alleviates the need for culturing, and samples are analyzed in a more direct manner. This method is relatively easy and many samples can be analyzed at the same time. It is also cheaper than AFLP fingerprinting. Moreover, the results obtained by SDS-PAGE of whole-cell proteins can discriminate at much the same level as DNA fingerprinting (Priest and Austin, 1993) in some cases.

Cluster analysis of the protein markers data placed the *Fusarium* species into main groups according to their previous species assignment. The genetic similarity between *Fusarium* spp. isolates, similarities ranged from 21 to 54% for inter-specific and 62 to 97% for intra-specific comparisons. The application of UPGMA clustering produced two large clusters within the population. These results suggested that protein profiles

data can clearly separate *Fusarium* spp. isolates with a few exceptions. A low correlation between protein dendrogram and geographic origin of tested isolates was found.

These results agree with those obtained by Mandeel et al. (1994) who compared SDS-PAGE patterns from eight isolates belonging to three *Fusarium* species. Protein profiles were distinct, and each isolate showed a uniquely characteristic profile. The data obtained from protein profiles support the potential use of this experimental approach to help distinguish between different *Fusarium* isolates. On the contrary, Belisario et al. (1998) found no differences when comparing total mycelium protein profiles (SDS-PAGE) of different species and formae species of *F. oxysporum*, *F. solani* and *F. culmorum*.

Isozyme analysis is frequently used for taxonomic purposes, especially when a taxon is morphologically diverse. In most cases, fungal species are easily differentiated by electrophoresis. The technique is commonly used to make recommendations on the separation or combination of species (Otrosina et al., 1992). Zambino and Harrington (1989) have also separated subspecies, varieties, and intersterility groups.

Simple band-counting procedures can be used to distinguish taxa, although cladistic and phylogenetic information can be derived from the allelic frequencies and ratios derived from a genetic interpretation of the data. Isozyme analysis is most successful in distinguishing species and subspecies when the amount of intra-specific genetic variation is limited within a population. Esterase isozyme fingerprinting have been frequently used because of the relatively large differentiation between *Fusarium* spp. in contrast to other enzyme systems (Baayen, 1997).

In the present study, UPGMA cluster analysis of esterase isozyme produced two large clusters within the population, each consisting of several subclusters. Low correlation was observed between clustering in the esterase dendrogram and sampling date of tested isolates. These results suggested that esterase patterns data clearly separated *Fusarium* spp. isolates with a few exceptions. These results, similar to those obtained by Ye and Wu (1985) who showed that the esterase patterns are distinct for 24 *F. graminearum* isolates. The patterns of 7 isolates of *Gibberella zeae* collected from different geographic localities appeared little different from one another and there were some identical bands of esterase isozymes in their patterns. However, no relationship was observed between isozyme patterns and geographic origin, phenotypic distance or virulence of isolates (Etebarian et al., 1996; Yli-Mattila and Hyvonen, 1996; Bosland and Williams, 1987). Also no clear-cut result was detected by malate dehydrogenase and peroxidase isozymes in discriminating between *Fusarium* spp. Isolates (Ye and Wu, 1985). In summary, the genomic diversity within the genus *Fusarium* was determined by using esterase isozyme and SDS-PAGE and each technique could prove useful for the rapid classification of *Fusarium* spp. isolates.

REFERENCES

- Aly AA, Hussein EM, Mostafa MA, Ismail AI (1996). Distribution, identification, and pathogenicity of *Fusarium* spp. isolated from some Egyptian cottons. *Menofiya J. Agric. Res.* 4: 819-836.
- Baayen RP, van-Dreven F, Krijger MC, Waalwijk A (1997). Genetic diversity in *Fusarium oxysporum* f. sp. *dianthi* and *Fusarium redolens* f. sp. *dianthi*. *Eur. J. Plant Pathol.* 103:395-408.
- Belisario A, Luongo L, Balmas V, Pezza L, Corazza L (1998). *Fusarium* wilts of winter melon. Sixth SIPaV Annual Meeting "Plant Pathology and Sustainable Agriculture" Campobasso, 17-18 September 1998 (Abst.)
- Bonde MR, Peterson GL, Emmett RW, Menge JA (1991) Isozyme comparison of *Septoria* isolates associated with citrus in Australia and the United States. *Phytopathology* 81: 517-521.
- Bosland PW, Williams PH (1987). An evolution of *Fusarium oxysporum* from crucifers based on pathogenicity, isozymes polymorphisms, vegetative compatibility, and geographic origin. *Can. J. Bot.* 65: 2067-2073.
- Bradford MM (1976). A rapid and sensitive for the quantification of microgram quantities of protein utilizing the principle of protein dye-binding. *Anal. Biochem.* 72: 248-254.
- Brown JKM (1996). The choice of molecular marker methods for population genetic studies of plant pathogens. *New Phytologist* 133: 183-195.
- Colyer PD (1988). Frequency and pathogenicity of *Fusarium* spp. associated with seeding diseases of cotton in Louisiana. *Plant Dis.* 72: 400-402.
- Damaj M, Jabaji-Hare SH, Charest PM (1993) Isozyme variation and genetic relatedness in binucleate *Rhizoctonia* species. *Phytopathology* 83: 864-871.
- Elias KS, Schneider RW (1992). Genetic diversity within and among races and vegetative compatibility groups of *Fusarium oxysporum* f. sp. *lycopersici* as determined by isozymes analysis. *Phytopathology* 82: 1421-1427.
- Etebarian HR, Wilcoxson RD, Stewart EL, Woodward RP (1996). Some isozyme patterns of *Fusarium culmorum*. *Iran. J. of Plant Pathol.* 32: 9-21.
- Gall C, Balesdent MH, Desthieux I, Robin P, Rouxel T (1995) Polymorphism of Tox, *Leptosphaeria maculans* isolates as revealed by soluble protein and isozyme electrophoresis. *Mycol. Res.* 99: 221-229.
- Hames BD (1995). One-dimensional polyacrylamide gel electrophoresis. In: *Gel electrophoresis of proteins. A practical approach.* Hames, B.D. and Rickwood, D. (eds.) Oxford University Press, New York 1-147 pp.
- Iglesias LH Lima Simon LP (1974). Isozyme identification of zygotic and nucellar seedlings in citrus. *J. Hered.* 65: 81-84.
- Johnson LF, JH Doyle (1986). Relationships of seeding disease of cotton to characteristics of loessial soil in Tennessee. *Phytopathology* 76:86-290.
- Julian AM, Lucas JA (1990). Isozyme polymorphism in pathotypes of *Pseudocercospora herpotrichoides* and related species from cereals. *Plant Pathol.* 39: 178-190.
- Laemmli UK (1970) Cleaving of the structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- Majer D, Mithen R, Lewis BG, Vos P, Oliver RP (1996). The use of AFLP fingerprinting for the detection of genetic variation in fungi. *Mycol. Res.* 100: 1107-1111.
- Mandeeel QA, El-Din AYG, Mohammed SA (1994). Analysis of SDS-dissociated proteins of pathogenic and nonpathogenic *Fusarium* species. *Mycopathologia* 127: 159-166.
- McDonald BA, McDermott JM (1993). Population genetics of plant pathogen fungi. *Bioscience* 43: 311-319.
- McDonald BA, Zhan J, Jarden O, Hogan K, Garton J, Pettway RE (1999). The population genetics of *Mycosphaerella graminicola* and *Phaeosphaeria nodorum*. In: Lucas J. A; P. Bowyer and H. M. Anderson (eds) *Septoria on cereals* (pp 44-69) CAB international, Wallingford, UK.
- Otrosina WJ, Chase TE, Cobb FWJ (1992). Allozyme differentiation of intersterility groups of *Heterobasidion annosum* isolated from conifers in western United States. *Phytopathology* 82: 540-545.
- Petrunk DM, Christ BJ (1992) Isozyme variability in *Alternaria solani* and *A. alternata*. *Phytopathology* 82: 1343-1347.
- Priest FG, Austin B (1993). *Modern Bacterial Taxonomy* (2nd ed.) Chapman & Hall, London.
- Rabilloud TG, Carpentier A, Tarroux P (1988). Improvement and simplification of low-background silver staining of proteins by using sodium dithionite. *Electrophoresis* 6: 288-291.
- Snider RD (1973). Electrophoresis and taxonomy of phytopathogenic fungi. *Bull. Torrey Bot. Club* 100: 272-276.
- Vauterin LA, Vauterin P (1992). Computer-aided objective comparison of electrophoresis patterns for grouping and identification of microorganisms. *Eur. Microbiol.* 1: 37-41.
- Ye HZ, Wu GQ (1985). Polyacrylamide gel electrophoresis studies of the esterase patterns of *Fusarium* sp. causing head blight of wheat and barley. *Acta Phytopathol. Sinica* 15: 87-94.
- Yli-Mattila T, Hyvonen J (1996). RAPD-PCR analysis of *Fusarium* strains - cladistic evaluation of the results. *Sydowia* 48: 184-195.
- Zambino FJ, Harrington TC (1989). Isozyme variation within and among host-specialized varieties of *Leptographium vageneri*. *Mycologia* 81: 122-128.
- Zhang J, Howell CR, Starr JL. (1996). Suppression of *Fusarium* colonization of cotton roots and *Fusarium* wilt by seed treatments with *Gliocladium virens* and *Bacillus subtilis*. *Biocontrol Sci. Technol.* 6:175-187.