

Review

Biological, ecological and agronomic significance of plant phenolic compounds in rhizosphere of the symbiotic legumes

Joachim H.J.R. Makoi¹ and Patrick A. Ndakidemi^{2*}

¹Faculty of Applied Science Cape Peninsula University of Technology, Cape Town Campus, Keizergracht, P.O. Box 652, Cape Town 8000, South Africa

²Research and Technology Promotion, Cape Peninsula University of Technology, Cape Town Campus, Keizergracht, P.O. Box 652, Cape Town 8000, South Africa.

Accepted 7 May, 2007

Phenolics are low molecular compounds ubiquitous in all tissues of higher plants with great significance in plant development. Our understanding of some phenolic compounds in the last few decades has greatly improved. However, their biological, ecological and agronomical significance in the rhizosphere of most symbiotic legumes is much less clear. Further understanding of these biomolecules will increase our knowledge of their contribution in soil and water conservation, weed management, mineral element nutrition, their impact as signal molecule in certain symbiotic relationships, and their role as defence molecules against soil pests and pathogens. This article reviews the bioecological and agronomical significance of phenolic compounds in the rhizosphere of a few studied symbiotic legumes and other plants.

Key words: Allelopathy, defence, flavonoids, growth inhibitors, soil structure, nutrients availability, pesticide, signal molecules, soil borne diseases, soil insects.

INTRODUCTION

Phenolic compounds are some of the most widespread molecules among plant secondary metabolites, and are of great significance in plant development. They are involved in diverse processes including rhizogenesis (Curir et al., 1990), vitrification (Kevers et al., 1984), resistance to biotic and abiotic stress (Delalonde et al., 1996), and redox reactions in soils (Takalama and Oniki, 1992). Additionally, they serve as flower pigments, act as constitutive protection agents against invading organisms, function as signal molecules, act as allelopathic compounds, and affect cell and plant growth (Dakora, 1995; Dakora and Phillips, 1996; Ndakidemi and Dakora, 2003), are important natural animal toxicants (Adams, 1989) and some may function as pesticides (Vidhyasekaran, 1988; Waterman and Mole, 1989; Beier, 1990). They are also functional components of the rhizosphere

and its soil organic matter (Haider et al., 1975; Martin and Focht, 1977). They have long been recognised as allelochemicals for weed control (Rice, 1984; Putnam and Tang, 1986) phytoestrogens in animals (Adams, 1989) and plant defence molecules (Vidhyasekaran, 1988). In the rhizosphere, they act as important precursors for the synthesis of soil humic substances (Haider et al., 1975).

The synthesis and release of phenolics are induced by various biotic and abiotic factors. Boron deficiency can, for example, induce the accumulation of phenolics in plants. Tissue injury, pathogen attack, herbivory, and infection by microsymbionts such as rhizobium can also cause synthesis and release of phenolics (Recourt et al., 1991; Dakora et al., 1993a, b; Lawson et al., 1996). Inside and outside plant tissues, these metabolites are known to function as phytoalexins, phytoanticipins and node gene inducers (Dakora and Phillips, 1996, 2002). Although several authoritative reviews have examined phenolic molecules in plant development (Long, 1989; Bekkara et al., 1998; Dakora and Phillips, 1996), few have assessed

*Corresponding author. E-mail: ndakidemi56@yahoo.co.uk. Tel: +27214603128. Fax: +27214603887.

their role in the quality and function of rhizosphere soil. This paper discusses phenolics as major players in soil function and plant growth.

PHENOLICS AS MODIFIERS OF RHIZOSPHERE SOIL STRUCTURE

In the rhizosphere, soil particles are bound together by phenolics and other exudate compounds from roots, microbes and decomposing organic matter to form aggregates, thus providing support for plant roots. As a living biological system, the soil consists of plant roots, micro flora and micro fauna whose chemical products bind with soil particles to enhance aggregation.

Phenolics and other organic molecules from root and seed exudates, leaf leachates, and decaying plant residues, play a major role in soil formation and pedogenesis. Because some phenolic metabolites are phytotoxic, their presence in higher concentrations can affect soil function (Patrick, 1971; Lodhi, 1975; Chou and Lin, 1976; Rice, 1984; Putnam and Tang, 1986; Elliott and Cheng, 1987; Waller, 1987). The quantity and quality of phenolics released by plants can differ from species to species. For example, ryegrass roots have a much higher tissue concentration of phenolics than red clover (Whitehead et al., 1979). The concentration of phenolic acids in soil can range from 2.1% to 4.4% for roots of monocots and 0.1% to 0.6% for dicots (Hartley and Harris, 1981; Smith and Hartley, 1983; Hartley and Whitehead, 1985).

The release of phenolic compounds by plants can vary with time, space, and location. In *Cistus albidus* green leaves, phenolic concentration ranged from 66.5 to 95.9 mg gallic acid g⁻¹ DM depending on the time of the year and the soil type, with maximum phenolic accumulation in winter and when growing on granitic and schistic soils (Eva and Josep, 2003). In valley bottom ecosystems, the concentration of individual phenolic acids (*p*-coumaric acid) was as high as 6.7% in January, but dropped to 2.0% in September (Lodhi, 1975), a decrease attributed to leaching, plant uptake, and microbial transformation (Martin et al., 1972; Turner and Rice, 1975; Martin and Haider, 1976). These situations suggest that phenolics in soil-plant continuum undergo a continuous cycle of deposition, decomposition, plant uptake, leaching and chemical immobilisation. In short, as phenolics moves through the rhizosphere, they may be transformed, metabolised by microbes, or bound by soil organic matter. In doing so, the soil structure is shaped, thus improving both soil porosity and plant growth.

Soil structure, soil macro and micro porosity in the rhizosphere may be improved by phenolic compounds in favour of air and water movement for both abiotic and biotic life in the soil ecosystem. Therefore, understanding their functional groups will be of vital importance for the overall improvement of agricultural production systems. Additionally, understanding which plant releases what type and amount of phenolic compound(s) will be of

utmost importance for better soil and water conservation.

ROLE OF PHENOLIC COMPOUNDS ON NUTRIENT AVAILABILITY IN THE RHIZOSPHERE

Phenolics are known to play a dominant role in rhizosphere mineral elements and organic matter dynamics by affecting organic matter degradation, humus formation, alteration of microbial activities, mineralization of N and its availability (Schimel et al., 1996; Northup et al., 1998; Hattenschwiler and Vitousek, 2000; Kraus et al., 2003). Although the uptake, metabolism and toxic effects of most phenolics in the rhizosphere of many plants are well documented (Curir et al., 1990; Klein and Blum, 1990), their biochemical mechanisms, biosynthesis, release and functioning as affected by the host plants such as legumes are to a large extent unknown.

Phenolics can have both negative and positive influence on plant mineral elements availability in the rhizosphere. There is evidence that green leaves and decomposing litter from plants can influence rhizosphere N dynamics through rainfall leachate of phenolics as chlorogenic acid (Eva and Josep, 2003), or through different mechanisms which alter phenolic synthesis. These mechanisms include formation of protein complexes, which in turn delays organic matter decomposition and mineralization (Horner et al., 1988; Nicolai, 1988; Oades, 1988; Palm and Sanchez, 1990; Bending and Read, 1995; Northup et al., 1995, 1998; Hoenschwiler and Vitousek, 2000); increase of microbial activity when used as carbon source (Sparling et al., 1981; Schimel et al., 1996; Schmidt et al., 1997; Blum, 1998; Fierer et al., 2001); N immobilization (Shafer and Blum, 1991; Sugai and Schimel, 1993; Schimel et al., 1996; Blum, 1998) and direct inhibition of nitrification (Rice, 1984; Boufalis and Pellissier, 1994). From these mechanisms, plants growing on infertile soils could benefit from the sequestered N into a large unavailable pool of recalcitrant organic matter.

Phosphate supply is one of the major constraints to plant growth due to its insolubility and high sorption capacity in soil (Marschner, 1995). However, there is now overwhelming evidence suggesting that some plants may directly modify their rhizosphere in order to gain access to unavailable soil P reserves (Morel and Hinsinger, 1999). These rhizosphere modifications include the release of phenolics, such as syringic, caffeic, and protocatechuic acids which then can solubilize and accumulate inorganic P. It has now been demonstrated that some plants such as *Lupinus albus* and *Brassica napus* are capable of releasing large amounts of phenolics into the rhizosphere in response to P deficiency (Laheurte and Berthelin, 1988; Hoffland et al., 1992; Gerke, 1994; Johnson et al., 1996; Imas et al., 1997). These phenolics may, in turn, improve plant P availability by interacting with metal oxides which could otherwise fix P, or desorb phosphate from sesquioxide surfaces by anion (ligand) exchange (Parfitt, 1979; Gerke, 1992). Detailed physiolo-

logical and biochemical studies to identify other specific role(s) phenolics can play in supplying P to plants are recommended.

Phenolics may also increase the availability of macro and micro elements (e.g. K, Ca, Mg, Cu, Zn, Mn, Mo, B) through formation of organic metal complexes and by providing sorption sites (Marschner, 1988; Hoffland, 1992; Slabbert, 1992; Marschner, 1995; Micales, 1997).

Phenolic acids are strong metal complexing agents representing a major group of organic acids that can affect solution concentration of cations, mobility, bioavailability and facilitation of nutrient uptake especially Fe (Tiarks et al., 1989; Slabbert, 1992; Scalbert et al., 1999; Seneviratne and Jayasinghearachchi, 2003). Under acid condition some plants such as legumes have been reported to release reductant compounds that enhance uptake of Fe (Osen et al., 1979). For example, the growth of excised pea roots (DeKock and Vaughan, 1975) and Fe-deficient tomato plants were promoted by enhancing Fe uptake when phenolics were added in the medium (Osen et al., 1981). The mechanism involved is that *p*-coumaric acid released by roots or decaying residue is hydroxylated into caffeic acid that can reduce Fe³⁺ and increase its mobilisation as shown in the following equation:



Copper (Cu) plays an important role in plant growth including nodulation and N₂-fixation in legumes (Cartwright and Hallsworth, 1970). Marked differences have been observed in Cu²⁺ nutrition between plants growing in Cu²⁺ enriched and non-contaminated soils (Ernst et al., 1992). The mechanism enabling plants to tolerate high concentrations of Cu²⁺ in their rhizosphere are still poorly understood. However, *in vitro* adsorption studies by Jung et al. (2003) showed that phenolics such as flavonoids and polyphenols mediated Cu²⁺ toxicity in the rhizosphere and roots apoplasm through complexations of Cu²⁺ ions, thus alleviating the Cu²⁺ toxicity. Further studies on the mechanisms involving Cu²⁺ nutrition and phenolic compounds should be undertaken in the rhizosphere of symbiotic legumes.

Aluminium (Al) rhizotoxicity is one of the biggest limitations to crop production in the world and is characterized by the inhibition of root cell elongation and to a lesser extent cell division (Kochian, 1995; Kochian and Jones, 1997). However, through phenolic compounds, some plants appear to be able to resist toxic concentrations of Al (>5 μM) in plants (Powel and Rate, 1987; Kochian, 1995; Ma et al., 1997). For instance, Kidd et al. (2001) showed that exposing maize roots to higher Al concentration resulted in higher exudation of penta-hydroxyflavone and morin phenolic compounds.

These molecules are known to have metal binding activity in plants (Kochian, 1995; Ma et al., 1997). More research efforts should explore other phenolic compou-

nds and the mechanism(s) involved in defending plants against this toxic element.

On the contrary, phenolic toxicity has also been reported to cause severe mineral elements deficiency in rhizosphere. For instance, maize plant growing in agricultural soil infested with *Agropyron repens*, a strong allelopathic weed rich in phenolics, suffered from severe deficiency of N, P and K (Buchholtz, 1971; Rice, 1984). The inhibition of these phenolics to K and P uptake was correlated with octanol-water partition coefficients and lipid solubility of the phenolics respectively. Thus, further studies on inhibition of other mineral elements by phenolic compounds are a subject that warrants further investigation. Understanding of these compounds and application may lead to greater revolution in soil fertility.

PHENOLICS AS PLANT GROWTH INHIBITORS IN THE RHIZOSPHERE

Growth-inhibiting compounds of agricultural importance have recently received considerable research attention (Siqueira et al., 1991; Inderjit, 1996). These compounds include phenolic compounds with allelopathic characteristics (Schenk et al., 1999; Callaway and Aschhoug, 2000). Allelopathy is an adverse influence of one plant or micro-organism on another (Rice, 1984) or production of active chemicals by living or decaying plant tissues which interfere with the growth of neighbouring plants (Muller, 1969; Rice, 1984; Putnam and Tang, 1986; Waller, 1987; Weidenhamer et al., 1989; Schenk et al., 1999; Callaway and Aschhoug, 2000). Studies have shown that these chemicals include simple phenols, phenolic acids derived from benzoic and cinnamic acids, coumarins, flavonoids, isoflavonoids, tannins and a variety of phenolic conjugates (Figures 1, 2 and 3). Their presence and accumulation in the soil may reach the threshold concentrations for inhibition of pre-emergence seed germination or post-germination, growth and other plant functions (Whitehead, 1964; Baskin et al., 1967; Einhellig, 1986; Leather and Einhellig, 1986, Fisher, 1987; Patterson, 1987). Different phenolics concentrations have been reported to inhibit seed germination and seedling growth in legumes (Beier et al., 1983; Zaynoun et al., 1984). For instance, 50% inhibition of germination and seedling growth required concentration of phenolic acids ranging from 150 – 1000 μg mL⁻¹ for rye grass and Lucerne (Guenzi and McCalla, 1966).

Other phenolics such as myrcetin, quercetin, and myrcitrin have been identified as seed coat rhizobium toxins (Fottrell et al., 1964) and prodelphindin, quercetin, rutin, catechol, *p*-coumaric, gentistic, salicylic and vanillic acids as rhizobia cell, bacteria and fungi inhibitors (Table 1; Debman and Smith, 1976; Skipp and Bailey, 1977; Kandasamy and Prasad, 1979; Pankhurst and Biggs, 1980; Adesanya et al., 1986; Thompson, 1988). It is, therefore, important to identify more phenolic compounds inhibiting plant growth and micro-organisms in the rhizo-




Figure 1. Chemical structures of selected phenolic compounds from legumes.

sphere, and have a better understanding of their roles and mechanisms involved in such interactions for better management of ecosystems.

Despite the fact that we have a considerable understanding of phenolics as allelochemicals and their allelopathic effects, there are still some noxious weeds such as *Striga* which retard plant growth and cause severe yield losses in cereals and legumes worldwide. Understanding of different phenolic compounds from various plant species including legumes and their allelopathic effects, and how best they can be used for weed management such as *Striga* and other parasitic weeds is vital.

PHENOLICS AS SIGNALS TO SYMBIOTIC RHIZOBIA AND VESICULAR-ARBUSCULAR MYCORRHIZAL

The exudation of low molecular weight compounds by seeds in the soil or in plant root is known to activate the expression of *nod* genes (Innes et al., 1985; Mulligan and Long, 1985; Rossen et al., 1985; Firmin et al., 1986; Peters et al., 1986; Redmond et al., 1986; Horvath et al., 1987; Spaink et al., 1987; Dakora et al., 1993; Leon-Barrios et al., 1993; Dakora, 1994; Phillips et al., 1994). These organic molecules include, amongst others, the phenolic compounds (Table 2). Many varieties of phenolic identified as *nod* gene inducers have been well described in Dakora et al. (1993); Leon-Barrios et al. (1993); Dakora, (1994); Phillips et al. (1994); Ndakidemi and

Dakora (2003). For instance, broad bean (*Vicia faba*) releases flavones and flavonols (Bekkara et al., 1998), soybean releases isoflavonoids (D'Arcy-Lameta and Jay, 1987) and peanut releases vanillin (Zawoznik et al., 2000). Such phenolics can accumulate in legume rhizosphere and facilitate *nod* gene induction and control the nodule formation (Dakora et al., 1993; Leon-Barrios et al., 1993; Dakora, 1994; Phillips et al., 1994; Dakora and Phillips, 1996).

The contribution of vesicular-arbuscular mycorrhizal to increased P acquisition and improvement of water relations of crop plants in our agricultural systems has been extensively reviewed (Smith and Read, 1997) and the role of phenolics in mycorrhizal establishment has also been reported (Anderson, 1988; Paula and Siqueira, 1990). For example, phenolic compound isoflavonoid from roots of soybean promoted the colonization of vesicular-arbuscular mycorrhizal fungi in soybean (Morandi and Bailey, 1984; Morandi, 1989; Wyss et al., 1989). Seed exudates from white clover identified as biochanin A, chrysin and formononetin also promoted the colonisation of AM fungi (Siqueira et al., 1991). Summing up these findings, the colonization of vesicular-arbuscular mycorrhizal at low soil P availability has been related to the accumulation of specific phenolic compounds capable of stimulating hyphal growth, appressorial formation and penetration of vesicular-arbuscular mycorrhizal fungi. Further understanding of the relationship between phenolic accumulation and vesicular-arbuscular mycorrhizal




Figure 2. Chemical structures of selected biologically active flavonoid-derived compounds: A) Isovitexin B) Maackiain C) Genistein D) Coumestrol E) Procyanidin F) Kaempferol G) Luteolin H) Naringenin.

formation is therefore, of utmost importance. This is because manipulation of phenolic compounds may lead to higher vesicular-arbuscular mycorrhizal infections, leading to greater P acquisition for plant nutrition, improved water relations and consequently resulting in better plant growth.

Although several studies have shown that a wide range of phenolic compounds such as flavonoids, alkaloids, isoflavonoids, flavonols, luteolin, quercetin, trigonelline, stachydrine and kaempferol are signal molecules in the rhizosphere (Peters et al., 1986; Hartwig et al., 1990; Hungria et al., 1991; Dakora, 1994; Phillips et al., 1995), it will not be surprising to uncover a large number of other

unknown phenolic compounds being involved in *nod* gene induction processes and mycorrhizal infections. Based on these facts, more research and understanding of the production and release of signal molecules from different legume species is needed.

PHENOLICS IN THE RHIZOSPHERE AS DEFENCE MOLECULES AGAINST SOIL-BORNE PATHOGENS

The consequences imposed by soilborne pathogens in our agricultural systems may favour different defence strategies by plants in their rhizosphere. There is a trem-




Figure 3. Chemical structures of selected phenolic compounds against plant pathogens: A) Glyceolin B) Kievitone C) Resverastrol D) Pisatin.

endous pressure on farmers to use safe control methods for soilborne diseases which do not pollute or degrade the environment (Duncan, 1991; Akhtar, 1995). Fortunately, phenolic compounds can offer an alternative to the chemical control of pathogens on agricultural crops (Tables 1 and 3). Studies have shown that plant defence against pathogens, nematodes, phytophagous insects is based on the synthesis, release and accumulation of various phenolic compounds (Rich et al., 1977; Russel et al., 1978; Dakora, 1995; Dakora and Phillips, 1996). For example, simple and complex phenolic compounds such as cajanin, medicarpin, glyceolin, rotenone, coumestrol, phaseolin, phaseolinin are known to accumulate in tissue

of tropical legumes and act as phytoalexins, phytoanticipins and nematicides against soilborne pathogens and phytophagous insects (Dakora and Phillips, 1996; Ndakidemi and Dakora, 2003; Tables 1 and 3). Alfalfa roots release isoflavonoid 2-(3'5'-dihydroxyphenyl)-5, 6-dihydroxybenzofuran which serves as phytoalexin against root pathogens such as *Fusarium oxysporum* f. sp. *Phaseoli* (Massaoka et al., 1993). Generally, when released into the soil from seeds, roots or residue decomposition, these molecules can act against soilborne pathogens and root-feeding insects (Ndakidemi and Dakora, 2003). Such natural defence provided by these biomolecules from symbiotic legumes in the rhizosphere deserves more scientific attention because of its ecological potential as a sustainable means of reducing soil borne infections in the ecosystems. To date, few studies have validated the role of phenolics against the soil borne pathogens. Therefore, establishing the biological significance of these molecules from unexplored legumes and other plant species could help in the integrated soilborne pathogen control programmes.

PHENOLICS AS PESTICIDAL MOLECULES AGAINST SOIL INSECTS

It is widely documented that plants produce a wide variety of secondary metabolites as defensive weapons against insect pests during attack (Dakora, 1995; Dakora and Phillips, 1996; Ndakidemi and Dakora 2003; Table 3). These include alkaloids, terpenoids, phenolic compounds, and many others (Ndakidemi and Dakora 2003). During the past few decades researchers have identified plant extracts highly effective against a good number of plant pests (Dakora, 1995; Dakora and Phillips, 1996; Ndakidemi and Dakora 2003). The most active documented compounds have been isolated from the Neem tree (Egunjobi and Afolami, 1976; Badra and Eligindi, 1979; Thakur et al., 1981, Muller and Gooch, 1982; Akhtar and Alam, 1993; Akhtar and Mahmood, 1996; Akhtar, 1998; Akhtar and Malik, 2000). Phenolic compounds extracted from different parts of Neem tree include: limo-noids, phenols, tannins, nimbin, salanin, thionemone, azadirachtin and various flavonoids. These compounds have been found to possess nematicidal properties and are also reported to be highly effective against nematodes and insects (Badra and Eligindi, 1979; Thakur et al., 1981, Akhtar and Malik, 2000; Table 3). Scientists working with symbiotic plants should find out if similar or other active compounds exist in their tissues.

Studies with a few symbiotic plants have also shown that the presence of the isoflavonoid vesitol in roots of legume *Lotus pedunculatus* made it resistant to the larvae of black beetle (*Heteronychus aratoo*) attack (Russel et al., 1978). Plants were susceptible to the same pest in a similar genus of different species (*Lotus corniculatus*) that was lacking the compound vesitol in their roots (Russel et al., 1978).

Table 1. Phenolics compound inhibitors of micro-organisms.

Micro-organism	Phenolic compound	Reference
Fungi	Catechol, sinarpic, phaseolin, phaseolidin, phaseolin isoflavan, kievitone, medicarpin, <i>p</i> -coumaric, ferulic, gentisic, benzoic, caffeic, <i>p</i> -coumaric, salicylic acid, benzoquinones, engenol, vanillin, gallic acid, pisatin, genistein, maackianin daidzein, isoprunitin, phaseoluteone, isoferrinin, isomecycarpin, glyceollin, glyceollin 1 coumestran, biochanin A, echinacin	Lingappa and Lockwood, 1962; Christie, 1965; Li et al., 1969; Choudhury et al., 1974; Keen, 1975; VanEtten, 1976; Debman and Smith, 1976; Skipp and Bailey, 1977; Wymann and VanEtten, 1978; Alfenas et al., 1982; Kramer et al., 1984; Stossel, 1985; Adesanya et al., 1986; Kasenberg and Traquar, 1988; Achenbach et al., 1988; Singh et al., 1988; Ravin et al., 1989; Weidenborner et al., 1989 & 1990.
Bacteria	Phaseolin, phaseolin isoflavan, kientone, quercetin-glucoside, myracetin quercetin, tanins, syringic, <i>p</i> -hydrobenzoic, ferulic, caffeic, chlorogenic, pisatin,	

Table 2. Phenolic compounds as *nod* gene inducers found in germinating seeds and root exudates of selected legume plants.

Legume	Phenolic compound	Functional role	Reference
Alfalfa (<i>Medicago sativa</i> L.)	3',4', 5, 7 tetrahydroxyflavone; 3 methoxyluteolin; luteolin; luteolin-7- <i>O</i> -glucoside; 5-methoxyluteolin; 5-methoxycrisoeriol; 3- <i>O</i> -quercetin galactoside; trigonelline; stachydrine 7,4'-dihydroxyflavone, 7-hydroxyflavone, 4,4'-dihydroxy-2-methoxychalcone, chrysoeriol, 2'-methoxychalcone	<i>nod</i> gene inducers	Peters et al., 1986; Kapulnik et al., 1987; Maxwell et al., 1989; Hartwig et al., 1990; Phillips et al., 1995;
	quercetin, 4,7'-dihydroxyflavonone,	AM hyphal growth inducer	Tsai and Phillips, 1991
Cowpea (<i>Vigna unguiculata</i> L.)	Daidzein, genistein, coumestrol	<i>nod</i> gene inducers	Dakora et al., 2000
Bean (<i>Phaseolus vulgaris</i> L.)	Genistein, genistein-3- <i>O</i> -glucoside, pentunidin, malvin, myricetin, quercetin, daidzein, coumestrol, kaempferol, naringenin, eriodictyol	<i>nod</i> gene inducers	Hungria et al., 1991, Dakora et al., 1993
Clover (<i>T.repens</i> L. and <i>T.incarnatum</i> L.)	7, 4'-dihydroxyflavone, geraldone, formononetin, umbelliferone.	<i>nod</i> gene inducers	Peters et al., 1986; Djoedjevic et al., 1987; Hartwig et al., 1990
Pea (<i>Pisium sativum</i> L.)	Eriodictyol, aspigenin-7- <i>O</i> -gluciside, apigenin	<i>nod</i> gene inducers	Firmin et al., 1986
Faba bean (<i>Vicia faba</i> L.)	Naringenin, eriodictyol, apigenin, luteolin hesperetin	<i>nod</i> gene inducers	Zaat et al., 1987; Zaat et al., 1989
Soybean (<i>Glycine max</i>)	Genistein, genistein-7- <i>O</i> -glucoside, genistein-7- <i>O</i> -(6"- <i>O</i> -malonylglucoside), isoliquiritigenin, narigenin daidzein, daidzein-7- <i>O</i> -(6"- <i>O</i> -malonylglucoside), coumestrol, 7,4'-dihydroxyflavone	<i>nod</i> gene inducers	Van Brussell et al., 1986; Kosslak et al., 1987; Bassan et al., 1988; Sadowsky et al., 1988

Although there is a high potential for finding phenolic compounds with insecticidal properties in symbiotic legumes and other plant species, to date, only a few molecules controlling soil insects have been identified. The available literature does not seem to offer enough information on the mechanisms and the effect of molecules extracted from different symbiotic plants on harmful soil pests. The possibility to uncover more and use them in soil pest control seems to be considerable. For this to become true, extensive research programmes must be initiated to identify more phenolic plant products with insecticidal activity towards harmful soil insects.

CONCLUSION

Understanding new roles of phenolics and their functional groups is vital for better rhizosphere fertility, soil pest control and the overall production ecosystems. Better understanding of which plants give what type and amount of which phenolics to be incorporated in our cultural practices will eventually solve problems such as soil and water conservation, soil fertility, nutrients conservation, and soil pests management. However, in order to exploit the rich knowledge achieved and to merit the efforts used so far, collective action among researchers is required to

Table 3. Phenolics compounds from common legume roots potential in defence against soil borne pests in cropping systems.

Legume	Phenolic compound	Functional role	Reference
<i>Lotus pendunculatus</i>	Vestitol	Insect deterrent	Russel et al., 1978
Soybean (<i>Glycine max</i> L.)	Glyceollin	Insect deterrent, phytoalexin	Dakora and Phillips, 1996
Cowpea (<i>Vigna unguiculata</i> L.)	Medicarpin	Insect deterrent, phytoalexin	Dakora and Phillips, 1996
	Kievitone	Antifungal	Keen, 1975
Bean (<i>Phaseolus vulgaris</i> L.)	Phaseolin	Insect deterrent, phytoalexin	Dakora and Phillips, 1996
Lima bean (<i>Phaseolus lunatus</i> L.)	Coumestrol	Nematicide	Rich et al., 1977
Pigeon pea (<i>Cajanus cajan</i>)	Cajanin	Insect deterrent, phytoalexin	Dakora and Phillips, 1996

develop detailed genetic, biochemical and physiological programmes involving phenolic compounds. Furthermore, our research should aim at the most appropriate combination of factors for the best use of the phenolics to improve plant and beneficial microbial growth as well as better the biological environments sustaining other life types.

ACKNOWLEDGEMENT

This study was supported by the Cape Peninsula University of Technology through University Research Funds 2007/RP03.

REFERENCES

- Achenbach H, Stocker M, Constenla MA (1988). Flavonoids and other constituents of *Bauhinia manca*. *Phytochem.* 27: 1835-1841.
- Adams NR (1989). Phytoestrogens. In: Toxicants of plants Origin, Vol. 4, Phenolics, Cheeke PR, Ed CRC Press, Boca Raton, FL, Chap. 2.
- Adesanya SA, O'Neill MJ, Roberts MF (1986). Structure related Fungi toxicity of isoflavonoids. *Physiol. Molec. Plant Pathol.* 29: 95-103.
- Akhtar M (1995). Economic aspects of nematode disease-crop loss management. *Everyman's Science.* 30: 71-75.
- Akhtar M (1998). Biological control of plant parasitic nematodes by neem products in agricultural soil. *Appl. Soil Ecol.* 7: 219-223.
- Akhtar M, Alam MM (1993). Control of plant parasitic nematodes by 'Nimin'- an urea coating agent and some plant oils. *Zeitschrift fur Pflanzenkrankhei ten Pflanzenschutz.* 100: 337-342.
- Akhtar M, Mahmood I (1996). Control of plant parasitic nematodes with organic and inorganic amendments in agricultural soil. *Appl. Soil Ecol.* 4: 243-247.
- Akhtar M, Malik A (2000). Role of organic soil amendments and soil organisms in the biological control of plant-parasitic nematodes: a review. *Biores. Technol.* 74: 35-47.
- Allen PJ (1972). Specificity of the cis-isomers of inhibitors of uredospore germination in the rust fungi. *Proc. Natl. Acad. Sci. U.S.A.* 69: 3497-3500
- Anderson AJ (1988). Mycorrhizae-host specificity and recognition. *Phytopathology.* 78: 375-378.
- Badra T, Eligindi DM (1979). The relationship between phenolic content and *Tylenchulus semipenetrans* populations in nitrogen amended citrus plants. *Revue de Nematologie.* 2: 161-164.
- Baskin JM, Ludlow CJ, Harris TM, Wolf FT (1967). Psoralen, an inhibitor in the seed of *Psoralea subacaulis* (*Lequinrinosa*). *Phytochem.* 6: 1209-1213.
- Bassan BJ, Djordjevic MA, Redmond JW, Batley M and Rolfe BG (1988). Identification of *nodD*-dependent locus in the rhizobium, strain NGR234 activated by phenolic factor secreted by soybeans and other legumes. *Mol. Plant-Microbe Interact.* 1: 161-168.
- Beier RC (1990). Natural pesticides and bioactive compounds in foods. *Rev. Environ. Contam. Toxicol.* 113: 47-137.
- Beier RC, Ivie GW, Oertli EH (1983). Proralens as phytoalexins in food plant family umbeliferae: Significance in relation to storage and processing. In: Xenobiotics in Food and feeds. *Amer. Chem. Soc. Symposium Ser. Vol. 234*, Finely JW, Schwass DE (eds) *Amer. Chem. Soc., Washington, D.C.* p. 295.
- Bekkara F, Jay M, Viricel MR, Rome S (1998). Distribution of phenolic compounds within seed and seedlings of tow *Vicia faba* cvs differing in their tannin content, and study of their seed and root phenolic exudations. *Plant Soil.* 203: 27-36.
- Bending GD, Read DJ (1995). The structure and function of the vegetative mycelium of ectomycorrhizal plants. VI. Activities of nutrient mobilizing enzymes in birch litter colonized by *Paxillus involutus* (Fr.) Fr. *New Phytol.* 130: 411-417.
- Blum U (1998). Effects of microbial utilization of phenolic acids and their phenolic acid breakdown products on allelopathic interactions. *J. Chem. Ecol.* 24: 685-708.
- Boufalis A, Pellissier F (1994). Allelopathic effects of phenolic mixtures on respiration of two spruce mycorrhizal fungi. *J. Chem. Ecol.* 20: 2283-2289.
- Buchholtz KP (1971). The influence of Allelopathy on mineral nutrition. In: Biochemical interaction among plants, US National Committee for IBP, Eds., National Academy of Sciences, Washington, D.C.
- Callaway RM, Aschehoug ET (2000). Invasive plant versus their new and old neighbours: a mechanism for exotic invasion. *Science.* 290: 521-523.
- Carballeira A, Cuervo A (1980). Seasonal variation in allelopathic potential of soils from *Erica australis* L. heathland. *Acta Oecol.* 1: 345-353.
- Cartwright B, Hallsworth EG (1970). Effect of Cu deficiency on root nodules of subterranean clover. *Plant Soil.* 33: 685-698.
- Chou CH, Lin HJ (1976). Autointoxication mechanism of *Oryza sativa* L. Phytotoxic effects of decomposing rice residues in soil. *J. Chem. Ecol.* 2: 353-376.
- Christie T (1965). The effect of some phenolic compounds upon growth of two species of *Phytophthora*. *N. Z. J. Agric. Res.* 8:630-635.
- Crowden RK (1967). Biosynthesis of polyphonic acid metabolites of *Polyporus tumulosus* Cooke. *Can. J. Microbiol.* 13: 181-197.
- Curir P, Van Sumere CF, Termini A, Barthe P, Marchesini A, Dolci M (1990). Flavonoid accumulation is correlated with adventitious roots formation in *Eucalyptus gunnii* Hook micropropagated through axillary bud stimulation. *Plant Physiol.* 92: 1148-1153.
- Cutler HG (1988). Unusual plant growth regulators from microorganisms. *Crit. Rev. Plant Sci.* 6: 323-343.
- D'Arcy-Lameta A, Jay M (1987). Study of soybean and lentil root exudates III. Influence of soybean isoflavonoids on the growth of rhizobia and some rhizosphere microorganisms. *Plant Soil.* 101: 267-272.
- Dakora FD (1994). Nodulation gene induction and genetic control in the legume-*Rhizobium* symbiosis. *S. Afr. J. Sci.* 90: 596-599.
- Dakora FD (1995). Plant Flavonoids: Biological Molecules for Useful Exploitation. *Aust. J. Plant Physiol.* 22: 87-99.
- Dakora FD (2000). Commonality of root nodulation signals and nitrogen assimilation in tropical grain legumes belonging to the tribe *Phaseo-*

- leae. Aust. J. Plant Physiol. 27: 885-892.
- Dakora FD, Phillips DA (2002). Root exudates as mediators of mineral acquisition in low-nutrient environments. *Plant Soil*. 245: 35-47.
- Dakora FD, Joseph CM, Phillips DA (1993). Common bean root exudates contain elevated levels of daidzein and coumestrol in response to *Rhizobium* inoculation. *Mol. Plant-Microbe Inter.* 6: 665-668.
- Dakora FD, Joseph CM, Phillips DA (1993a). Alfalfa (*Medicago sativa* L.) root exudates contain isoflavonoids in the presence of *Rhizobium melliloti*. *Plant Physiol.* 101: 819-824.
- Dakora FD, Joseph CM, Phillips DA (1993b). Common bean root exudates contain elevated levels of daidzein and coumestrol in response to *Rhizobium inoculation*. *Mol. Plant-Microbe Interact.* 6: 665-668.
- Dakora FD, Phillips DA (1996). Diverse functions of Isoflavonoids in legumes transcend ant-microbial definitions of phytoalexins. *Physiol. Mol. Plant Pathol.* 49: 1-20
- DeKock PC, Vaughan D (1975). Effect of some chelating and phenolic substances on the growth of excised pea root segments. *Planta*. 126: 187-195.
- Deladonde M, Barret Y, Coumans MP (1996). Development of phenolic compounds in maize anthers (*Zea mays*) during cold pre-treatment prior to endogenesis. *J. Plant Physiol.* 149: 612-616.
- Delhaize E, Rayan PR, Randall PJ (1993). Aluminium tolerance in wheat (*Triticum aestivum* L.). 2. Aluminium stimulated excretion of malic acid from root apices. *Plant Physiol.* 103: 695-702.
- Djordjevic MA, Redmond JW, Batley M, Rolfe BG (1987). Clovers secrete specific phenolic compounds, which either stimulate or repress *nod* gene expression in *Rhizobium trifolii*. *EMBO J.* 6: 1173-1179.
- Duncan LW (1991). Current options for nematode management. *Ann. Rev. Phytopathol.* 29: 467-490.
- Egunjobi OA, Afolami SO (1976). Effect of Neem (*Azadirachta indica*) leaf extracts on populations of *Pratylenchus brachyurus* and on the growth and yield of maize. *Nematologica*. 22: 125-132.
- Einhellig FA (1986). Mechanisms and mode of action of allelochemicals. In: The science of allelopathy, Putnam AR, Tang CS Eds. John Wiley and Sons, New York, Chap.10.
- Elliott LF, Cheng HH (1987). Assessment of Allelopathy among microbes and plants. In: Allelopathy: Role of agriculture and Forestry, Walker GR (eds) ACS Symposium. Ser., American Chemical Society, Washington, D.C. Chapter 45.
- Ernst WHO, Verkleij JAC, Schad H (1992). Metal tolerance in plants. *Acta Bot. Neerl.* 41(3): 229-248.
- Eva C, Josep P (2003). Is there a feedback between N availability in siliceous and calcareous soils and *Cistus albidus* leaf chemical composition. *Oecol.* 136: 183-192.
- Faudin AS, Macko V (1974). Identification of the self inhibitor and some germination characteristics of peanut rust uredospores. *Phytopathol.* 64:990-993
- Fierer N, Schimel JP, Cates RG, Zou J (2001). Influence of balsam poplar tannin fractions on carbon and nitrogen dynamics in Alaskan taiga floodplain soils. *Soil Biol. Biochem.* 33: 1827-1839.
- Firmin JL, Wilson KE, Rossen L, Johnstone AWB (1986). Flavonoid activation of nodulation genes in *Rhizobium* reversed by other compounds present in plants. *Nature (London)*. 324: 90-92.
- Fisher RF (1987). Allelopathy: a potential cause of forest regeneration failure. In: Allelochemicals, Role in agriculture and Forestry, Walker GR (eds) ACS Symposium. Ser., Am. Chemical Soc. Washington, D.C. Chapter 16.
- Fottrell PF, O'Conner S and Masterson CL (1964). Identification of the flavonol myricetin in legume seed and its toxicity to nodule bacteria. *Ir. J. Agric. Res.* 3: 246-249.
- French RC, Graham CL, Gale AW, Long RK (1977). Structural and exposure time requirement for chemical stimulation of germination of uredospores of *Uromyces phaseoli*. *J. Agric. Food Chem.* 25: 84-88.
- Gerke J (1992). Phosphate, Aluminium and ion in the soil solution of three different soils in relation to varying concentrations of citric acid. *Z. Pflanzenenahr. Bodenk.* 155: 339-343.
- Gerke J (1994). Kinetics of soil phosphate desorption as affected by citric acid. *Z. Pflanzenenahr Bodenk.* 157: 17-22.
- Guenzi WD, McCalla TM (1966). Phenolic acids in oats, wheat, sorghum and corn residues and their phytotoxicity. *Agron. J.* 58: 303-304.
- Guenzi WD, McCalla TM (1966). Phytotoxic substances extracted from soil. *Soil Sci. Amer. Proc.* 30: 214-216.
- Haider K, Martin JP, Filip Z (1975). Humus biochemistry. In: Soil Biochemistry, Vol. 4, Paul EA Ed., Marcel Dekker, New York. Chap. 6.
- Harborne JB (1980). Plant phenolics. In: Secondary plant products. Bell EA and Charlwood BV (eds) Springer Verlag, Berlin, chap. 6.
- Hartley RD, Harris PJ (1981). Phenolic constituents of the cell walls of dicotyledons. *Biochem. Syst. Ecol.* 9: 189-203.
- Hartley RD, Whitehead DC (1985). Phenolic acids in soils and their influence in growth and soil microbial processes. In: Soil organic matter and Biological Activity, Vaughan D, Malcolm RE (eds) Martinus Nijhoff/Dr W Junk Publishers, Dordrecht.
- Hartwig UA, Maxwell CA, Joseph CM, Phillips DA (1990). Chrysoeriol and luteolin released from alfalfa seeds induce *nod* genes in *Rhizobium melliloti*. *Plant Physiol.* 92: 116-122.
- Hattenschwiler S, Vitousek PM (2000). The role of polyphenols in terrestrial ecosystem nutrient cycling. *Tree*. 15: 238-243.
- Hoenschwiler S, Vitousek PM (2000). The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends Ecol. Evol.* 15: 238-243.
- Hoffland E (1992). Quantitative evaluation of the role of organic-acid exudation in the mobilization of rock phosphate by rape. *Plant Soil*. 140: 279-289.
- Hoffland E, Van den Boogaard R, Nelemans J, Findenegg G (1992). Biosynthesis and root exudation of citric and malic acids in phosphate starved rape plants. *New Phytol.* 122: 675-680.
- Horner JD, Gosz JR, Cates RG (1988). The role of carbon-based plant secondary metabolites in decomposition in terrestrial ecosystems. *Am. Nat.* 132: 869-883.
- Horvath B, Bachem C, Schell J and Kondorosi A (1987). Host specific regulation of nodulation genes in *Rhizobium* is mediated by a plant signal interacting with the *nodD* gene product. *EMBO J.* 6: 841-848.
- Hungria M, Joseph CM, Phillips DA (1991). Anthocyanins and flavonols, major *nod*-gene inducers from seed of black seeded common bean (*Phaseolus vulgaris* L.). *Plant Physiol.* 97: 751-758.
- Imas P, Bar Yosef B, Kafkafi U, Ganmore-Neumann R (1997). Phosphate induced carboxylate and proton release by tomato roots. *Plant Soil*. 191: 35-39.
- Inderjit (1996). Plant phenolics in allelopathy. *Bot. Rev.* 62: 186-202.
- Innes RW, Kuempel PL, Plazinski J, Canter-Cremers H, Rolfe GB, Djordjevic MA (1985). Plant factors induce expression nodulation and host range genes in *Rhizobium trifolii*. *Mol. Gen. Genet.* 201: 426-432.
- Johnson JF, Allan DL, Vance CP, Weiblen G (1996). Root carbon dioxide fixation by phosphorus deficient *Lupinus albus*: Contribution to organic acid exudation by proteoid roots. *Plant Physiol.* 112: 19-30.
- Jung C, Maeder V, Funk F, Frey B, Sticher H, Frossard E (2003). Release of phenols from *Lupinus albus* L. roots exposed to Cu and their possible role in Cu detoxification. *Plant Soil*. 252: 301-312.
- Kandasamy D, Prasad NN (1979). Colonisation of rhizobia of the seed and roots of legumes in relation to exudation of phenolics. *Soil Biol. Biochem.* 11: 73-75.
- Kapulnik Y, Joseph CM, Phillips DA (1987). Flavone limitation to root nodulation and symbiotic nitrogen fixation in alfalfa. *Plant Physiol.* 84: 1193-1196.
- Kasenberg TR and Traquar JA (1988). Effects of phenolics on growth of *Fusarium oxysporum* f. sp. *radicis lycopersici* *in vitro*. *Can. J. Bot.* 66: 1174-1177.
- Katase T (1981). The different forms in which *p*-coumaric acid exists in a peat soil. *Soil Sci.* 131: 271-275.
- Kevers, Coumans M, Coumans Gilles M, Gasper T (1984). Physiological and Biochemical events leading to vitrification of plants cultured *in vitro*. *Physiol. Plant.* 61:69-74.
- Kidd PS, Llugany M, Poschenrieder C, Gunse B, Barcelo J (2001). The role of root exudates in Aluminium resistance and silicon-induced amelioration of aluminium toxicity in three varieties of maize (*Zea mays* L.). *J. Exp. Bot.* 52: 1339-1352.
- Klein K, Blum U (1990). Inhibition of cucumber leaf expansion by ferulic acid in spit root experiments. *J. Chem. Ecol.* 16: 455-463.
- Kochian LV (1995). Cellular mechanisms of Aluminium toxicity and resistance in plants. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 46: 237-260.

- Kochian LV, Jones DL (1997). Aluminium toxicity and resistance in plants. In: Research issues in Aluminium toxicity (eds) Yokel RA and Golub MS Taylor and Francis Publishers, Bristol, PA.
- Kosslak RM, Bookland R, Barkei J, Paaren HE, Appelbaum E. (1987). Induction of *Bradyrhizobium Japonicum* common *nod* gene by isoflavones isolated from *Glycine max*. Proc. Natl. Acad. Sci. USA. 84: 7428-7432.
- Kovacs MF Jr (1971). Identification of aliphatic and aromatic acids in root and seed exudates of peas, cotton, and barley. Plant Soil. 34: 441-451.
- Kraus TEC, Dahlgren RA, Zasoski RJ (2003). Tannins in nutrients dynamics of forest ecosystems-a review. Plant Soil. 256: 41-66.
- Laheurte F, Berthelin J (1988). Effects of phosphate solubilizing bacteria on maize growth and root exudation over four levels of labile phosphorus. Plant Soil. 105: 11-17.
- Lane GA, Sutherland ORW, Skipp RA (1987). Isoflavonoids as insect feeding deterrents and antifungal compounds from roots of *L. angustifolius* L. J. Chem. Eco. 13(4): 771-783
- Lawson CGR, Rolfe BG, Djordjevic MA (1996). *Rhizobium* inoculation induces condition dependent changes in the flavonoid composition of root exudates from *Trifolium subterraneum*. Aust. J. Plant Physiol. 23: 93-101.
- Leather GR, Einhellig FA (1986). Bioassay in the study of allelopathy. In: The science of allelopathy, Putnam AR and Tang CS (eds) John Wiley and Sons, New York, Chap.8.
- Leon-Barrios M, Dakora FD, Joseph CM, Phillips DA (1993). Isolation of *Rhizobium meliloti nod* gene inducers from alfalfa rhizosphere soil. Appl. Environ. Microbiol. 59: 636-639.
- Lingappa BT, Lockwood JL (1962). Fungitoxicity of lignin monomers, model substances, and decomposition products. Phytopathology 52:295-299.
- Lodhi MAK (1975). Soil-plant phytotoxicity and its possible significance in patterning of herbaceous vegetation in a bottomland forest. Am. J. Bot. 62: 618-622.
- Long SR (1989). Rhizobium legume regulation: Life together in the underground. Cell. 56: 203-214.
- Ma JF, Hiradate S, Nomoto K, Iwashita T, Matsumoto H (1997). Internal detoxification mechanism of Aluminium in Hydrangea – Identification of Al form in the leaves. Plant Physiol. 113: 1033-1039.
- Marschner H (1988). Mechanism of Mn acquisition by roots from soils. In: manganese in soils and Plants Graham RD, Hannam RJ and Uren NC (eds), pp. 191-204. Kluwer Academic, Dordrecht, The Netherlands
- Marschner H (1995). Mineral Nutrition of Higher plants. Academic Press, London.
- Martin F (1977). Biological properties of soils, In: Soils for Management of Organic Wastes and Waste Water Elliot LF and Stevenson FJ (eds) American Society of Agronomy, Madison, WI, Chap. 20.
- Martin JP, Heider K (1976). Decomposition of specifically carbon-14-labelled ferulic acid: free and linked into model humic acid-type polymers. Soil Sci. Soc. Amer. 40: 377-380.
- Martin JP, Heider K, Wolf D (1972). Synthesis of phenol and phenolic polymers by *Hendersonula toruloidea* in relation to humic acid formation. Soil Sci. Soc. Amer. Proc. 36: 311-315.
- Masaoka Y, Kojima M, Sugihara S, Yoshihara T, Koshino M, Ichihara A (1993). Dissolution of ferric phosphate by alfalfa (*Medicago sativa* L.) root exudates. Plant Soil. 155/156: 75-78.
- Maxwell CA, Hartwig UA, Joseph CM, Phillips DA (1989). A chalcone and two related flavonoids released from Alfalfa seeds and roots induced *nod* genes of *Rhizobium meliloti*. Plant Physiol. 91: 842-847.
- Micales JA (1997). Localization and induction of oxalate decarboxylase in the brown rot wood decay fungus *Postia placenta*. Int. Biodereioration Biodegrad. 39: 125-132.
- Morandi D (1989). Effect of Xenobiotics on endomycorrhizal infection and isoflavonoid accumulation in soybean roots. Plant Physiol. Biochem. 27: 697-701.
- Morandi D, Bailey JA (1984). Isoflavonoids accumulation in soybean roots infected with vesicular-arbuscular mycorrhizal fungi. Physiol. Plant Pathol. 24: 357-364.
- Morel C, Hinsinger P (1999). Root induced modifications of the exchange of phosphate ion between soil solution and solid phase. Plant Soil. 211: 103-110.
- Morris HR, Taylor GW, Jermyn KA, Kay RR (1987). Chemical structure of the morphogen differentiation inducing factor from *Dictyostelium discoideum*. Nature. 328: 811-814.
- Muller CH (1969). Allelopathy as a factor in ecological process. Vegetation. 18: 348-357.
- Muller R, Gooch PS (1982). Organic amendment in nematode control. An examination of the literature. Nematropica. 12: 319-326.
- Mulligan JT, Long SR (1985). Induction of *Rhizobium meliloti nodC* expression by plant exudate requires *nodD*. Proc. Natl. Acad. Sci. USA. 82: 6609-6613.
- Ndakidemi PA, Dakora FD (2003). Legume seed flavonoids and nitrogenous metabolites as signals and protectants in early seedling development. Review. Functional Plant Biol. 30: 729-745.
- Nicolai V (1988). Phenolic and mineral content of leaves influences decomposition in European forest ecosystems. Oenology. 75:575-579.
- Northup RR, Dahlgren RA, McColl JG (1998). Polyphenols as regulators of plant-litter-soil interactions in northern California's pygmy forest: a positive feedback? Biogeochem. 42: 189-220.
- Northup RR, Yu Z, Dahlgren RA, Vogt KA (1995). Polyphenol control of nitrogen release from pine litter. Nature. 377:227-229.
- Oades JM (1988). The retention of organic matter in soils. Biogeochem. 5: 35-70.
- Osen RA, Bennett JH, Blume D, Brown JC (1981). Chemical aspects of the Fe stress response mechanisms in tomatoes. J. Plant Nutr. 3: 905-921.
- Palm CA, Sanchez PA, (1990). Decomposition and nutrient release patterns of the leaves of three tropical legumes. Biotropica. 22: 330-338.
- Pankhurst CE, Biggs DR (1980). Sensitivity of *Rhizobium* to selected isoflavonoids. Can. J. Microbiol. 26: 542-545.
- Parfitt RL (1979). The availability of P from phosphate-goethite bridging complexes. Description and uptake by ryegrass. Plant Soil. 53: 55-65.
- Patrick ZA (1971). Phytotoxic substances associated with the decomposition in soil plant residue. Soil Sci. 111: 13-18.
- Patterson DT (1987). Effects of allelopathic chemicals on growth and physiological responses of soybean (*Glycine max*). Weed Sci. 29:53-59.
- Paula MA, Siqueira JO (1990). Stimulation of hyphal growth of the VA mycorrhizal fungus *Gigaspora margarita* by suspension cultured *Pueraria phaseoloides* cells and cell products. New Phytol. 155: 69-75.
- Peters NK, Frost JW, Long SR (1986). A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. Science. 233: 977-980.
- Phillips DA (1992). Plant root signals to soil microbes. Recent Adv. Phytochem. 26: 201-231.
- Phillips DA, Dakora FD, Sande E, Joseph CM, Zon J (1994). Synthesis, release, and transmission of alfalfa signals to rhizobial symbionts. Plant Soil. 161: 69-80.
- Phillips DA, Wery J, Joseph C, Jones AD, Teuber, LR (1995). Release of flavonoids and betaines from seeds of seven *Medicago* species. Crop Sci. 35: 805-808.
- Powel H, Rate AW (1987). Aluminium-tannin equilibrium: a potentiometric study. Aust. J. Chem. 40: 2015-2022.
- Putnam AR, Tang CS (1986). The Science of Allelopathy, John Wiley and Sons, New York.
- Ravin H, Andary C, Kovacs G, Molgaard P (1989). Caffeic acid esters as *in vitro* inhibitors of plant pathogenic bacteria and fungi. Biochem. Syst. Ecol. 17: 175-184.
- Recourt K, Schripsema J, Kijne JW, Van Brussel AAN, Lugtenberg BJJ (1991). Inoculation of *Vicia sativa* subsp. *nigra* roots with *Rhizobium leguminosarum* biovar *viciae* results in release of *nod* gene activating flavonones and chalcones. Plant Mol. Biol. 16: 841-852.
- Reddy MN, Rao AS, Rao KN (1975). Production of phenolic compounds by rhizoctonia. Trans. Br. Mycol. Soc. 64: 146.
- Redmond JW, Batley M, Innes RW, Kuempel PL, Rolfe GB, Djordjevic MA (1986). Flavones induce expression of nodulation gene in *Rhizobium*. Nature (London). 323: 632-635.
- Rice EL (1984). Allelopathy. Academic Press, Orlando, FL.
- Rich JR, Keen NT, Thomason IJ (1977). Associations of coumestans

- with the hypersensitivity of lima bean roots to *Pratylenchus scribneri*, *Physiol. Plant Pathol.* 10: 105-116.
- Richards A, Christopher LS (1999). Flavonoids and isoflavonoids – a gold mine for metabolic engineering. *Trends Plant Sci.* 4(10): 394-400.
- Robbins MP, Thomas B, Morris P (1995). Phenylpropanoid defence responses in transgenic *Lotus corniculatus*. II Modelling plant defence in transgenic root cultures using thiol and carbohydrate elicitors. *J. Exp. Bot.* 46: 513-524.
- Rossen L, Shearman CA, Johnston AWB, Downie JA (1985). The *nodD* gene of *Rhizobium leguminosarum* is autoregulatory and in the presence of plant exudate induces the *nodA*, *B*, *C* genes. *EMBO J.* 4: 3369-3373.
- Russel GB, Sutherland ORW, Hutchinson RFN, Christmas PE (1978). Vestitol: a phytoalexin with insect feeding-deterrent activity. *J. Chem. Ecol.* 4: 571-580.
- Sadowsky MJ, Gregan PB, Keyser HH (1988). Nodulation and nitrogen fixation efficiency of *Rhizobium fredii* with *Phaseolus vulgaris* genotypes. *Appl. Environ. Microbiol.* 54: 1907-1910.
- Scalbert A, Miller I, Expert D, Marmolle F, Albretch A, Hurrel R, Huneau J, Tome D (1999). Polyphenols, metal ion complexation and biological consequences. In: *Plant polyphenols 2. Chemistry, Biology, Pharmacology, Ecology* Gross GG, Hemingway RW, Yoshida R, Branham S (eds) pp. 545-553.
- Schenk HJ, Callaway RM, Mahal BE (1999). Spatial root segregation: are plants territorial. *Adv. Ecol. Res.* 28: 146-180.
- Schimel JP, van Cleve K, Cates RG, Clausen TP, Reichardt PB (1996). Effects of balsam poplar (*Populus balsamifera*) tannins and low molecular weight phenolics on microbial activity in taiga floodplain soil: implications for changes in N cycling during succession. *Can J. Bot.* 74: 84-90.
- Schmidt IK, Michelsen A, Jonasson S (1997). Effect of labile soil carbon on nutrient partitioning between arctic graminoid and microbes. *Oecol.* 112: 557-565.
- Seneviratne G, Jayasingheachchi HS (2003). Phenolic acids: Possible agents of modifying N₂-fixing symbiosis through rhizobial alteration. *Plant Soil.* 252:385-395.
- Shafer SR, Blum U (1991). Influence of phenolic acids on microbial populations in the rhizosphere of cucumber. *J. Chem. Ecol.* 17: 369-388.
- Siqueira JO, Nair MG, Hammerschmidt R, Safir GR (1991). Significance of phenolic compounds in Plant-Soil-Microbial systems. *Cri. Rev. Plant Sci.* 10(1): 63-121.
- Siqueira JO, Safir GR, Nair MG (1991). Simulation of VAM formation and growth of white clover by flavonoid compound. *New Phytol.* 118:87-93.
- Slabbert N (1992). Complexation of tannins with metal ions. In: *plant polyphenols. Synthesis, properties, significance.* Eds. RW Hemingway, PE Laks pp. 421-436. Plenum press New York.
- Smith MM, Hartley RD (1983). Occurrence and nature of ferulic acid substitution of cell walls polysaccharides in graminaceous plants. *Carbohydr. Res.* 118: 65-80.
- Smith SE, Read DJ (1997). 'Mycorrhizal symbiosis.' (Academic press Ltd: London)
- Spaink HP, Wijffelman CA, Pees E, Okker RJH, Lugtenberg BJJ (1987). Rhizobium nodulation gene *nodD* as a determinant of host specificity. *Nature (London).* 328: 337-340.
- Sparling GP, Ord BG, Vaughan D (1981). Changes in microbial biomass and activity in soils amended with phenolic acids. *Soil Biol. Biochem.* 13: 455-460.
- Sugai SF, Schimel JP (1993). Decomposition and biomass incorporation of ¹⁴C-labeled glucose and phenolics in taiga forest-floor: effect of substrate quality, successional state, and season. *Soil Biol. Biochem.* 25: 379-1389.
- Takahama U, Oniki T (1992). Regulation of peroxidase dependent oxidation of phenols in the apoplast of spinach leaves by ascorbate. *Plant Cell Physiol.* 33: 379-387.
- Thakur RS, Singh BS, Goswami A (1981). Review article. E. *Azadirachta indica* A. Juss. CROMAP. 3.
- Thompson JA (1988). Survival of root-nodule bacteria on inoculated seed. In: *Microbial in action*, Murrell, W.G. and Kennedy, I.R., eds., John Wiley and sons, New York, Chap 5.
- Tiarks AE, Bridges JR, Hemingway RW, Shoulders E (1989). Condensed Tannins in southern pines and their interactions with the ecosystem. In: *Chemistry and significance of condensed Tannins* Hemingway RW and Karchesy JJ (eds) pp. 369-390. Plenum Press New York.
- Tsai SM, Phillips DA (1991). Flavonoids released naturally from alfalfa promote development of symbiotic *Glomus* spores *in vitro*. *Appl. Environ. Microbiol.* 57: 85-88
- Turner JA, Rice EL (1975). Microbial decomposition of ferulic acid in soil. *J. Chem. Ecol.* 1: 41-58.
- Van Brussell, AAN, Zaat SAJ, Canter Cremers HCJ, Wijffelman CA, Pees E, Tak T and Lugtenberg BJJ (1986). Role of plant root exudates and sym plasmid localized nodulation genes in the synthesis by *Rhizobium leguminosarium* of Tsr factor, which causes thick and short roots on common vetch. *J. Bacteriol.* 165: 517-529.
- Vidhyasekaran P (1988). Physiology of disease resistance in plants, Vol. 1, CRC Press, Boca Raton, FL.
- Waller GR (1987). Allelochemicals role in agriculture and forestry, ACS symposium. 330, American Chemical society, Washington, D.C.
- Waterman PG, Mole S (1989). Extrinsic factors influencing production of metabolites in plants. In: *Insect-Plant interactions* Bernays EA (eds) CRC Press, Boca Raton, FL, Chap. 4.
- Whitehead DC (1964). Identification of *p*-hydroxybenzoic vanillic, *p*-coumaric and ferulic acid in soils. *Nature.* 202: 417-418.
- Whitehead DC, Buchan H, Hartley RD (1979). Composition and decomposition of roots of ryegrass and red clover. *Soil Biochem.* 11: 619-628.
- Wymann JG, VanEtten HD (1978). Antibacterial activity of selected isoflavonoids. *Phytopathology* 68: 583-589.
- Wyss P, Boller T, Wiemken A (1989). Glyceollin production in soybean during the process of infection by *Glomus mosseae* and *Rhizoctonia solani*. *Agric. Ecosyst. Environ.* 29: 451-458.
- Zaat SAJ, Wijffelman CA, Spaink HP, Van Brussel AAN, Okker RJH, Lugtenberg BJJ (1989). Analysis of the major inducers of the *Rhizobium nodA* promoter from *Vicia sativa* root exudates and their activity with different *nodD* genes. *Plant Mol. Biol.* 13: 175-188.
- Zaat SAJ, Wijffelman CA, Spaink HP, Van Brussel AAN, Okker RJH, Lugtenberg BJJ (1987). Induction of *nodA* promoter of *Rhizobium leguminosarum* Sym plasmid pRL1J1 by plant flavonones and flavones. *J. Bacteriol.* 169: 198-204.
- Zawoznik MS, Garrido LM, del Pero Martinez MA and Tomaro ML (2000). Effect of vanillin on growth and symbiotic ability of *Bradyrhizobium* sp. (Arachis) strain. In: *Proc. Int. Plant growth-Promoting Rhizobacteria* Electronic publication (<http://www.ag.uburn.edu/argentina>).
- Zaynoun ST, Aftimos BG, Ali LA, Tenekjiankk KK, Khalid U, Kurban AK (1984). *Ficus carica*, isolation and quantification of the photoactive components. *Contact Dermatit.* 11: 21-25.