

Full Length Research Paper

Response of microalgae from mud-flats to petroleum hydrocarbons in the presence of nitrogenous fertilizer effluents

A. Adoki

Health Safety and Environment, Shell Petroleum Development Co. Ltd, P.O. Box 263, Old Aba Road, Port Harcourt, Nigeria. E-mail: akuro.adoki@shell.com.

Accepted 6 February, 2008

Untreated industrial waste (effluent) from out-fall of the National fertilizer company of Nigeria (NAFCON) was found to be weakly alkaline with a pH of 7.8. Total dissolved solids and sulphate were relatively high with levels of 2052 and 1599 mg/l, respectively. Nitrogen as ammonia was high in comparison to nitrate nitrogen. Phosphate was also found to be low in concentration. Using indigenous microalgae and aeration, there was a reduction in nutrients; NH_4^+ , NO_3^- , PO_4^{3-} and SO_4^{2-} . However, in the presence of the hydrocarbon, there was delayed nutrient uptake. The consequence of this response was the observed reductions in chlorophyll content, biomass, etc. There was however a quick recovery in the oiled systems.

Key words: Industrial, effluent, petroleum, mudflat, microalgae, nitrogenous, fertilizer, response.

INTRODUCTION

The increasing deterioration of the environment and the need for energy and food is forcing the exploration of the feasibility of wastewater recycling and resource recovery. Within this context, bio-treatment with microalgae has been particularly attractive because of their photosynthetic capabilities converting solar energy into useful biomass and incorporating nutrients such as nitrogen and phosphorus which cause pollution and eutrophication (De la Noue and De Pauw, 1988). They play a major role as producers in waste stabilization ponds and other polluted aquatic environments. Microalgae thus have been used extensively for domestic as well as industrial waste water treatment. Emphasis may be put on wastewater treatment and/or algae biomass production. Processes involved in the removal of nutrients are precipitation, stripping and uptake by algal biomass. The efficiency of removal of a particular nutrient (for example N or P) will also depend on whether or not these nutrients are limiting in the wastewater to be treated. In practice, with adequate stirring, more than 90% nitrogen and/or phosphorus can be removed (Shelef et al., 1980).

During the past years, various systems and species, mainly unicellular species have been investigated in order to improve the efficacy and economic feasibility of

such bio-treatment systems. Soler et al. (1991) studied temporal changes in physico-chemical parameters and photosynthetic microorganisms in a self-depuration waste water lagoon. In the study, some phytoplanktons played vital roles in the bio-treatment or stabilization. These microalgae include *Chlamydomonas* sp., chlorophytes mainly of the genera *Golenkia*, *Chlorella* and *Scenedesmus* and the cyanophyte of the genus *Synechococcus*. These microalgae were characterized by proliferation with corresponding reduction of nutrients levels such as PO_4^{3-} and NH_4^+ .

A large number of nitrogenous fertilizer industries refining, coking and organic chemical industries generate high concentrations of ammonia and urea. Effluent discharges from these sources are the main cause of eutrophication – the fertilization of lakes and waterways by nitrogen and phosphorus compounds. Eutrophication occurs naturally as nutrients move from field to inland waters, but the use of inorganic or organic nitrogen in agriculture and release from industries undoubtedly enhance the enrichment of those bodies of water in which these elements limit photosynthetic populations. Only a low concentration of nitrate is necessary, and lakes with 0.3 ppm nitrate nitrogen sometimes may support unwanted

ted algal blooms, provided other nutrients are in adequate supply (Alexander, 1977).

Excessive blooms result in high cost of water treatment, fish kills as oxygen is depleted, offensive odour and bad taste of water and navigational problems for ships. Because of toxic and other effects on aquatic life, in addition to deterioration of water quality, the limit of discharging of waste water containing ammoniacal nitrogen in aquatic environments was fixed at 50 mg N/L (Gupta and Sharma, 1996) in India. Thus there is need to remove ammoniacal nitrogen prior to discharge. Biological nitrification-denitrification is the process commonly used to accomplish this objective. The use of aquatic plants as nutrient removal agents has also been effective. The water hyacinth, *Eichhornia crassipes*, is the most studied in this aspect. It has been shown to remove 40 – 45% of the orthophosphates in water effluents in 25 – 30 days and also 94% removal of the nitrate–nitrogen (NO_3^- -N) and NH_3 -N in 10 days (Cornwell et al., 1977).

Microalgae have been identified in environments polluted with oil. An estimated 6.1 million metric tons of hydrocarbons enter the marine environment annually (Alexander and Schwarz, 1980). While some oil seepage may occur naturally, the current pollution of the oceans by oil at the estimated rate of 5 million metric tons annually is clearly a consequence of man's activities (Atlas and Bartha, 1973). The local impact of major, or even small oil spills is well documented (Blumer et al., 1971; Atlas and Bartha, 1973) but such accidents contribute only about 4% of the total oil pollution load of the oceans (Atlas and Bartha, 1973). The remaining 96% are contributed by 'normal' operations in the production, transportation and disposal of petroleum and its refined fractions

A considerable number of studies have been carried out on the biodegradation of these hydrocarbons on terrestrial soils (Jobson et al., 1974; Westlake et al., 1978) and in marine and estuarine waters and sediments (Blumer et al., 1971; Atlas and Bartha, 1973; Hollaway et al., 1980; Colwell and Walker, 1977; Gibson et al., 1975; Schwartz et al., 1974).

There have also been studies on eutrophication, waste water treatment and biomass production using microalgae cultures (De la Noue and De Pauw, 1988; Soler et al., 1991; Azov and Shelef, 1982; Fallowfield and Garret, 1985; Shelef et al., 1980). These studies have been necessitated by the need for environmental protection and preservation, domestic and industrial wastewater treatment and also resource recovery.

However, studies on the response of the microalgae which are the primary producers in these environments, to crude pollution in the presence of nitrogenous fertilizer effluents has not been given much attention.

The objective of this study, therefore, was: 1. To study the various aspects of microalgae composition and dynamics i.e. species diversity (distribution and succession) that are involved in the treatability of the NAFCON efflu-

ent. 2. Evaluate the nutrient levels and other physico-chemical parameters that affect self-depuration or biotreatability of the NAFCON effluent and 3. Study the response of the microalgae to different levels of crude pollution in the presence of the NAFCON effluent.

MATERIALS AND METHODS

Effluent waste water and sediment

The industrial wastewater (effluent) and sediment were obtained from the outfall point of the National Fertilizer Company of Nigeria, (NAFCON), Onne, Rivers State. Samples of Bonny light crude used in this study were obtained from the Shell Petroleum Development Company (SPDC) flow station at Agbada near Port Harcourt.

Collection of samples

The NAFCON industrial wastewater (effluent) was collected in clean 25 L plastic containers at the out-fall of the NAFCON plant. The sediment samples were collected at low tide near the out-fall in open plastic containers while crude petroleum was supplied in a four-litre metal container.

Chemical and reagents

All the chemicals and reagents used in this study were of analytical grade and obtained from the Institute of Pollution Studies (I.P.S.) Laboratory (1986), the Chemistry and the Biology Department Laboratories, all of the Rivers State University of Science and Technology, Nkpolu, Rivers State.

Reactor vessels

The Institute of Pollution Studies (I.P.S.) Laboratory of the Rivers State University of Science and Technology, Port Harcourt provided eight Coca-Cola concentrate glass bottles of 3.5- L capacity.

Light source

The light source was artificial illumination from two parallel 4 ft white fluorescent lamps emitting about $15 \mu \text{E m}^{-2}/\text{s}$ of light each and mounted in a box-like chamber behind the reactor vessels.

Aeration and agitation

Aeration and agitation were by means of three aquarium aerators /air pumps (Tecax AP 1500 models). They were linked to one another by means of plastic Teflon tubes of about 6 mm diameter.

Experimental design

2 L of the NAFCON effluent collected was introduced into each of the eight Coca-Cola concentrate bottles (3.5 L capacity). The bottles were aerated by means of three aquarium air pumps and illuminated by two parallel 4 ft fluorescent lamps emitting about $15 \mu \text{E m}^{-2} \text{S}^{-1}$ light and left at ambient temperature in the laboratory (plate 1). Different levels of crude 0.2, 0.6, 0.8 and 1.0% were introduced into the vessels whilst another four without crude served

Table 1. Physico-chemical characteristics of untreated fertilizer factory effluent.

Parameter	Value
pH	7.80
Temperature	34°C
Conductivity	3537 μ mhos/cm
Total dissolved solids (T.D.S.)	2052 mg/L
Ammonia	2.98 mg/L
Nitrate	0.35 mg/L
Phosphate	0.25 mg/L
Sulphate	1599 mg/L

as control. 20 g of sediment was introduced into each vessel. The vessels were set up as follows:

A	medium contained only sediment
B	medium contained only sediment
C	medium contained only sediment
D	medium contained only sediment
AA	medium contained only sediment and 0.2% crude petroleum
BB	medium contained sediment and 0.6% crude petroleum
CC	medium contained sediment and 0.8% crude petroleum
DD	medium contained sediment and 1.0% crude petroleum

Analysis

The NAFCON effluent and sediment were analysed on the day of collection, a day after that and subsequently, weekly for nine weeks for each of the oil concentrations. The following parameters were monitored, using modified methods from the standard methods for water and waste water (APHA, 1985): pH, temperature, conductivity, total dissolved solids (T.D.S.), Ammonia, (NH_3), nitrate (NO_3^-), sulphate (SO_4^{2-}), Phosphate (PO_4^{3-}) and biomass estimations.

Biomass determination and cell numbers

The biomass (growth) was determined as dry weight, chlorophyll a estimation and the number of cells (cell count).

Viable cell counts

The counting of cells was done using the improved Neubauer cytometer (counting chamber).

Dry weight

50 ml of sample was harvested by centrifugation at 3,000 r.p.m in a bench-top centrifuge for 15 m. The cells were washed three times with distilled water, transferred to a pre-weighed filter paper and dried to a constant weight in a hot air oven at 70°C.

Chlorophyll estimation

The chlorophyll content was measured according to the standard method of water and wastewater analysis (APHA, 1985).

Identification of microalgal cells- algal diversity

Cells were identified, using the x 10 and x 40 objectives of the microscope (HM-Lux Leitz Wetzlar, Germany), on the basis of cell morphology and colonial characteristics (Round, 1973; APHA, 1985).

RESULTS AND DISCUSSION

Physico-chemical parameters

Untreated industrial waste (effluent) from out-fall of the National fertilizer company of Nigeria (NAFCON), had the characteristics outlined in Table 1. The effluent at collection was found to be weakly alkaline with a pH of 7.8. Total dissolved solids and sulphate were relatively high with levels of 2052 and 1599 mg/l, respectively. Nitrogen as ammonia was high in comparison to nitrate nitrogen. Phosphate was also found to be low in concentration.

Nutrient uptake studies

In solutions containing NH_4NO_3 , ammonia is usually preferentially absorbed and the pH of the medium falls. Assimilation of nitrate ions on the other hand tends to raise the pH. Marine phytoplankton has been shown to contain nitrogen in proportion to carbon and phosphorus in the ratio of 7:4:2 (Round, 1973). Apart from ammonium and nitrate ions, elemental nitrogen is also utilized by some species of cyanophyta e.g. *Phormidium*, *Oscillatoria* etc. during nitrogen fixation (Gallon, 1989). Nitrogen is considered to be the most important macronutrient in the control of growth of marine phytoplankton (Gilbert, 1988; McCarthy, 1980; Ietswaart et al., 1994). Many species of marine algae have also been shown to use amino acids and other organic nitrogen sources (Antia et al., 1991), as they possess an uptake system for amino acids through amino acid oxidases (Palenik and Morel, 1990). In this study, there was a significant difference in the uptake and utilization of nitrate and NH_4 ions between the stressed and non-stressed samples. The non-stressed samples experienced a rapid decline in the concentration of ammonia mostly after 2 weeks signifying a rapid uninhibited uptake system in contrast to the stressed samples which still had high concentrations of ammonia till 7 weeks after which decreases were noticed. This could be due to a delayed or inhibited uptake system occasioned by the presence of the hydrocarbon. There were increases in concentration of ammonia after 7 weeks for most samples probably due to the death of some microalgae releasing free ammonia with a consequent rise in pH (Figures 1 and 2).

The utilization of nitrate had a similar profile to that of ammonia probably because it is a mixed microalgal community and both macronutrients were used up simultaneously though the ammonium was preferred (De la Noue and De Pauw, 1988). There was equally a de-




Figure 1. Changes in pH levels of fertilizer factory effluent exposed to crude petroleum hydrocarbon.




Figure 2. Changes in ammoniacal nitrogen in nitrogenous fertilizer plant effluent exposed to crude petroleum hydrocarbon

layed utilization of the nitrate ions till after 7 weeks and this was probably due to the presence of the hydrocarbon (Figure 3). The rise and fall of pH in this study could be indicative of the utilization dynamics of these ions. The hydrocarbon in the various samples could have stressed the microalgae causing a nutrient-dependent lag period of microalgal growth leading to increased excretion (Letswaart et al., 1994) and interferred nutrient supply. Kinetic studies have indicated that algae generally have low specific affinities for ions (Button, 1985) which in this case could have been compounded by the presence of

the hydrocarbon. It is however well known that under nitrogen-limiting conditions, algae are capable of utilizing alternative nitrogen sources (Flynn and Byrett, 1986; North and Stephens, 1972). Recent studies have also indicated that inorganic nitrogen and phosphorus can directly limit phytoplanktonic growth. In system supplemented with NH_4NO_3 , phosphorus addition resulted in up to two to five fold increases in total phytoplankton biomass (Jianhua Le et al., 1994).

In this study, there was no significant difference between the means of most of the samples suggesting that




Figure 3. Changes in nitrate nitrogen levels of fertilizer factory effluent exposed to crude petroleum hydrocarbon.




Figure 4. Changes in phosphate levels of fertilizer factory effluent exposed to crude petroleum hydrocarbon.

the PO_4^{3-} was probably not a limiting factor in the experiment. The phosphate levels were also not constant in the two sample groups. The increases observed beyond 7 to 9 weeks (Figure 4) and could be attributed to death of certain taxa particularly the diatoms. Sulphur is present in small quantities in all microalgal cells (Round, 1973). The SO_4^{2-} concentration showed a gradual decline in both oil and non-oil samples till the last week of study when equally, there were increases possibly due to microalgal

deaths through putrefaction leading to a slightly higher mineral water (Figure 5). The diatoms particularly require the SO_4^{2-} for silica uptake which is required in soluble form for wall silification. Utilization of silica by diatoms is linked with their sulphur metabolism, so that shortages of this element could indirectly be growth limiting. The decreases in SO_4^{2-} level could therefore be linked to utilization by diatoms. The decline in the population of the diatoms probably by death through putrefaction could have led to




Figure 5. Sulphate levels in fertilizer factory effluent exposed to crude petroleum hydrocarbon.




Figure 6. Biomass (dry weight) production by microalgae in fertilizer factory effluent exposed to crude petroleum hydrocarbon.

the release of the SO_4^{2-} into the medium.

This succession pattern and nutrient uptake were almost similar to that reported by Soler et al. (1991). The biomass equally was in line with the work of Jianhua Le et al. (1994) where two to five fold increases in total phytoplankton biomass were reported in response to N and P fertilizations (Figure 6). However, since chlorophyll a content is affected by the different abilities of the organisms to obtain PO_4^{3-} (Jianhua Le et al., 1994), the non-stressed samples showed greater increases in chlorophyll a content at each given time in the study (Figure 7). Stressed phytoplankton communities also increased in abundance and biomass. This suggested obvious recovery of the systems. The changes in conductivity of fertilizer factory effluent exposed to crude petro-

leum hydrocarbon is presented in Figure 8.

In this study overall, there was a reduction in nutrients; NH_4^+ , NO_3^- , PO_4^{3-} and SO_4^{2-} between 2 days to 7 weeks for the non-stressed samples and between 5 days to 9 weeks for the stressed samples, leading to microalgae proliferation. However, due to the presence of the hydrocarbon (which showed no significant difference between samples with the different crude concentrations), there was delayed nutrient uptake. This is in line with the findings of Baker (1970) and Amakiri and Onofeghara (1983) for higher plants. The consequence of this response is the observed reductions in chlorophyll content, biomass etc. There seemed however to be a quick recovery in the oiled systems as in the case of the Oshika oil spill where there was reported cases of blue-green




Figure 7. Chlorophyll-a production (mg/l) by microalgae in fertilizer factory effluent exposed to crude petroleum hydrocarbon.




Figure 8. Changes in conductivity of fertilizer factory effluent exposed to crude petroleum hydrocarbon.

algae bloom a few weeks after the spill (IPS, 1986).

Biomass measurements

Dried biomass measurements of microalgae cultures indicated increases in dry weight and chlorophyll a content particularly after one week when full bloom was attained in the samples without crude additions. Significant increases in biomass and chlorophyll a content was observed in the samples with crude additions only after 4 weeks (Figure 7). The original NAFCON effluent/sediment culture on the day of collection had high num-

bers and diverse species of microalgae probably because of the high nutrient concentrations of the effluent that is discharged into the river.

Taxonomic studies showed that three main groups were present during the course of the experiments. The microalgae profile included genera of cyanobacteria (blue-green algae), Bacillariophyceae (the diatoms) and the chlorophyceae (green algae). In the raw effluent, the diatoms had the least percentage distribution of 29.08% followed by the green algae with 35.29%. The blue-green algae had the highest percentage distribution of 35.62%. (Table 2) The Bacillariophyceae include species of *Ampiprora*, *Cyclotella*, *Navicula pinnularia* and *Nitzschia*.

Table 2. Microalgal diversity, species characterization and percentage occurrence.

Organism	Untreated sample	A	B	C	D	AA	BB	CC	DD
Diatoms									
<i>Amphiprora sp.</i>	20	48	69	21	11	9	18	71	14
<i>Cyclotella striata</i>	33	22	23	28	-	2	35	36	28
<i>Coscinodiscus radiata</i>	41	13	18	55	24	-	28	22	33
<i>Dictylum brighwelli</i>	65	39	27	34	17	8	12	14	21
<i>Diatoma liemale</i>	36	10	94	-	28	10	16	21	26
<i>Melosira granulata</i>	9	66	15	46	11	14	-	10	20
<i>Navicula cuspidata</i>	59	37	7	109	61	-	65	39	19
<i>Navicula minima</i>	80	24	3	42	5	54	22	17	20
<i>Pinnularia gibba</i>	3	-	3	28	19	18	4	11	24
<i>Nitzschia closterium</i>	7	10	2	39	36	55	9	9	10
<i>Naviculabacillum</i>	3	10	-	65	81	28	11	-	3
Total	356	370	261	467	293	198	220	250	218
%	29.08	39.15	32.22	41.92	34.03	38.82	35.43	34.29	46.09
Blue-green algae									
<i>Anacystis Geroginosa</i>	68	55	-	-	52	11	10	-	4
<i>Anabaena flos-aguae</i>	45	36	25	94	35	19	-	36	13
<i>Oscillatoria Sancta</i>	20	-	39	58	-	28	27	20	1
<i>Oscillatoria Chadybaca</i>	37	57	41	24	-	9	15	6	16
Phormidium Uncinatum	42	21	11	-	21	14	20	3	5
<i>Pleurosyna donyatum</i>	20	13	28	33	28	6	5	12	6
<i>Oscillatoria sp.</i>	204	26	26	64	27	6	9	19	10
Total	436	208	170	273	163	93	86	96	55
%	35.62	22.01	20.99	24.51	18.93	18.24	13.85	13.17	11.63
Green algae									
<i>Schiothrix Pulvinata</i>	261	154	96	112	143	112	106	149	130
<i>Scenedesmus</i>	44	135	140	195	140	67	160	97	37
<i>Quadricauda</i>									
<i>Chlorella Sp.</i>	127	78	143	67	122	40	49	137	33
Total	432	367	379	374	405	219	315	383	200
%	35.29	38.84	46.79	33.57	47.04	42.94	50.72	52.54	42.28
Overall Total	1224	945	810	1114	861	510	621	729	473

The cyanobacteria included species of *Anacystis*, *Anabaena*, *Oscillatoria phormidium* and *pleurosyna*, while the chlorophytes were made up of three genera *Schizothrix*, *Scenedesmus* and *Chlorella*. From the 8th day of study, when samples with no crude additions attained full bloom, the chlorophytes dominated the other groups till the end of the study attaining percentages that ranged from 33.57% on day 8 to 89.99% after 9 weeks. In this, the *Chlorella* followed by *Scenedesmus quadricauda* were particularly dominant from week 1 up till week 5 when the species *Schizothrix pulinata* appreciated in number and from 6 to 9 weeks, all three species dominated in the culture. The chlorophytes in the samples with hydrocarbon were dominant at day zero till day 8 when the cyanobacteria gained control of the culture. By week 5, the chlorophytes once again became the dominant species till end followed by the cyanobacteria.

The NAFCON industrial waste water or effluent was

found from this study suitable to support the blooming of microalgae including *Chlorella*, *Scenedesmus* etc. as was previously done by Anaga and Abu (1996), Abu and Epegu (1992) suggesting use for the waste in microalgae biomass production and effluent biotreatment. This is in consonance with the work of Chapman and Gellenbeck (1989) which showed that some species of microalgae such as *Porphyridium*, *Scenedesmus*, *Chlorella* and *Spirulina* could actually be cultivated in the laboratory.

The biomass measurements as illustrated by the dry weight and chlorophyll contents and also the number of cells indicated that there were increases in biomass indicating growth. However, this was much more in the non-stressed than in the stressed samples where a much longer lag phase of growth was noticed. In the non-stressed samples, nutrient uptake was much more rapid and immediate. Cross-system analysis of results however, showed that phytoplankton chlorophyll-a, dry weight

and total number of cells were non-significant in all systems. In microalgal diversity, microalgal populations showed 3 main taxa with the cyanobacteria dominating at the commencement of study. The profile later showed the chlorophyta particularly, *Chlorella* sp. and *Scenedesmus quadricauda* dominating till end of the study during which time the diatoms had almost become extinct (Table 2).

REFERENCES

- Abu GO, Epegu CD (1992). Polymer Production by microalgae isolated from aqua culture facilities at Aluu in Nigeria. Abstract Ninth International Biotechnology Symp. (Subsection: Biotechnology in Developing Countries), Nov. crystal city, VA USA. 866: 16-21
- Alexander M (1977). Introduction to soil microbiology (2nd ed.) John Wiley and Sons, Inc., p. 267.
- Alexander SJ, Schwarz JR (1980). Short-Term effects of south Louisiana and Kuwait crude oils on glucose utilization by marine bacterial populations – Appl. Environ. Microbiol. 7: 341-345.
- Amakiri JO, Onofeghara FA (1983). Effect of crude oil pollution on the growth of *zea mays*, *Abelmoschus esculentus* and *capsicum frutescens*. Oil Petrochem. Pollut. 1: 199-205.
- American Public Health Association, APHA (1985), 13th August, 1993. APHA. (1985). Standard Methods for the examination of Water and Waste Water, 6th Edition. American Public Health Association, Washington, D.C.
- Anaga A, Abu GO (1996). A laboratory scale cultivation of *chlorella* Sp. and *spirulina* sp. using waste effluent from a fertilizer company in Nigeria. Bioresour. Technol. 58: 93-95.
- Antia NJ, Harrison PJ, Oliviera L (1991). The role of dissolved organic nitrogen in phytoplankton nutrition, cell biology and ecology. Phycologia 30: 1-89.
- Atlas RM, Bartha R (1973). Abundance, distribution and oil biodegradation potential of microorganisms in Raritan Bay. Environ. Pollut. 4: 291-300.
- Azov Y, Shelef G (1982). Operation of high-rate oxidation ponds; theory and experiments. Water Res. 16: 1153-1160.
- Baker JM (1970). Effects of oil on plants. Environ. Pollut. 1: 27-44.
- Blumer M, Sanders HL, Grassle JF, Hampson GR (1971). A small oil spill. Environment 13(2): 2-12.
- Button DK (1985). Kinetics of nutrient limited transport and microbial growth. Microbiol. Rev., 49: 270-297.
- Chapman DJ, Gellenbeck KW (1989). An historical perspective of algae biotechnology. In Cresswell RC, Rees TA, Shab N (eds.) Algae and cyanobacterial Biotechnology; John Wiley and sons.
- Colwell, RR, Walker JD (1977). Ecological aspects of microbial degradation of petroleum in the marine environment. CRC Crit. Rev. Microbiol. 5: 423-445.
- Cornwell DA, Zoitek J, Patrineiy CD, Furman TD, Kim JI (1977). Nutrient removal of water hyacinths. J. Water Pollut. Contr. Fed. 49: 1.
- De La Noue J, De Pauw N (1988). The potential of microalgae biotechnology. A review of production and uses of microalgae Biotechnol. Adv. 6: 725-770.
- Fallowfield HJ, Garret MK (1985). The treatment of wastes by algae culture. J. Appl. Bact. Symp. Suppl., pp. 187-205.
- Flynn KJ, Byrett PJ (1986). Utilization of L-lysine and L-arginine by the diatom *phaeodactylum tricornutum*. Mar. Biol. (Berlin) 90: 157-163.
- Gallon JR (1989). The industrial potential of cyanobacteria multipurpose organism. In: Greenshields Rod (ed), Resources and Application of Biotech. The New Wave, pp. 13-26.
- Gibson DT, Mahadevan V, Jerina DM, Hagi H, Yeh HJC (1975). Oxidation of the carcinogens benzo (α) pyrene and benzo (α) anthracene to dihydrodiolol by a bacterium, Science, 189: 295.
- Gilbert PM (1988). Primary productivity and pelagic nitrogen Cycling, In Blackburn TH, Sorensen J (eds) Nitrogen cycling in coastal marine environments. Wiley, New York, pp. 3-31.
- Gupta SK, Sharma R (1996). Biological oxidation of high strength Nitrogenous waste water. Water Res. 30(3): 593-600.
- Hollaway SI, Faw GM, Sizemore RK (1980). The bacterial community composition of an active oil field in the Northwestern gulf of Mexico. Mar. Pollut. Bull. 2: 193-196.
- Ietswaart T, Schweider PJ, Prins RA (1994). Utilization of organic Nitrogen source by two phytoplankton species and a bacterial isolate in pure and mixed culture. Appl. Environ. Microbiol., pp. 1554-1560.
- Institute of Pollution Studies (1986). "Oshika Oil Spill (NAOC pipeline) Environmental Impact Assessment. Final Report". Submitted to the petroleum Inspectorate, NNPC by Institute of Pollution studies, Rivers State University of Science and Technology, Port Harcourt. RSUST/IPS/R/86/022xx + p. 181.
- Jianhua Lee J, Wehr D, Campbell L (1994). Uncoupling of Bacterioplankton and phytoplankton production in fresh water is affected by Inorganic Nutrient Limitation. Appl. Environ. Microbiol, pp. 2086-2093.
- Jobson AM, Mclawghlin M, Cook FD, Westlake DW (1974). Effect of amendments on the microbial utilization of oil applied to soil. App. Microbiol 27: 166-171.
- McCarthy JJ (1980). Nitrogen and phytoplankton ecology In I. Morris (ed.) The physiological ecology of phytoplankton. Blackwell scientific publications, Oxford, pp. 191-233.
- North BB, Stephens GC (1972). Amino acid transport in *Nitzschia ovalis* Arnott. J. Phycol. 8: 54-68.
- Palenik BF, Morel MM (1990). Comparison of cell-surface L. amino acid oxidases from several marine phytoplankton. Mar. Ecol. Prog. Ser. 59: 195-201.
- Round FE (1973). The Biology of the Algae. Edward Arnold Publishers. London.
- Schwartz JR, Walker JD, Colwell RR (1974). Growth of deep sea bacteria on hydrocarbon at ambient and *in situ* pressure, J. Appl. Microbiol. 28: 982-986.
- Shelef G, Azov Y, Moraine R, Oron G (1980). Algae mass production as an integral part of a wastewater treatment and reclamation system. In Shelef G, Soeder CJ (eds). Algal Biomass Production and use, Elsevier/North Holland Biomedical press pp. 183-189.
- Shelef G, Soeder CJ (eds) (1980). Algae biomass production and use. Elsevier /North Holland Biomedical press.
- Soler J, Saez M, Llorens T, Martinez FT, Berna LN (1991). Changes in physicochemical parameters and photosynthetic microorganisms in a deep waste water self-depuration lagoon. Water Res. 25: 689-695.
- Westlake DWS, Jobson AM, Cook FD (1978). In situ degradation of oil in a soil of the aboreal region of the North West territories can. J. Microbiol. 24: 254-260.