

Full Length Research Paper

The antibacterial potentials of *Nauclea latifolia*

Okwori, A. E. J.^{1*}, Okeke, C. I.¹, Uzoehina, A.¹, Etukudoh, N. S.², Amali, M. N.³, Adetunji, J. A.⁴
and Olabode, A. O.¹

¹Department of Medical Microbiology, Federal College of Veterinary and Medical Laboratory Technology, Vom, Nigeria.

²Federal School of Medical Laboratory Technology, (JUTH), Jos, Nigeria.

³Department of Physiology, College of Health Sciences, Benue State University, Makurdi, Nigeria.

⁴Department of Chemical Pathology, Jos University Teaching Hospital, Jos, Nigeria.

Accepted 12 March, 2008

Herbal medicine is readily available in diverse African vegetation with the potentials of introducing new templates into medicine worldwide. Evaluating plants from the traditional African system of medicine provides us with clues on how these plants can be used in the treatment of diseases. *In vitro* effect of *Nauclea latifolia* extract in hot water, cold water, petroleum ether and chloroform at concentrations of 200, 150, 100, 50% were tested on some pathogenic bacteria such as *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi* and *Pseudomonas aeruginosa*. Using agar diffusion punch hole method, both the aqueous and alcoholic extracts of the leaves and roots showed appreciable inhibitory effect when compared to the positive control on *S. aureus* and *P. aeruginosa* while *S. typhi* and *E. coli* were resistant to the extracts. Using serial doubling dilution, the minimum inhibitory concentration (MIC) was determined. The minimum bacteria concentration was determined by plating various dilutions of the extracts without turbidity. Aqueous and alcoholic extracts of *N. latifolia* showed inhibitory and bactericidal activity on the test organisms. The alcoholic extracts showed larger zone of inhibition on the test organisms. The alcohol leaf extracts showed a higher percentage of growth inhibition when compared to the positive control. The MIC ranges from 6.25 – 150 mg/ml on *S. aureus* and 12.5 – 150 mg/ml for *P. aeruginosa*. The MBC ranges from 100 – 150 mg/ml. The phytochemical analysis revealed the presence of saponin, resins, alkaloids, and carbohydrate.

Key words: Antimicrobial potentials, *Nauclea latifolia*, bacterial pathogens, Nigeria.

INTRODUCTION

Bacterial and fungal pathogens have evolved numerous defense mechanisms against antimicrobial agents and resistance to old and newly produced drugs are on the rise. Traditional medicine is a comprehensive term used to refer both to traditional medicine systems such as traditional Chinese medicine, Indian ayurveda and Arabic unani medicine, and to various forms of indigenous medicine (WHO, 1978). Historically, plant has provided a good source of anti-infective agents; emetine, quinine and berberine remain highly effective instruments in the fight against microbial infections. Phytomedicines derived from plants have shown great promise in the treatment of intractable infectious diseases including opportunistic

AIDS infections. Plants containing protoberberines and related alkaloids, picralima-type indole alkaloids and *Garcinia biflaronones* used in traditional African system of medicine, have been found to be active against a wide variety of microorganisms.

Seeking remedies for human ailments from the environment has formed the basis for therapeutics (Potier et al., 1990). African plants constitute a rich untapped pool of natural products (Bringmann and Pokorny, 1995). Extracts of different parts of the plant *Nauclea latifolia* (e.g. fruits, leaves, stem-bark and roots) in hot water or alcohol are used in form of infusions, decoctions or concoctions (Irvine, 1961; Agoha, 1974).

Scientific investigations of medicinal plants have been initiated in many countries because of their contributions to health care. The continual search for, and the interest in natural plant products for use as medicines has acted as the catalyst for exploring methodologies involved in

*Corresponding author. E-mail: Okwori2001@yahoo.com. Tel: 234 80 37 00 11 72. Fax: 234 73 280271.

obtaining the required plant materials and probing their constituents. The plant, *N. latifolia*, is used in many African countries by traditional medical practitioners for the treatment of various ailments including bacterial diseases. It is most effective against *Corynebacterium diphtheriae*, *Streptobacillus spp*, *Streptococcus spp*, *Neisseria spp*, *Pseudomonas aeruginosa*, *Salmonella spp* (Deeni and Hassan, 1991).

N. latifolia commonly known as pin cushion tree is a straggling shrub or small tree native to tropical Africa and Asia. It is a tropical plant that grows commonly in Akwa Ibom and Cross River State of Nigeria and also in most parts of the northern Nigeria. In Akwa Ibom and Cross River states, it is called "Mbom-mbog" whilst in the northern Nigeria it is called "Tabasiya". It is known as "Uche" by Igede people. It is found in the forest and fringe tropical forest. The plant is also used in the treatment of ailments like malaria (Kokwaro, 1976; Akabue and Mittal, 1982; Boye, 1990), gastrointestinal tract disorders (Maduabunyi, 1995), sleeping sickness (Kerharo, 1974), prolong menstrual flow (Elujoba, 1995), hypertension (Akabue and Mittal, 1982) and as a chewing stick (Asubiojo et al., 1982). Aqueous extracts from *N. latifolia* S.M. (*Rubiaceae*) is also a common plant used in Ivory Coast by traditional healers for the treatment of malaria. In Burkina Faso, the plant is used for the treatment of several diseases as jaundice, malaria, infant gastroenteritis, dysentery etc (Fernandez de la pradisilla, 1982a,b). The plant is also used as a tonic and fever medicine, chewing stick, toothaches, dental cares, septic mouth and malaria, diarrhea and dysentery (Lamidi et al., 1995).

Many commercially proven drugs used in modern medicine were initially used in crude form in traditional or folk healing practices, or for other purposes that suggested potentially useful biological activity. The primary benefits of using plant-derived medicines are that they are relatively safer than synthetic alternatives, offering profound therapeutic benefits and more affordable treatment. This study was aimed at establishing the antimicrobial potentials of widely acclaimed medicinal value of *N. latifolia* extracts on bacterial pathogens isolated in Vom.

MATERIALS AND METHODS

The plants were obtained within the premises of National Veterinary Research Institute, Vom. The botanical identity was determined and authenticated at the Department of Botany, University of Jos, Nigeria where a voucher specimen has been deposited.

Processing of the plant

The leaves and roots of the plant were washed with tap water and rinsed in distilled water. They were dried using hot air oven at 40°C for three days. The dried plant parts were pulverized using sterile laboratory mortar to obtain the powdered form. These were stored in airtight sterile bottles until required for analysis.

Crude extraction

The method of Junaid et al. (2006) and Okwori et al. (2007) were adopted for this study.

Preparation of the extract

The aqueous extracts were reconstituted using sterile distilled water while chloroform and petroleum ether were reconstituted using 50% acetone to obtain concentrations of 200, 150, 100 and 50 mg/ml.

Sources of bacterial isolates

The stock bacterial isolates used were obtained from the Bacteriology laboratory of the Federal College of Veterinary and Medical laboratory Technology, Vom and the Bacterial research Laboratory, National Veterinary research Institute, Vom. Purity plate for each of the bacterial isolates was used for this study.

Preparation of the test bacteria

Fresh purity plates of the test bacteria were made from the isolate cultures obtained on agar slants. The isolates were sub-cultured on selective and differential solid media and re identified phenotypically (Okwori et al., 2007).

With the aid of a sterile wire loop, colonies of fresh cultures of the different bacterial isolates were picked and suspended on 5 ml nutrient broth in different well labeled sterile 10 ml bijou bottles. 1 ml of the isolate from the broth was diluted in peptone water using the two-fold dilution. For *Staphylococcus aureus* which is a slow growing organisms, 0.02 ml of (10^{-1} to 10^{-7}) dilutions was picked using a single channel pipette and plated out in triplicate on nutrient agar to obtain the mean viable count of the isolates. The same procedure was done for *Salmonella typhi*, *Escherichia coli* and *P. aeruginosa*, but since they are fast growing organisms, 0.02 ml of (10^{-4} to 10^{-10}) dilutions were plated out on nutrient agar and incubated at 37°C for 24 h (Miles and Misra, 1938).

After 24 h, different numbers of colonies were obtained at different dilutions for the different bacterial isolates. The mean numbers of colonies were calculated to obtain the colony-forming unit per ml (cfu/ml) for the different isolates. 1 ml of the standard inoculum of the different bacterial isolates was used in flooding sterile nutrient agar plates in the agar diffusion method of *in vitro* antimicrobial sensitivity test.

Antimicrobial potential

A standardized inoculum of $1 - 2 \times 10^7$ cfu/ml was used for the inoculation of plates. The plates were allowed to dry in the incubator for 30 min at 37°C and with the aid of a sterile standard cork borer, 6 wells were bored at equidistant. The bottoms of the wells were sealed with sterile molten nutrient agar (Biotec, Suffolk, UK) to prevent seepage of the extract under the agar. The 5 and 6th wells served as positive and negative controls. The sterile distilled water served as negative control. Ciprofloxacin (10 ug/ml) was used as the positive control. 0.2 ml of each prepared concentration of the extracts was aseptically introduced into wells 1 – 4. The plates were allowed on the bench for 40 min for pre-diffusion followed by an overnight incubation at 37°C. The resulting zones of inhibition were measured using a ruler calibrated in millimeters. The average of the three readings was taken to be zone of inhibition of the bacterial isolates in question at that particular concentration (Abayomi, 1982; Junaid et al., 2006).

Table 1. Antimicrobial activity of different solvent extracts (zone of inhibition measured in mm).

Isolate	Conc.	HWR	CWR	DR	CR	HWL	CWL	CL	PDL
<i>S. aureus</i>	200	16	16	20	20	18	17	20	18
	150	15	15	18	19	17	15	15	15
	100	13	12	17	17	15	12	15	15
	50	10	10	13	13	12	10	10	12
<i>S. typhi</i>	200	0	0	0	0	0	0	0	0
	150	0	0	0	0	0	0	0	0
	100	0	0	0	0	0	0	0	0
	50	0	0	0	0	0	0	0	0
<i>P. aeruginosa</i>	200	15	12	20	15	23	23	22	24
	150	14	11	20	15	22	22	20	22
	100	10	11	15	13	23	22	18	20
	50	0	0	10	10	20	20	26	26

HWR – Hot water root; CWR – Cold water root; DR – Diether root; CR – Chloroform root; HWL – Hot water leaves; CWL – Cold water leaves; PDL – Petroleum ether leaves; CL – Chloroform leaves.

Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

MIC of the extracts was determined by dilution to various concentrations according to the macro broth dilution technique (Baron and Finegold, 1990; Akinyemi et al., 2005). Standardized inoculum was added to series of sterile tubes of nutrient broth containing two fold dilution of the extracts and incubated at 37°C for 24 h. The MIC was read as the least concentration that inhibited the growth of the test organisms. However, the MBC was determined by sub-culturing the test dilutions onto fresh drug-free solid medium and incubated further for 18 – 24 h. The highest dilution that yielded no single bacterial colony on a solid medium was taken as MBC (Okwori et al., 2007).

Phytochemical screening

The presence of alkaloids, resins, saponins, glycosides, tannins, flavonoids, cardiac glycoside, steroidal ring, steroidal terpenes, anthraquinone and carbohydrates were determined as described by Akinyemi et al. (2005) and Junaid et al. (2006) and Okwori et al. (2007).

RESULTS

Alcoholic and aqueous extracts of the roots and leaves extracts of *N. latifolia* were tested on four known clinical bacterial isolates using agar diffusion (punch hole method; see Figure 1). Alcoholic root extracts at concentration ranging from (200 - 50%) produced mean inhibitory zone ranging from 13 – 20 mm on *S. aureus* while alcoholic leaf extract produced mean inhibitory zone ranging from 10 – 20 mm on *S. aureus*. Alcoholic root extract produced mean inhibitory zone ranging from 10 – 20 mm on *P. aeruginosa* while alcoholic leaf extract produced mean inhibitory zone ranging from 16 – 24 mm on *P. aeruginosa* which was significant compared with the control.

Aqueous root extract at concentration ranging from




Figure 1. Inhibitory effect of the extracts at different concentrations on sensitivity test agar.

(200 - 50%) produced mean inhibitory zone ranging from 10 – 16 mm on *S. aureus* while aqueous leaf extract produced a mean zone of inhibition ranging from 10 – 18 mm on *S. aureus*. Aqueous leaf extract produced a significant zone of inhibition ranging from 20 – 23 mm on *P. aeruginosa* when compared with the control while the aqueous root extract produced zone of inhibition ranging from 10 – 15 mm on *P. aeruginosa* (Table 1).

S. typhi and *E. coli* were resistant on both alcohol and aqueous leaves and roots extract at all concentration. The alcoholic and aqueous extracts exhibited both bacteriocidal and bacteriostatic activities on the positive

Table 2. MIC based on bacterial turbidity at various conc. (mg/ml) using two-fold dilution.

Isolate	Extract	200	150	100	50	25	12.5	6.25	3.12	1.56	0.78	0.39	0.19
<i>S. aureus</i>	HWL	-	-	-	-	-	-	+	+	++	++	++	++
	CWL	-	-	-	-	-	-	-	+	++	++	++	++
	DL	-	-	-	-	-	-	-	+	++	++	++	++
	CL	-	-	-	-	-	-	+	+	++	++	++	++
	HWR	-	-	-	-	-	-	+	+	++	++	++	++
	CWR	-	-	-	-	-	-	+	+	++	++	++	++
	CR	-	-	-	-	-	-	-	+	++	++	++	++
	DR	-	-	-	-	-	-	-	+	++	++	++	++
<i>P. aeruginosa</i>	HWL	-	-	-	-	-	-	++	++	++	++	++	++
	CWL	-	-	-	-	-	-	++	++	++	++	++	++
	DL	-	-	-	-	-	-	++	++	++	++	++	++
	CL	-	-	-	-	-	-	++	++	++	++	++	++
	HWR	-	-	-	-	-	-	++	++	++	++	++	++
	CWR	-	-	-	-	-	-	++	++	++	++	++	++
	CR	-	-	-	-	-	-	++	++	++	++	++	++
	DR	-	-	-	-	-	-	++	++	++	++	++	++

- = Not turbid; + = slightly turbid; ++ = very turbid.

Table 3. MIC based on bacterial growth.

Isolate	Extract	Concentration					MBC
		200	150	100	50	25	
<i>S. aureus</i>	HWL	-	-	+	+	+	150
	CWL	-	-	-	+	+	100
	DL	-	-	-	+	+	100
	CL	-	-	-	+	+	100
	HWR	-	-	+	+	+	150
	CWR	-	-	+	+	+	150
	CR	-	-	+	+	+	150
	DR	-	-	+	+	+	150
<i>P. aeruginosa</i>	HWL	-	-	-	+	+	100
	CWL	-	-	-	+	+	100
	DL	-	-	+	+	+	150
	CL	-	-	+	+	+	150
	HHR	-	-	+	+	+	150
	CHR	-	-	+	+	+	150
	CR	-	-	+	+	+	150
	DR	-	-	+	+	+	150

- = No bacterial growth; + = bacterial growth.

test organisms (Tables 2 and 3).

The phytochemical analysis of both root and leaf extracts revealed the presence of carbohydrate, saponins, resins, alkaloids.

DISCUSSION

From the results obtained, both the alcohol and aqueous

extracts of leaves and roots of *N. latifolia* were active against 2 out of the 4 test bacterial organisms. The exception to the rule was *E. coli* and *S. typhi*, which was found to be resistant to both, though a gram negative bacterium, has been reported to be sensitive to extracts of *N. latifolia* (Omer et al., 1998) but this was not in agreement with this work. This may be as a result of natural resistance, genetic variability or mutational changes which occur even before the introduction of

drug. The resistance can be transferred from one bacterium to another of the same species, and also between different species and sometimes, even between related general. Deeni and Hussain (1991) documented the susceptibility of *P. aeruginosa* and *Salmonella* species to plant extracts. However, our finding is in agreement with this report except for *S. typhi* and this could be as a result of the species of *Salmonella* used in the previous work.

Both the aqueous and alcohol leaf extracts showed a wide range of antibacterial activity when compared to the positive control. Umeh et al. (2005) suggests in his work that the antibacterial constituents of the plant are preferentially concentrated in the leaves and in this work the leaf extracts showed a higher percentage of growth inhibition.

Both aqueous and alcohol extracts of the root showed zones of inhibition. This work is also in agreement with similar reports documented by Iwu (1993) whose studies showed that the root has antibacterial activity against gram positive and negative bacteria. It was found in this work that test bacterial organisms were more susceptible to alcoholic extracts than the aqueous extracts. This may be due to its phytochemical extraction capability (Junaid et al., 2006). Similarly, the studies of Umeh et al. (2005) showed that petroleum ether and chloroform extracts of the plants studied have more inhibitory effect than methanolic and aqueous extracts, although the inhibitory effects of aqueous and methanolic extracts of medicinal plants are usually reported (Ogunlana and Ramstad, 1975; Omer et al., 1998; Olayinka et al., 1992). This report is also in line with the report of Obi and Onuoha (2000), who reported alcohol to be the best solvent for the extraction of most plant active principles of medical importance.

It was also observed from this work that the higher the concentration, the more their activity and as the concentration decreases the lower the antibacterial activity. Hence an acceptable and effective dosage can be prepared by traditional healers for the control and eradication of bacterial pathogens.

The variation in results imply that the MBC results obtained after plating various dilutions of extracts is more reliable compared to MIC results obtained usually using turbidity as an index (Junaid et al., 2006).

One of the greatest service which can be rendered to any country is to add a useful plant to its culture. It is therefore highly essential that medicinal plants whose properties have not been fully characterized should form a top agenda of top management in developing nations whose citizens are sometimes unable to afford expensive orthodox medicine. This policy if pursued will not only preserve the scarce foreign exchange but also promote the spirit of plant conservation.

ACKNOWLEDGEMENT

The authors thank the Provost, Federal College of Veteri-

nary and Medical Laboratory Technology (FCVMLT), National Veterinary Research Institute, Vom, Nigeria for permission to publish this paper.

REFERENCES

- Abayomi S (1982). The State of Medicinal Plants Research in Nigeria. University of Ife Press, p. 200.
- Agoha RC (1974). Medicinal plants in Nigeria. Faculterider wiskurde en Natuurwesteris Chappen Nijeggan, The Netherlands.
- Akabue P, Mittal GC (1982). Clinical Evaluation of a Traditional Herbal Practice in Nigeria: a Preliminary Rep. J. Ethnopharmacol 6(3): 355-359.
- Akinyemi KO, Oladapo O, Okwara CE, Ibe CC, Fasura KA (2005). Screening of Crude Extracts of Six Medicinal Plants Used in South – West Nigeria unorthodox medicine anti- methicillin resistant *Staphylococcus aureus* activity. BMC Complement. Alternat. Med. 5(6): 1-7.
- Asubiojo OI, Guinn VP, Okunuga A (1982). Multi-element Analysis of Nigerian chewing sticks by Instrumental Neuron Activation Analysis. J. Radio Anal. Chem. 74: 149-156.
- Baron JE, Fingold SM (1990). Methods for Testing Antimicrobial Effectiveness. In: Bailey Scotts Diagnostic Microbiology, Mosby CV (ed), Missouri, pp. 171-194.
- Boye GL (1990). Studies on Antimalarial Action of *Cryptolepis sanguinolenta* Extract. Proc. Int. sym. On East – West Med. Seoul, Korea: 243-251.
- Bringmann G, Pokorny F (1995). The Naphthyliso quinoline alkaloids in: The Alkaloids. Vol. 46: Cordell G (Ed) Academic University press, New York, p. 127.
- Deeni Y, Hussain H (1991). Screening for antimicrobial activity and for alkaloids of *Nauclea latifolia*. J. Ethnopharmacol. 35: 91-96.
- Elujoba AAA (1995). Female infertility in the Hands of Traditional Birth Attendants in South – West Nigeria. Fitoterapia 66(3): 239-248.
- Fernandez de la pradilla C (1982a). Plantes medicinales contre les hepatitis: 51 especes traopicales. Polygr., Ed. Librairie Jeunesse d' Afrique, Ouagadougou, 61 pges.
- Fernandez de la pradilla C (1982b). Plantes medicinales vendue sur les marchès de Ouagadougou. Polygr., Ed. Librairie Jeunesse d' Afrique, Ouagadougou, p. 22.
- Irvine F (1961). Woody plants of Ghana. Oxford Univ. Press, London, p. 969.
- Iwu M (1993). Handbook of African Medicinal plants. CRC Press, Boca Raton, FL.
- Junaid SA, Olabode AO, Onwuliri FC, Okwori AEJ, Agina SE (2006). The antimicrobial properties of *Ocimum gratissimum* extracts on some selected bacterial gastrointestinal isolates. Afr. J. Biotechnol. 5(22): 2315-2321.
- Kerharo J (1974). Historic and Ethnopharmacognosic Review on the Belief and Traditional Practices in the Treatment of sleeping sickness in West Africa. Bull. Soc. Med. Afr. Noire Lang FR. 19: 400.
- Kokwaro JO (1976). Medicinal plants of East Africa. East African Literature Bureau, Nairobi.
- Lamidi ME, Olivier R, Faure L, Debrauwer L, Nze E, Balandsard G (1995). Quinovic acid glycosides from *Nauclea diderichic*. Planta Med. 61: 280-281.
- Maduabunyi II (1995). Anti – hepatocidal and Trypanocidal activities of the Ethanolic Extract of *Nauclea latifolia* Root Bark. J. Herbs Spices Med. Plants 3(2): 23-53.
- Miles AA, Misra SS (1938). The estimation of the bactericidal power of the blood. J. Hyg. 38: 732-749.
- Obi VI, Onuoha C (2000). Extraction and characterization methods of plants and products. In: Biological and agriscultural techniques. Ogbulie JN, Ojiako OJ (eds). Websmedia publications, Owerri. pp. 271-286.
- Ogunlana EO, Ramstad E (1975). Investigations Into. The antibacterial activities of local plants. Planta Med. 27: 354.
- Okwori AEJ, Dina CO, Junaid SA, Okeke IO, Adetunji JA, Olabode AO (2007). Antimicrobial activities of *Ageratum conyzoides* on some selected bacterial pathogens. Internet J. Microbiol. Vol. 4 No.1.

- Olayinka DO, Onoruwe O, Lot TY (1992). Cardiovascular effects in rodents of methanolic extracts of *Khaya Senegalensis*. *Phytother. Res.* 6: 282-284.
- Omer MEA, Al Magboul AZ, EL Egani AA (1998). Sudanese plants used in folkloric medicine: Screening for antimicrobial activity. *Fitoterapia* 69: 542-545.
- Potier P, Mangatal L, Le Goff MT, Gueritte VF, Guenard O, Lavelle F (1990). Relationships between the structure of taxol analogues and their antibiotic activity. *J. Med. Chem.* 34: 992-998.
- Umeh EU, Oluma HOA, Igoli JO (2005). Antibacterial screening of four local plants using an indicator – based microdilution technique. *Afr. J. Tradit.* 2(3): 238-243.
- World Health Organization (WHO) (1978). The Promotion and Development of Traditional Medicine. Technical Report Series, p. 622.