

*Full Length Research Paper*

# Height suppression of tomato plug seedlings by an environment friendly seed treatment of plant growth retardants

Woo Gun Shin<sup>1,2,3</sup>, Seung Jae Hwang<sup>1,2</sup>, Iyyakkannu Sivanesan<sup>1</sup> and Byoung Ryong Jeong<sup>1,2\*</sup>

<sup>1</sup>Department of Horticulture, Division of Applied Life Science (BK21 Program), Graduate School, Gyeongsang National University, Jinju 660 - 701, Korea

<sup>2</sup>Research Institute Agriculture and Life Science, Gyeongsang National University, Jinju 660-701, South Korea.

<sup>3</sup>Korea Seed and Variety Service, Ministry for Food, Agriculture, Forestry and Fisheries, Milyang, Korea.

Accepted 27 July, 2009

Experiments were conducted to investigate appropriate concentrations of plant growth retardants (PGRs) and duration of seed soaking in order to suppress hypocotyl length and plug seedling height of 2 tomato cultivars (*Lycopersicon esculentum* Mill. cv. Seogeon and Seokwang). Daminozide (B-9), uniconazole (Sumagic) and ethephon (Florel) were used as PGRs. Seeds were first soaked in 15 ml PGR solutions in an environment controlled chamber (25°C) for 1, 3, or 5 days. Then seeds were washed in tap water and were dried in a 5°C chamber for 1 day. Finally, dried seeds were used in both a germination test in a chamber and a growing test in a greenhouse. Differences among cultivars in response were observed in germination and seedling growth. Although germination in petri dishes and seedling emergence in plug trays declined, suppression of hypocotyl length and seedling height was evident in uniconazole treatments in both cultivars, duration of seed soaking in PGRs solutions had greater influence than the concentration. Differences in percent germination and percent emergence were also observed as affected by PGRs. Uniconazole treatments suppressed hypocotyl length and seedling height very significantly. Seeds soaked in PGRs solutions germinated in petri dishes, but tended not to emerge well in plug trays.

**Key words:** Daminozide (B-9), emergence, ethephon (florel), germination, *Lycopersicon esculentum*, uniconazole (sumagic).

## INTRODUCTION

Because of the merits of reduction of labor in raising transplants mass production of uniform transplants and specialization of transplant production, plug transplants have been used by many growers. From the early 90 s, specialized nurseries in Korea have produced an increasing proportion of plug seedlings of vegetables and flowers for the horticulture industry due to their efficiency and economies of scale (Choi et al., 1997). Unfortunately, the high intensities at which the seedlings are grown, together with either low natural radiation levels during the rainy season (June - July) and winter or high greenhouse

temperature in summer, often result in batches of badly stretched seedlings being produced (Kim et al., 2008).

Chemical control of growth and/or flower of commercial flower crops became a reality about 30 years ago. Therefore, many people nowadays take it for granted to use these chemicals at lower concentrations. Many methods including PGR, withholding water or nutrients, temperature control, clipping shoots and mechanical stimulation (brushing) are used to control transplant height (Garner and Bjorkman, 1996). Many PGR could be used for height control of many plant species. If PGRs are used to control plant height, height can be influenced by concentration (LeCain et al., 1986; McDanniel, 1986; Ruter, 1992), time of application (Miranda and Carlson, 1980; Gilbertz, 1992), mode of application (Cathey, 1975), formulation (McDaniel, 1986; Ruter, 1992) and media com-

\*Corresponding author. E-mail: brjeong@gnu.ac.kr. Tel.: +82-55-751-5489. Fax: +82-55-757-7542.

**Table 1.** The chemicals and their concentrations used in the nutrient solution for the culture of tomato plug seedlings.

Chemical	Concentration (mg·L <sup>-1</sup> )
Ca(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	472
KH <sub>2</sub> PO <sub>4</sub>	272
MgSO <sub>4</sub> ·7H <sub>2</sub> O	246
KNO <sub>3</sub>	202
NH <sub>4</sub> NO <sub>3</sub>	80
Fe-EDTA	15
MnSO <sub>4</sub> ·4H <sub>2</sub> O	2.1
H <sub>3</sub> BO <sub>3</sub>	1.4
ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.8
CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.2
Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	0.1

position (Barrett, 1982). In many cases, by the time growers apply PGR, stretching of the hypocotyl has already occurred and application to seedlings are ineffective (Pasian and Bennett, 2001). Hence, time of application is very important to use Plant growth regulators (PGRs) because the effect of PGRs depends on time. In addition, the optimal rate of application and sensitivity of plants to each PGR may vary greatly from one species to another (Wang and Blessington, 1990).

Seedling height can be controlled by applying the PGR directly to the seed (Pasian and Bennett, 2001; Pill and Gunter, 2001; Still and Pill, 2003). Treatment to tomato seeds with hypertonic solutions containing uniconazole (0.1, 1, or 10 mg l<sup>-1</sup>) was of little practical value in protecting seedlings from freeze damage, although T<sub>50</sub> (days) and seedling height was reduced (Davis et al., 1990). Soaking seeds in 1000 mg l<sup>-1</sup> paclobutrazol for 6, 16, or 24 h, growth restriction was 61, 37 and 76% in geranium, 30, 38 and 41% in marigold and 31, 31 and 40% respectively. Therefore, PGR application to geranium, marigold and tomato seeds may be feasible using a 6 or 16 h soak in 500 mg l<sup>-1</sup> paclobutrazol (Pasian and Bennett, 2001). Soaking pepper seeds in 1-100 mg l<sup>-1</sup> uniconazole for 24, 72 or 120 h significantly suppressed hypocotyls length and seedling growth (Shin and Jeong, 2002). Cucumber seeds soaked in 250, 500, 1000 or 2000 mg l<sup>-1</sup> paclobutrazol for 6, 12 or 24 h reduced hypocotyls length, stem elongation, leaf area, fresh and dry weight (Cho et al., 2002). Soaking tomato seeds in 50 - 1000 mg l<sup>-1</sup> paclobutrazol for 48 h resulted in lower percentage of germination or emergence than soaking seeds for 24 h (Still and Pill, 2003). Soaking seeds in 50 mg l<sup>-1</sup> paclobutrazol effectively controlled plug height in tomato (Still and Pill, 2006), whereas soaking tomato seeds in 100 mg l<sup>-1</sup> paclobutrazol for 1 h prevented early hypocotyls stretch of tomato seedlings with no long term effects on plant growth (Brigard et al., 2006). Despite these previous studies, there is little information available on the effects of daminozide, ethephon and uniconazole application to control stretching in

seedlings. This study was conducted to evaluate the effect of seed soaking treatment with PGRs for controlling stretching of tomato seedling.

## MATERIALS AND METHODS

Tomato (*Lycopersicon esculentum* Mill. cv. Seogeon and Seokwang) seeds were put in a petri dish (87 mm × 15 mm) filled with 15 ml of PGR solutions and were placed in an environment controlled chamber (25 °C, 80% RH, dark) for 1, 3, or 5 days. Daminozide (B-9) at 10,000, 20,000 or 30,000 mg l<sup>-1</sup>, uniconazole (sumagic) at 100, 200 or 300 mg l<sup>-1</sup>, and ethephon (flore) at 1,000, 2,000 and 3,000 mg l<sup>-1</sup> were used. After soaking treatment, seeds were washed in tap water and were dried in growth chamber (25 °C, 80% RH, dark) for 1 day.

Seeds treated with PGRs solutions were placed in a petri dish with a sheet of filter paper (Whatman no. 2). Then seeds were placed in an environment controlled chamber (25 °C, 80% RH, dark) in a completely randomized design. Equal amount of distilled water was supplied to all treatments. Germination of tomato seeds was checked once a day for 9 days. Percent germination, mean daily germination (MDG), T<sub>50</sub>, and mean germination time (MGT) were evaluated.

Treatments which caused precocious germination during the seed soaking process and treatments which gave percent germination less than 50% in germination experiment were excluded in growth experiment. Seeds were sown in 200-cell (11cc) plug trays containing plug medium (Tosilee medium, Shinan Grow Co., Korea). 3 replicates per treatment and 50 seeds per replicate were used. Seed-sown trays were placed on germination beds with a fogging system for three days in a glasshouse. After seedlings had been emerged, trays were laid out in a completely randomized block design on beds in a glasshouse. A nutrient solution was supplied uniformly for all treatments once a day through a mat subirrigation system. The composition of nutrient solution was based on the formulation used in commercial plug greenhouses (Table 1).

Temperatures of a greenhouse were measured during the experimental period by digital thermometers (Thermo Recorder TR-71S, T and D Crop., Japan). Maximum, minimum and mean temperatures of a greenhouse during the culture period were 35.4, 12.6 and 24.4 °C, respectively.

Emergence was checked for 10 days and growth measurement on seedlings was conducted at 32 days after sowing. Emergence, hypocotyl length, plant height, leaf area, number of leaves, % dry matter, T/R ratio, fresh and dry weights of shoot, root and whole plant and chlorophyll concentration were measured. Leaf area was determined with a leaf area meter (LI-3100 Area meter, LI.COR. Inc., Lincoln, Nebraska USA). Dry weight was measured after 72 h of drying at 60 °C in a dry oven. For measurement of chlorophyll concentration, vigorous and uniform leaves were sampled and extracted with 80% (v/v) acetone for 24 h. Total chlorophyll concentration was determined by measuring absorbance of the extracted solution at 645 nm and 663 nm with a spectrophotometer (Uvikon 922, Kotron Instruments, Italy) according to the procedure developed by Arnon (1949). Chlorophyll concentration (μg·mg<sup>-1</sup>fw) = [(20.29 × A<sub>645</sub>) + (8.02 × A<sub>663</sub>)] × [volume of acetone (ml) ÷ fresh weight (mg)], where A<sub>645</sub> and A<sub>663</sub> are absorbance at 645 and 663 nm, respectively.

Data collected were analyzed for statistical significance by the SAS (Statistical analysis system, V. 6.12, Cary, NC, USA) program. The experimental results were submitted to an analysis of variance (ANOVA).

## RESULTS AND DISCUSSION

Soaking seeds in water or PGRs solution reduced germi-

**Table 2.** Effect of seed treatment of plant growth retardants on percent germination, mean daily germination (MDG), number of days to 50% germination ( $T_{50}$ ), mean germination time (MGT) and percent emergence of 'Seogeon' tomato seeds.

Growth retardant	Concentration (mg l <sup>-1</sup> )	Soaking Duration (day)	Germination (%)	MDG (day)	$T_{50}$ (day)	MGT (day)	Emergence (%)
<b>Control (unsoaked)</b>		0	98.7 a <sup>1</sup>	5.0 a	2.6 de	4.7 cd	96.3 a
<b>Distilled water</b>		1	94.0 ab	4.7 ab	2.9 de	3.4 dg	91.7 a
		3	89.7 bc	4.5 ab	2.7 de	3.2 dg	97.0 a
<b>Daminozide</b>	10,000	1	27.3 e	1.4 e	5.6 c	5.9 bc	- <sup>2</sup>
		3	8.0 g	0.4 g	6.1 c	6.6 b	-
		5	2.0 g	0.1 g	6.7 bc	6.7 b	-
	20,000	1	34.7 e	1.7 e	5.8 c	6.3 b	-
	30,000	1	19.3 f	1.0 f	6.7 bc	6.8 b	-
<b>Uniconazole</b>	100	1	81.3 c	4.1 c	5.9 c	6.6 b	54.7 b
		3	3.0 g	0.2 g	7.3 ab	8.3 a	-
	200	1	51.3 d	2.6 d	7.8 a	8.1 a	30.0 c
<b>Ethephon</b>	1,000	1	95.3 ab	4.8 ab	3.2 de	3.9 f	95.3 a
		3	98.7 a	4.9 a	2.4 e	2.4 g	9.6 d
	2,000	1	96.7 ab	4.8 ab	3.6 d	4.1 de	95.3 a
		3	81.3 g	4.1 c	2.6 de	3.0 eg	7.4 d
	3,000	1	94.7 ab	4.7 ab	3.1 de	3.6 dg	94.0 a
		3	88.7 bc	4.4 bc	3.4 e	2.6 fg	5.3 d
<b>F-test</b>			***	***	***	***	***

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at P = 0.05.

<sup>2</sup> Not measured because of no germination.

NS, \*, \*\*, \*\*\* Nonsignificant or significant at P = 0.05, 0.01, or 0.001, respectively.

nation percentage in both cultivars when compared to the control. Table 2 show a progressive decrease in the rate of germination resulting from the seed treatment with water. Soaking for 5 days killed all the seeds. The water causes the cells to become turgid, the entire seed grows in volume and the seed coat becomes more permeable to oxygen and carbon dioxide. However, over soaking of seeds can damage seeds and unable to exchange gases, which caused inhibition in the metabolic activities, thus germination percent reduced significantly. Soaking seeds of tomato resulted in a gradual loss of germination, however, increased emergence percentage and reduced mean germination time (MGT) when compared to the control. These results are in agreement with previous report (Sabongari and Aliero, 2004).

Seeds of 'Seogeon' tomato did not germinate in 100 mg l<sup>-1</sup> (5 days soaking), 200 mg l<sup>-1</sup> (3 and 5 days soaking) and 300 mg l<sup>-1</sup> (1, 3 and 5 days soaking) uniconazole solutions and 2,000 and 3,000 mg l<sup>-1</sup> (3 and 5 days soaking) daminozide solutions. These results are in agreement with (Pasian and Bennett, 2001; Shin and Jeong, 2002; Still and Pill, 2003), they reported that seed germination was lower with an increase in PGRs concentration and soaking duration of seeds. Seeds of 'Seogeon' tomato soaked in ethephon solution for 5 days germinated during soaking process (Table 2). Percent germination in ethe-

phon treatment was slightly lower than that in the control, but was higher than that in other PGRs treatments. The MDG,  $T_{50}$ , and MGT in ethephon treatment were similar to those in the control. Percent seedling emergence was lower than percent germination in all treatments in the one day ethephon treatment which had similar percent germination and percent emergence as those in the control (Table 2).

Seeds of 'Seogeon' tomato in 100 mg l<sup>-1</sup> (1 and 3 days soaking) and 200 mg l<sup>-1</sup> (1 day soaking) uniconazole solutions germinated, but percent germination was greater than 80% only in the 100 mg l<sup>-1</sup> (one day soaking) uniconazole solution. In addition, percent emergence in these treatments was less than 50%. Although seeds of 'Seogeon' tomato germinated in petri dishes, seedlings did not emerge in trays, especially in 10,000 mg l<sup>-1</sup> and 30,000 mg l<sup>-1</sup> daminozide solution and 100 mg l<sup>-1</sup> (3 days soaking) uniconazole treatment (Table 2).

Table 3 shows similar trends in 'Seokwang' tomato. Most seeds of 'Seokwang' tomato soaked in uniconazole solutions did not germinate except for uniconazole 100 mg l<sup>-1</sup> (one day soaking) and percent germination was less than 3%. Percent germination more than 88% was obtained only in the 100 mg l<sup>-1</sup> (one day soaking) uniconazole solution. In the present study, soaking seeds in PGRs reduced germination percentage significantly in

**Table 3.** Effect of seed treatment of plant growth retardants on percent germination, mean daily germination (MDG), number of days to 50% germination ( $T_{50}$ ), mean germination time (MGT) and percent emergence of 'Seokwang' tomato seeds.

Growth retardant	Concentration ( $\text{mg l}^{-1}$ )	Soaking Duration (day)	Germination (%)	MDG (day)	$T_{50}$ (day)	MGT (day)	Emergence (%)
<b>Control (unsoaked)</b>		0	99.3 a <sup>1</sup>	5.0 a	2.6 de	4.7 cd	97.3 a
<b>Distilled water</b>		1	95.3 ab	4.8 ab	3.0 de	3.5 dg	92.7 a
		3	92.7 ab	4.5 ab	2.8 de	3.3 dg	98.0 a
<b>Daminozide</b>	10,000	1	84.7 c	4.2 c	5.5 bd	5.4 be	86.7 a
		3	20.0 e	1.0 e	6.2 ac	6.7 ac	- <sup>2</sup>
		5	22.7 e	1.1 e	6.1 ac	6.3 ad	-
	20,000	1	67.3 d	3.4 d	5.7 bd	6.2 ad	62.0 bc
		3	2.0 f	0.1 f	7.5 ab	7.5 ab	-
		5	6.7 f	0.3 f	4.5 cg	4.8 cf	-
30,000	1	82.0 c	4.1 c	5.6 bd	6.1 ad	52.0 c	
	3	6.7 f	0.3 f	5.3 be	6.5 ac	-	
	<b>Uniconazole</b>	100	1	88.0 bc	4.4 bc	6.0 ac	6.7 ac
3			2.7 f	0.1 f	4.2 ch	3.3 eg	-
200		1	2.7 f	0.1 f	8.2 a	8.2 a	-
		3	2.0 f	0.1 f	5.0 cf	3.5 eg	-
<b>Ethephon</b>	1,000	1	98.7 a	4.9 a	3.0 ei	3.6 eg	98.7 a
		3	96.0 a	4.8 a	2.6 gi	2.9 eg	87.3 a
	2,000	1	97.3 a	4.9 a	3.4 di	3.9 dg	95.3 a
		3	97.3 a	4.9 a	1.5 l	2.1 g	62.0 bc
	3,000	1	96.7 a	4.8 a	3.4 di	3.9 dg	97.3 a
		3	93.3 ab	4.7 ab	3.6 di	3.9 dg	68.7 b
<b>F-test</b>			***	***	***	***	***

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at  $P = 0.05$ .

<sup>2</sup> Not measured because of no germination.

NS, \*, \*\*, \*\*\* Nonsignificant or significant at  $P = 0.05, 0.01, \text{ or } 0.001$ , respectively.

both cultivars when compared to the water soaking. This might be due to inhibition of gibberellins biosynthesis. Gibberellins are known to stimulate germination of dormant as well as non-dormant seeds of several plant species (Bewley and Black, 1982). The growth retardants paclobutrazol, ancymidol, uniconazole are known to block the oxidation of ent-kaurene to ent-kaurenoic acid which are steps in the GA synthesis pathway of gibberellins (Pressman and Shaked, 1991).

Of the 3 PGRs tested, uniconazole was shown to be the most effective retardant followed by ethephon and daminozide. Table 4 shows that hypocotyl length and plant height of 'Seogeon' tomato seedlings were remarkably suppressed in the  $100 \text{ mg l}^{-1}$  (1 day soaking) and  $200 \text{ mg l}^{-1}$  (1 day soaking) uniconazole treatments as compared to the control and other treatments. The plant growth retardant uniconazole is known to affect the levels of endogenous gibberellins. Gibberellins are known to induce elongation (Kurepin et al., 2006), while triazole PGRs reduces gibberellins levels and causes a decrease in shoot growth (Davis and Curry, 1991). Uniconazole reduce stem length in many plant species such as hibis-

cus (Wang and Gregg, 1991), chrysanthemum (Schuch, 1994), kalanchoe (Hwang et al., 2008, 2009), pepper (Shin and Jeong, 2002). An increase in ethephon concentration from  $1000 \text{ to } 3000 \text{ mg l}^{-1}$  resulted in a significant decrease in hypocotyls length when compared to the control, however, plant height did not affect significantly at any concentrations of ethephon. Fresh and dry weights of shoot significantly decreased when seeds were soaked in  $100 \text{ or } 200 \text{ mg l}^{-1}$  uniconazole. At  $100 \text{ mg l}^{-1}$  uniconazole fresh weight of root did not affect when compared to the control, but dry weight reduced significantly. When seeds were treated in  $1000 \text{ mg l}^{-1}$  ethephon root fresh weight increased significantly, but dry weight of root was not changed. At  $2000 \text{ mg l}^{-1}$  ethephon fresh and dry weights of shoot significantly decreased (Table 4). Leaf area and number of leaves were not reduced when the seeds were treated with ethephon, but significantly reduced by uniconazole treatment. Ethephon treatment ( $1000 \text{ or } 2000 \text{ mg}$ ) increased chlorophyll content when compared to the control. However, at  $3000 \text{ mg}$  ethapon chlorophyll content was not increased. When the seeds were soaked in uniconazole for 1 day chlorophyll content was reduced when

**Table 4.** Effect of seed treatment of plant growth retardants on growth of 'Seogeon' tomato seedlings measured at 32 days after sowing in plug trays.

Growth retardant	Concentration (mg·L <sup>-1</sup> )	Soaking Duration (day)	Hypocotyl Length (cm)	Plant Height (cm)	Fresh wt. (g)		Dry wt. (g)	
					Shoot	Root	Shoot	Root
<b>Control (unsoaked)</b>		0	8.5 a <sup>1</sup>	30.1 a	3.3 a	0.29 c	0.25 a	0.02 ab
<b>Distilled water</b>		1	7.0 c	27.9 a	2.6 b	0.44 ab	0.20 ab	0.02 a
		3	7.7 bc	30.4 a	3.3 a	0.35 c	0.22 ab	0.02 ab
<b>Uniconazole</b>	100	1	3.4 d	6.7 b	1.0 c	0.30 c	0.08 c	0.01 b
	200	1	3.4 d	5.4 b	0.5 c	0.12 d	0.03 d	0.01 c
<b>Ethephon</b>	1,000	1	8.0 ab	28.2 a	3.2 a	0.48 a	0.22 ab	0.02 a
	2,000	1	7.2 bc	28.8 a	2.5 b	0.31 c	0.18 b	0.02 b
	3,000	1	6.8 c	28.4 a	3.1 a	0.33 c	0.20 ab	0.02ab
<b>F-test</b>			***	***	***	***	***	***

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at  $P = 0.05$ .  
NS,\*, \*\*, \*\*\*Nonsignificant or significant at  $P = 0.05, 0.01, \text{ or } 0.001$ , respectively.

**Table 5.** Effect of seed treatment of plant growth retardants on growth of 'Seogeon' tomato seedlings measured at 32 days after sowing in plug trays.

Growth retardant	Concentration (mg·L <sup>-1</sup> )	Soaking Duration (day)	Leaf area (cm <sup>2</sup> )	No. of leaves	Chlorophyll (µg·mg <sup>-1</sup> fw)	Dry matter (%)	T/R ratio
<b>Control (unsoaked)</b>		0	44.1 a <sup>1</sup>	4.1 a	2.64 bc	7.24 ab	13.94 a
<b>Distilled water</b>		1	40.3 a	4.0 ab	3.12 a	7.37 a	9.62 bd
		3	42.9 a	4.0 ab	2.75 ab	6.51 ac	11.53 ab
<b>Uniconazole</b>	100	1	21.3 b	3.6 c	2.20 d	7.35 a	5.64 d
	200	1	4.5 c	3.0 d	2.27 cd	5.85 c	7.05 cd
<b>Ethephon</b>	1,000	1	43.4 a	4.0 ab	3.09 a	6.73 ac	9.62 bd
	2,000	1	39.9 a	4.0 ab	3.08 a	6.79 ac	11.84 ab
	3,000	1	43.8 a	3.9 b	2.79 ab	6.30 bc	11.15 ac
<b>F-test</b>			***	***	***	*	***

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at  $P = 0.05$ .  
NS,\*, \*\*, \*\*\*Nonsignificant or significant at  $P = 0.05, 0.01, \text{ or } 0.001$ , respectively.

compared to the control (Table 5). Chlorophyll content was enhanced by PGRs treatment (Starman et al., 1990; Tekalign and Hammes, 2004). In contrast, chlorophyll content was reduced by PGRs treatment (Knypl, 1969; Davis et al., 1988). Differential effect of PGR on chlorophyll content may be due to its accumulation and endogenous level of cytokinins. Percent dry matter was significantly lower in 200 mg l<sup>-1</sup> uniconazole treatment. T/R ratio slightly decreased in 100 mg l<sup>-1</sup> (1 day soaking) and 200 mg l<sup>-1</sup> (1 day soaking) uniconazole treatments (Table 5).

Table 6 and Figure 1 show that hypocotyl length and plant height of 'Seokwang' tomato seedlings were remarkably suppressed in 100 mg l<sup>-1</sup> (1 day soaking) uniconazole treatment as compared to the control and other treatments. Plant height was increased in 10,000 mg l<sup>-1</sup> daminozide for 1 day when compared to the control. Soaking seeds in 2000 or 3000 mg l<sup>-1</sup> ethephon for 1 day plant height increased. However, 3 days soaking decreased plant height. The results are in agreement with

(Shin and Jeong, 2002; Still and Pill, 2003) they reported that plant height decreased as soaking period increased. Leaf area, number of leaves, and T/R ratio slightly decreased in 100 mg l<sup>-1</sup> (1 day soaking) (Table 6).

Chlorophyll concentration and percent dry matter were not significantly different in all treatments (Table 7). Uniconazole was effective in inhibiting seedling height of both cultivars at a very low concentration than that of daminozide and ethephon, but significantly affect the emergence percentage. Hence, further studies are needed in order to obtain maximum emergence and this may be possible by reducing uniconazole concentration.

In conclusion, the growth of tomato seedlings was efficiently regulated by uniconazole 100 mg l<sup>-1</sup> (1 day soaking) treatment. These results should be helpful in allowing plant nurseries to control the stretching of tomato seedling. Although these methods successfully control stretching, seeds soaked in PGR solutions germinated, it tended not to emerge well. A seed soaking

**Table 6.** Effect of seed treatment of plant growth retardants on growth of 'Seokwang' tomato seedlings measured at 32 days after sowing in plug trays.

Growth retardant	Concentration (mg l <sup>-1</sup> )	Soaking Duration (day)	Hypocotyl Length (cm)	Plant Height (cm)	Fresh wt. (g)		Dry wt. (g)	
					Shoot	Root	Shoot	Root
<b>Control (unsoaked)</b>		0	7.8 a <sup>1</sup>	28.7 a	2.7 a	0.37bc	0.18 a	0.02 b
<b>Distilled water</b>		1	6.8 ac	28.8ac	2.4 a	0.34bd	0.16 c	0.02 b
		3	7.3 ab	30.0 b	2.7 a	0.52 a	0.19 a	0.09 a
<b>Daminozide</b>	10,000	1	7.1 ac	31.5 a	2.8 a	0.31bd	0.19 a	0.02 b
	20,000	1	6.3 bc	28.1ac	2.8 a	0.31bd	0.15ac	0.02 b
	30,000	1	6.6 ac	24.4 d	2.3 a	0.41ab	0.16ab	0.02 b
<b>Uniconazole</b>	100	1	3.2 d	6.0 e	0.8 b	0.22 d	0.05 d	0.01 b
<b>Ethephon</b>	1,000	1	6.9 ac	28.2ac	2.4 a	0.39ac	0.16ac	0.02 b
		3	6.0 c	26.7bd	2.2 a	0.27bd	0.12 c	0.01 b
	2,000	1	7.4 ab	30.4 a	2.8 a	0.41ab	0.19 a	0.02 b
		3	5.9 c	24.6 d	2.2 a	0.25cd	0.13bc	0.01 b
	3,000	1	7.0 ac	29.4ac	2.7 a	0.39ac	0.19 a	0.02 b
		3	6.3 bc	26.4cd	2.3 a	0.31bd	0.12 c	0.02 b
<b>F-test</b>			***	***	*	**	***	NS

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at  $P = 0.05$ .

<sup>N</sup>S, \*, \*\*, \*\*\* Nonsignificant or significant at  $P = 0.05, 0.01, \text{ or } 0.001$ , respectively.



**Figure 1.** Tomato 'Seokwang' cultivar grown with different plant growth retardant (daminozide, uniconazole, and ethephon) soaking concentration treatment harvested at 32 days after sowing in plug tray. Soaked seeds were placed for one day.

**Table 7.** Effect of seed treatment of plant growth retardants on growth of 'Seokwang' tomato seedlings measured at 32 days after sowing in plug trays.

Growth retardant	Concentration (mg l <sup>-1</sup> )	Soaking Duration (day)	Leaf area (cm <sup>2</sup> )	No. of leaves	Chlorophyll (µg mg <sup>-1</sup> fw)	Dry matter (%)	T/R ratio
<b>Control (unsoaked)</b>		0	48.4 b <sup>1</sup>	4.0 a	2.54 a	6.79 b	11.89 ab
<b>Distilled water</b>		1	38.6 df	4.0 ab	2.53 a	6.46 b	9.00 ae
		3	46.5 bc	4.1 ab	2.53 a	8.88 a	9.00 ae
<b>Daminozide</b>	10,000	1	41.5 cd	4.0 a	2.19 a	6.79 b	10.92 ad
	20,000	1	33.7 f	4.1 ab	2.34 a	5.42 b	12.08 a
	30,000	1	39.4 de	4.0 b	2.38 a	6.74 b	7.40 cg
<b>Uniconazole</b>	100	1	13.8 h	3.1 c	2.26 a	6.42 b	4.15 fi
<b>Ethephon</b>	1,000	1	42.0 cd	4.1 ab	2.32 a	6.41 b	7.91 bf
		3	28.6 g	4.0 ab	2.36 a	5.46 b	8.95 ae
	2,000	1	45.0 bc	4.2 a	2.33 a	6.82 b	9.00 ae
		3	35.3 ef	4.1 ab	2.25 a	5.93 b	7.22 cg
	3,000	1	41.6 cd	4.0 b	2.21 a	6.72 b	6.08 eh
		3	56.1 a	4.0 b	2.46 a	5.40 b	8.77 ae
<b>F-test</b>			***	***	NS	*	*

<sup>1</sup> Mean separation within columns by Duncan's multiple range test at  $P = 0.05$ .

NS, \*, \*\*, \*\*\* Nonsignificant or significant at  $P = 0.05, 0.01, \text{ or } 0.001$ , respectively.

method which is effective in suppressing hypocotyl length and seedling height without affecting germination and seedling emergence significantly would be ideal and it is necessary to research effects that a seed soaking method makes on flowering and fruit setting as well as general growth.

## ACKNOWLEDGEMENTS

This work partially supported by the BK21 Program, the Ministry of Education, Science and Technology, Korea.

## REFERENCES

- Arnon DI (1949). Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. Plant Physiol. 24: 1-15.
- Barrett JE (1982). Chrysanthemum height control by ancymidol, pp333, and EL-500 dependent on medium composition. Hort. Sci. 17: 896-897.
- Bewley JD, Black M (1982). Physiology and biochemistry of seeds in relation to germination. II. Viability, dormancy and environment control. Springer-Verlag, Berlin.
- Brigard JP, Harkess RL, Baldwin BS (2006). Tomato early seedling height control using a paclobutrazol seed soak. Hort. Sci. 41: 768-772.
- Cathey HM (1975). Comparative plant growth retarding activities of ancymidol with ACPC, phosfon, chlormequat, and SADH on ornamental plant species. Hort. Sci. 10: 204-216.
- Cho KC, Yang WM, Kim WS, Chung SJ (2002). Growth retardation of cucumber (*Cucumis sativus* L.) seedlings by treatment of paclobutrazol and removal of retardation by foliar spray of gibberellins. J. Korean Soc. Hort. Sci. 43: 415-420.
- Choi JM, Ahn JW, Ku JH, Lee YB (1997). Effect of medium composition of physical properties of soil and seedling growth of red-pepper in plug system. J. Korean Soc. Hort. Sci. 38: 618-624.
- Davis TD, Curry EA (1991). Chemical regulation of vegetative growth. Crit. Rev. Plant Sci. 10: 151-188.
- Davis TD, Eills JE, Walser RH (1990). Emergence, growth, and freezing tolerance of tomato seedlings grown from uniconazole-treated seed. Hort. Sci. 25: 312-313.
- Davis TD, Steffens GL, Sankkhla N (1988). Triazole plant growth regulators. Hort. Rev. 10: 63-105.
- Garner LC, Bjorkman T (1996). Mechanical conditioning for controlling excessive elongation in tomato transplants: Sensitivity to dose, frequency, and timing of brushing. J. Am. Soc. Hort. Sci. 121: 894-900.
- Gilbert DA (1992). Chrysanthemum response to timing of paclobutrazol and uniconazole sprays. Hort. Sci. 27: 322-323.
- Hwang SJ, Lee MY, Sivanesan I, Jeong BR (2008). Growth control of kalanchoe cultivars Rako and Gold Strike by application of paclobutrazol and uniconazole as soaking treatment of cuttings.
- Hwang SJ, Lee MY, Sivanesan I, Jeong BR (2009). Suppression of stretchiness in pot *Kalanchoe blossfeldiana* Poelln. Rako by application of plant growth retardants as recycled subirrigational supply. Propagation Ornamental Plants, 9: 26-34.
- Kim IS, Zhang C, Kang HM, Mackay B (2008). Control of Stretching of cucumber and tomato plug seedlings using supplement light. Hort. Environ. Biotechnol. 49: 287-292.
- Knypl JS (1969). Inhibition of chlorophyll synthesis by growth retardants and coumarin, and its reversal by potassium. Nature, 224: 1025-1026.
- Kurepin LV, Pharis RP, Richard P, Reeid DM, Chinnappa CC (2006). Involvement of gibberellins in the stem elongation of sun and shade ecotypes of *Stellaria longipes* that is induced by low light irradiance. Plant Cell Environ. 29: 1319-1328.
- LeCain DR, Scheke KA, Wample RL (1986). Growth-retarding effects of paclobutrazol on weeping fig. Hort. Sci. 21: 1150-11520.
- McDaniel GL (1986) Comparison of paclobutrazol, flurprimidol and tetcyclacis for controlling poinsettia height. Hort. Sci. 21: 1161-1163.
- Miranda RM, Carlson WH (1980). Effect of timing and number of applications of chlormequat and ancymidol on the growth and flowering of seed geraniums. J. Am. Soc. Hort. Sci. 105: 273-277.
- Pasian CC, Bennett MA (2001). Paclobutrazol soaked marigold, geranium, and tomato seeds produce short seedlings. Hort. Sci. 36: 721-723.

- Pill WG, Gunter JA (2001). Emergence and shoot growth of cosmos and marigold from paclobutrazol-treated seed. *J. Environ. Hort.* 19: 11-14.
- Pressman E, Shaked R (1991). Interactive effects of Gas, CKs and growth retardants on the germination of celery seeds. *Plant Growth Regul.* 10: 65-72.
- Ruter JM (1992). Growth and flowering response of butterfly-bush to paclobutrazol formulation and rate of application. *Hort. Sci.* 27: 929.
- Sabongari S, Aljero BL (2004). Effects of soaking duration on germination and seedling growth of tomato (*Lycopersicon esculentum* Mill.). *Afr. J. Biotechnol.* 3: 47-51.
- Schuch UK (1994). Response of chrysanthemum to uniconazole and daminozide applied as dip to cuttings or as foliar spray. *J. Plant. Growth Regul.* 13: 115-121.
- Shin WG, Jeong BR (2002). Seed treatment of growth retardants for height suppression of pepper plug seedlings. *J. Korean Soc. Hort. Sci.* 43: 565-570.
- Starman TW, Kelly JW, Pemberton HB (1990). The influence of ancymidol on morphology, anatomy, and chlorophyll levels in developing and mature *Helianthus annuus* leaves. *Plant growth Regul.* 9: 193-200.
- Still JR, Pill WG (2003). Germination and emergence, and seedling growth of tomato and impatiens in response to seed treatment with paclobutrazol. *Hort. Sci.* 38:1201-1204.
- Still JR, Pill WG (2006). Growth and stress tolerance of tomato seedlings (*Lycopersicon esculentum* Mill.) in response to seed treatment with paclobutrazol. *J. Hort. Sci. Biotechnol.* 81: 197-203.
- Tekalign T, Hammes PS (2004). Response of potato grown under non-inductive condition to paclobutrazol: shoot growth, chlorophyll content, net photosynthesis, assimilate partitioning, tuber yield, quality, and dormancy. *Plant Growth Regul.* 43: 227-236.
- Wang YT, Blessington TM (1990). Growth of four tropical foliage species treated with paclobutrazol or uniconazole. *Hort. Sci.* 25: 202-204.
- Wang YT, Gregg (1991). Modification of Hibiscus growth by treating unrooted cuttings and potted plants with uniconazole or paclobutrazol. *J. Plant Growth Regul.* 10: 47-51.