

Full Length Research Paper

Influence of volatile compounds and food preservatives in the management of fumonisins (B₁) production by *Fusarium moniliforme*

Narasimha Rao Kollu*, Vijaypal Reddy Baireddy, Girisham Sivadevuni and Madhusudhan Reddy Solipuram

Department of Microbiology, Kakatiya University, Warangal 506009, Andhra Pradesh, India.

Accepted 15 June, 2009

Efficacy of volatile compounds and food preservatives in the management of fumonisins B₁ production by *Fusarium moniliforme* was assessed *in vitro*. The toxin production was analyzed by thin layer chromatographic technique. Among different volatile compounds studied, formic acid and propionic acid inhibited the growth and fumonisins (B₁) production by *F. moniliforme*. Aniline, benzaldehyde, and formaldehyde can suppress fumonisin B₁ production to maximum extent. Camphor at a concentration of 15 mg/ml could inhibit fumonisin B₁ production among different food preservatives studied. From the above results it is concluded that volatile compounds and food preservatives can be employed in the management of fumonisins (B₁).

Key words: Food preservatives, volatile compounds, *Fusarium moniliforme*, fumonisins.

INTRODUCTION

Fumonisins are a group of structurally related mycotoxins produced by the fungus *Fusarium moniliforme* and several other *Fusarium* species (Nelson et al., 1991, 1992), which occur in corn and corn-based foods and feeds (Shephard et al., 1996a). Fumonisins, primarily fumonisin B₁ (FB₁) are hepatotoxic and nephrotoxic in many animal species. They have been shown to cause farm animal diseases such as, equine leuko encephalomalacia (ELEM) and porcine pulmonary oedema (PPE) (Norred, 1993). Fumonisin B₁ has been shown to be a liver and kidney carcinogen in rodents (Gelderblom et al., 1996; NTP 1999). Although the role of fumonisins in human health is unclear, consumption of *F. moniliforme* contaminated corn has been correlated with human oesophageal cancer in areas of southern Africa and China (Marasas, 1996). Galvano et al. (2001) discussed elegantly the strategies to be adopted to control fumonisins contamination of food grains. Fumonisins are generally concentrated on the pericarp layer of corn kernels (Sydenham et al.,

1993). Sydenham et al. (1994) could reduce the fumonisins level very significantly by sorting out fine particles of less than 3 mm size. Sydenham et al. (1993) reduced the fumonisins level in corn by treating with 0.1 M calcium hydroxide for 24 h at 25°C. Norred et al. (1991) could reduce the fumonisins level in corn by ammoniation. Detoxification of fumonisins in maize by ammonia could reduce to 30% (Chourasia, 2001). Bhat et al. (1997) recorded variation in fumonisins B₁ level in hybrids and hybrids of maize. Yamgish et al. (1997) could produce mutant of *F. proliferatum* which produced little or no fumonisins and can be used as a competitor for fumonisins producing strains and thus minimize fumonisins contamination. Beekrun et al. (2003) indicated that naturally occurring phenolics of plants such as chlorophyllin, iroka, benzoic acid, caffeic acid, ferulic acid and vanillic acid are effective inhibitors of fumonisins production by *F. verticillioides*. Ono et al. (2002) postulated that the decreased seed moisture of freshly harvested maize through predrying of storage grains can control fungal growth and fumonisins contamination (Figure 1).

In the light of the above facts some food preservatives and different volatile compounds on the growth and pro-

*Corresponding author. E-mail: kollu2006@yahoo.com




Figure 1. Structure of fumonisins B₁.

duction of fumonisins (B₁) by *F. moniliforme* was studied.

MATERIALS AND METHODS

Chemicals and microorganism

Fumonisins (B₁) was purchased from Sigma-Aldrich, Mumbai, India. Yeast extract, NaNO₃, KH₂PO₄, MgSO₄·7H₂O, KCl and all other chemicals of highest available purity were obtained from Himedia, Mumbai, India.

F. moniliforme MRC 826 was purchased from *Fusarium* research center collection (Pennsylvania State University, University park, Pa.). Stock cultures were maintained on potato dextrose agar slants at 4°C and sub cultured for every 3 months.

Influence of volatile compounds

50 ml of Czapeck's medium (Yeast extract 10 g; NaNO₃ 2 g, KH₂PO₄ 1 g; MgSO₄·7H₂O 0.5 g and distilled water 1000 ml) was taken in 250 ml Erlenmeyer conical flasks containing glass vial hanged from mouth of conical flask (5 ml capacity) and sterilized. Volatile compound viz. propanol, butanol-1, isoamyl alcohol, ethyl alcohol, chloroform, formaldehyde, benzaldehyde, formic acid, acetic acid, ethyl acetate, acetone, benzene, toluene, hexane, acetonitrile, kerosene, petrol, propionic acid, xylene and aniline, control are poured into glass vial separately after sterilization and flasks were inoculated with *F. moniliforme* and incubated at 26 ± 2°C for 20 days.

Influence of food preservatives

Food preservatives viz boric acid, eucalyptus oil, camphor and naphthalene were added individually to 50 ml of sterilized Czapeck's medium contained in 250 ml conical flasks to get 2.5, 5.0, 10.0, 15.0 mg/ml and inoculated with *F. moniliforme*. All the flasks thus prepared were incubated at 26 ± 2°C for 20 days.

Qualitative and quantitative analysis of fumonisins B₁

At the end of incubation period cultures were harvested and the mycelial mat was dried and extracted with ethyl acetate in soxhlet apparatus for 24 h. Ethyl acetate was evaporated to near dryness and redissolved in methanol. A portion of methanol solution was spotted on TLC plates and developed in water: methanol (1:3) mixture. The plates thus developed were sprayed with p-anisaldehyde (0.2% in ethanol) and heated at 120°C until colour developed. Fumonisins appeared as brown colour spot. Fumonisins was estimated quantitatively as suggested by Sydenham (1989). The biomass attained by the fungus was estimated by drying the mycelium to complete dryness at 65 - 70°C for 48 h and weighed after cooling to room temperature.

RESULTS AND DISCUSSION

Influence of different volatile compounds

Efficacy of different volatile compounds was studied on the growth and fumonisins B₁ production by *F. moniliforme* and the results are presented in Table 1. Propionic acid and formic acid were responsible for total inhibition of growth of *F. moniliforme* and fumonisins B₁ production. Whereas, aniline, formaldehyde and benzaldehyde were also effective as they were responsible for inhibition of fumonisins production by *F. moniliforme* to maximum extent. On the other hand, vapors of propanol failed to inhibit the growth and fumonisins production. Acetic acid, acetone, butanol and benzene were also not much effective in the control of growth and fumonisins production by *F. moniliforme*. The rest of the volatile compounds were mild in their toxicity as they were responsible for decreased amount of fumonisins production.

Similarly, different volatiles inhibited the growth of *F.*

Table 1. Effect of volatile compounds on growth and fumonisins production by *F. moniliforme*.

Name of the compound	Final pH	Dry weight (mg/ml)	Fumonisin B ₁ (µg/ml)
Propanol	7.0	14.23	0.90
Butanol-1	5.8	13.89	0.64
Isoamyl alcohol	5.3	8.43	0.23
Ethyl alcohol	6.8	10.49	0.33
Chloroform	5.4	1.23	0.34
Formaldehyde	5.2	2.09	0.15
Benzaldehyde	4.8	4.80	0.18
Formic acid	3.9	--	--
Acetic acid	3.4	0.18	0.72
Ethyl acetate	6.6	8.42	0.28
Acetone	7.3	4.20	0.71
Benzene	6.1	12.01	0.63
Toluene	5.4	8.20	0.33
Hexane	6.8	11.80	0.31
Acetonitrile	6.3	10.18	0.30
Kerosene	5.4	7.14	0.21
Petrol	5.1	8.30	0.24
Propionic acid	3.4	--	--
Xylene	5.6	6.22	0.21
Aniline	5.8	5.81	0.19
Control	6.5	14.20	1.09

moniliforme to varying level. Acetic acid and chloroform were responsible for maximum inhibition of growth of *F. moniliforme* while, propanol, butanol, benzene, hexane, ethyl alcohol and acetonitrile failed to inhibit the growth of *F. moniliforme*. The rest of the volatile compounds were responsible for intermediate degree of inhibition of growth of *F. moniliforme*. The present observations are in agreement with those of Surekha and Reddy (1992), Laxma and Reddy (1992) and Al-Hilli et al. (1992) who also recorded the control of mould infestation and mycotoxin contamination by the volatiles tested by them.

The final pH recorded was highly variable. The final pH in media containing propanol, ethyl alcohol, ethyl acetate, acetone and hexane was near neutral, while medium containing propionic acid and acetic acid and formic acid was highly acidic.

Influence of food preservatives

Influence of different food preservatives were studied on the growth and fumonisins B₁ production by *F. moniliforme* and the results were recorded in Table 2. From the Table 2 it is evident that different food preservatives exerted significant influence on fumonisins production by *F. moniliforme*. Boric acid at 2.5 mg/ml of concentration stimulated the fumonisins (B₁) production, while at higher concentrations it inhibits fumonisins (B₁) production. The degree of inhibition increased with the increase in boric

acid concentration. Eucalyptus oil was toxic and its degree of inhibition increased with the increase in eucalyptus oil concentration. Vegetative growth of the fungus, which was very good at 2.5 mg/ml concentration, decreased with increase in concentration of eucalyptus oil up to 15 mg/ml. Similarly, camphor inhibited the fumonisins (B₁) production by *F. moniliforme* at 15 mg/ml. Napthalene was also responsible for increased inhibition of fumonisins production and vegetative growth of *F. moniliforme* with increased concentration.

Conclusion

Based on the above results, it can be concluded that food preservatives tried in the present investigations were effective in checking *F. moniliforme* infestation and fumonisins (B₁) production and can be exploited in protection of food grains from infestation of *F. moniliforme* and fumonisin contamination.

ACKNOWLEDGEMENTS

The authors thank the Head, Department of Microbiology, Kakatiya University, Warangal 506009, Andhra Pradesh, India, for providing necessary facilities and Council of Scientific and Industrial Research (CSIR) for financial assistance.

Table 2. Effect of some food preservatives on growth and fumonisins production by *F. moniliforme*.

Compound	Conc. (mg/ml)	Dry weight (mg/ml)	Fumonisins B ₁ (µg/ml)
Boric acid	2.5	18.3	1.00
	5.0	15.6	0.74
	10.0	9.8	0.65
	15.0	4.3	0.43
Eucalyptus oil	2.5	17.6	0.73
	5.0	16.4	0.67
	10.0	8.0	0.42
	15.0	6.3	0.32
Camphor	2.5	8.6	0.66
	5.0	7.2	0.46
	10.0	4.8	0.41
	15.0	2.9	0.10
Napthalene	2.5	13.4	0.72
	5.0	12.8	0.71
	10.0	8.3	0.40
	15.0	4.3	0.42
Control	--	14.9	0.91

REFERENCES

- Ali-Hilli AL, Smith JE (1992). Influence of propionic acid on growth and aflatoxin production by *Aspergillus flavus* in liquid submerged and solid substrate conditions. *J. Environ. Path. Toxi. Oncol.*, 11: 121-124.
- Beekrun S, Govinden R, Padayachoe T, Odhav B (2003). Naturally occurring phenols: A detoxification strategy for fumonisin B₁. *Food additives contaminants*, 20: 490-493.
- Bhat RV, Shetty PHI, Amruth RP, Sudershan RV (1997). A food borne disease outbreak due to the consumption of mouldy sorghum and maize containing fumonisins mycotoxins. *Clinic. Toxicol.* 35: 249-255.
- Chourasia HK (2001). Efficacy of ammonia in detoxification of fumonisin contaminated corn. *Indian J. Exp. Biol.* 39: 493-495.
- Galvano F, Piva A, Ritiene A, Galvano G (2001). Dietary strategies to counteract the effects of mycotoxins. A review: *J. Food Prot.* 64: 120-131.
- Gelderblom WCA, Snyman SD, Abel S, Lebepe-Mazur S, Smuts CM, Vander Westhuizen L, Marasas WFO, Victor TC, Knasmuller S, Huber W (1996). Hepatotoxicity and carcinogenicity of the fumonisins in rats: A review regarding mechanistic implications for establishing risk in humans. *Adv. Exp. Med. Biol.* 392: 279-296.
- Laxma Reddy G, Reddy SM (1992). Control of satratoxin production by *Stachybatrys atra* through food preservatives. *Nat. Acad. Sci. Lett.* 15: 175-177.
- Marasas WFO (1996). Fumonisins: History, world wide occurrence and impact. *Adv. Exp. Med. Biol.* 392: 1-17.
- Nelson PE, Plattner RD, Shackelford DD, Desjardin AE (1991). Production of fumonisins by *Fusarium moniliforme* strains from various substrates and geographic areas. *Appl. Environ. Microbiol.* 57: 2410-2412.
- Nelson PE, Plattner RD, Desjardin AE (1992). Production of fumonisins by *Fusarium* species other than *F. moniliforme* in section *Liseda* and some related species. *Appl. Environ. Microbiol.* 58: 984-989.
- Norred WP, Vos KA, Bacon CW, Rirrlay RT (1991). Effectiveness of ammonia treatment in detoxification of fumonisin-contaminated corn. *Food chem. Toxicol.* 29: 815-819.
- Norred WP (1993). Fumonisins mycotoxins produced by *Fusarium moniliforme*. *J. Toxicol. Environ. Health*, 38: 309-328.
- NTP (1999). Technical Report - Toxicology and carcinogenesis studies of fumonisins B₁ in F344/N rats and B₆C₃F₁ mice. NTPTR 496 NIH publication No. 99-3955.
- Ono EVS, Sasaki EY, Hashimoto EH, Hara LN, Correa B, Itano EN, Sugiura T, Ueno Y and Hirroka EY (2002). Post harvest stage of corn: Effect of beginning moisture content on mycoflora and fumonisin contamination. *Food add. Contam.* 19: 1081-1090.
- Shephard GS, Thiel PG, Stockenstrom S and Sydenham EW (1996a). World wide survey of fumonisin contamination of corn and corn-based products. *J. AOAC int.* 79: 671-687.
- Surekha M, Reddy SM (1992). Efficacy of food preservatives in the control of penitrem B production by *penicillium aurantiogriseum*. *J. food Sci. Technol.* 14: 377-378.
- Sydenham EW, Ishephard GS, Thiel PG, Marasas WFO, Rheeder LEP, Peralta S, Gonzalez HHL, Resnik SL (1993). Fumonisins in Argentina field-trial corn. *J. Agric. Food Chem.* 41: 891-895.
- Sydenham EW, Vander Westhuizen L, Stockenstrom S, Shephard GS, Thiel PG (1994). Fumonisin contaminated maize: physical treatment for the partial decontamination of bulk shipments. *Food Add. Conta.* 11: 25-32.
- Sydenham EW (1989). The chromatographic determination of fusarium toxins in maize associated with human/oesophageal cancer. M.Sc Thesis, University of Capetown, South Africa.