

Full Length Research Paper

Antimicrobial activity of the fruit extracts of *Coccinia indica*

Syed Zeenat Shaheen*, Krishna Bolla, Kandukuri Vasu and M. A. Singara Charya

Department of Microbiology, Kakatiya University, Warangal - 506009, Andhra Pradesh, India.

Accepted 26 October, 2009

The bioactive compounds of fruits of *Coccinia indica* were investigated for antibacterial activity against some pathogenic bacteria. The aqueous extracts did not show much significant activity, while the organic extracts (petroleum ether and methanol) showed the highest activity against the test bacteria. The activity was more pronounced on gram-positive organisms with *Staphylococcus aureus* being more susceptible and *Salmonella paratyphi A* being more resistant. Phytochemical analysis showed that the extracts contain alkaloids, tannins, saponins, flavonoids, glycosides and phenols.

Key words: *Coccinia indica*, antibacterial activity, phytochemical analysis.

INTRODUCTION

Most of the modern researches on herbal medicine have hinged around traditional folklore medicine. The modern medicine has brought with it an array of drugs, none of which is non-toxic and quite safer for human consumption. There are hundreds of medicinal plants that have a long history of curative properties against various diseases and ailments. However, screening of plants for their activity is very essential and needs urgent attention in order to know the value of the plant. The screening of the plants for their biological activity is done on the basis of either their chemotaxonomic investigation or ethnobotanical knowledge for a particular disease. Identification of a particular compound against a specific disease is a challenging long process. Importance of the plant lies in their biologically active principles. There are two types of plant chemicals, primary metabolites such as sugars, proteins, amino acids, chlorophylls, etc. The other category of chemicals is called secondary metabolites, which includes alkaloids, terpenoids, saponins and phenolic compounds. These chemicals exert a significant physiological effect on the mammalian system.

Microorganisms have developed resistance to many antibiotics and this has created immense clinical problem in the treatment of infectious diseases (Davis, 1994). The increase in resistance to microorganisms due to indiscri-

minate use of antimicrobial drugs forced scientists to search for new antimicrobial substances from various sources including medicinal plants (Karaman et al., 2003). Another driving factor for the renewed interest in past 20 years has been the rapid rate of plant species extinction.

Around 12,000 plant secondary metabolites of antimicrobial importance have been isolated. These compounds fall in one of the major groups of compounds like phenols, quinines, flavonoids, tannins, terpenoids, alkaloids and other mixtures (Scultes, 1978). Infectious diseases account for high proportion of health problems in the developing countries (Sashi et al., 2003). In India, about 2,500 species of plants are used for medicinal purposes and about 90% of the medicinal plants provide raw materials for the herbal pharmaceuticals, which are collected from wild habitats. The rich knowledge base of countries like India in medicinal plants and healthcare has led to the keen interest by pharmaceutical companies to use this knowledge as a resource for research and development programs in the pursuit of discovering novel drugs (Rajasekharan and Ganeshan, 2002). However, several plants are used in India in the form of crude extracts without scientific evidence of efficacy (Ahmed et al., 1998). At this juncture it is of interest to determine the scientific basis for the traditional use of these plants.

Coccinia indica Wight and Arnold (Cucurbitaceae) commonly known as 'Ivy gourd' and 'Kundru' in Hindi is a perennial tendril climber, available in wild and cultivated form. It is the native of Central Africa, India and Asia and

*Corresponding author. E-mail: zeenatsd@gmail.com.

Table 1. Various medicinal uses of different parts of *Coccinia indica*.

Part used	Medicinal value
Leaves	When mixed with gingely oil treats ring worm, psoriasis and itch; when mixed with ghee cures sores, skin diseases, skin eruptions of small pox; causes cooling effect to eyes, heals big ulcers, small lesions of scabies, anuria and alleviate body heat.
Stem	Antispasmodic effect.
Fruit	Green fruit when chewed cures sores on tongue; raw fruit used as vegetable; dried fruit removes eczema.
Bark	Dried bark has cathartic properties
Juice	Juice of tuberous roots, stem and leaves serve as a cure for diabetes, intermittent glycosuria, enlarged glands and skin diseases such as pityriasis and also treats urinary tract infection, related troubles.
Powder	Used in treating gastro – intestinal disturbances, liver weakness, dysentery, vomiting, wormal infestation, purifies blood, curb infection in the body, effective against chronic cough and cold and gives good results for bronchitis and asthma.
Root	Tubers remove pain in joints, diabetes, skin lesions (Tenia), aphthous ulcers, wheezing and phlegm.
Decoction	Decoction of stem and leaf cures bronchitis.

Table 2. Qualitative analysis of phytochemicals in *Coccinia indica* fruit.

Phytochemical	Methanolic extract
Alkaloids	+
Steroids	+
Tannins	+
Saponins	+
Ellagic acid	+
Phenols	+
Glycosides	+
Lignans	-
Triterpenoids	+
Flavonoids	+

+ Indicates presence; - indicates absence.

distributed naturally in China, Tropical Asia, India, Australia and Africa. It is considered as a valuable wild vegetable by the indigenous people of Southeast Asia and India. Every part of this plant is valuable in medicine for ring worm, psoriasis, small pox, scabies (Perry, 1980) and other itchy skin eruptions and ulcers (Behl et al., 1993). *C. indica* has antidiabetic (Purintrapiban et al., 2006), hypoglycemic (Akhtar et al., 2007), anti-inflammatory (Rao et al., 2003; Juneja et al., 2007), analgesic (Rao et al., 2003), hepatoprotective (Rao et al., 2003), antioxidant (Venkateshwaran and Pari, 2003), antilithic (Jayaweera, 1980), and antimutagenic (Kusamran et al., 1998) activities.

The aim of the present study is to reveal the antimicrobial properties and phytochemicals of different extracts of *C. indica* fruit through *in vitro* investigation.

MATERIALS AND METHODS

Preparation of plant extracts

The plant material was collected during September – October, 2007 from different places of Warangal District, Andhra Pradesh, India and authenticated at Plant Systematics Laboratory, Department of Botany, Kakatiya University, Warangal. Coarse powder from the shade dried fruits of *C. indica* (50 g) was extracted to exhaustion successively with petroleum ether, diethyl ether, chloroform, ethyl acetate, acetone, methanol, ethanol and aqueous extracts using a soxhlet apparatus. These extracts thus obtained were dried under reduced pressure at room temperature not exceeding 40°C and were used for the assays. The solvents used were purchased from Merck (Mumbai) and all the solvents were of analytical grade and all the media ingredients were from Hi media (Mumbai).

Phytochemical screening

Chemical tests were carried out on the methanol extract to identify the constituents utilizing standard methods of analysis (Gibbs, 1974; Trease and Evans, 1989).

Bacterial cultures

Six bacteria, two gram positive, *Staphylococcus aureus*, *Bacillus cereus* and four gram negative, *Pseudomonas putida*, *Salmonella paratyphi A*, *S. paratyphi B* and *Klebsiella pneumoniae* were used for the bioassay. The pure strains were obtained from Microbial Type Culture Collection and Gene bank (MTCC), Institute of Microbial Technology, Chandigarh, India. The organisms were maintained on agar slopes at 4°C and subcultured for 24 h before use.

Bacterial susceptibility testing

The agar plate well diffusion method was used as described by

Table 3. Antibacterial activity of different solvent extracts of *Coccinia indica* fruit by well diffusion method (Zone of inhibition in mm).

Solvent (100 mg/ml)						
	Gram positive		Gram negative			
	<i>Staphylococcus aureus</i>	<i>Bacillus cereus</i>	<i>Pseudomona sputida</i>	<i>Salmonella paratyphi A</i>	<i>Salmonella paratyphi B</i>	<i>Klebsiella pneumoniae</i>
Petroleum ether	19 ± 0.2	12 ± 0.2	17 ± 0.1	9 ± 0.1	8 ± 0.2	15 ± 0.2
Diethyl ether	11 ± 0.1	10 ± 0.2	13 ± 0.1	1 ± 0.1	1 ± 0.2	9 ± 0.2
Chloroform	7 ± 0.2	1 ± 0.2	1 ± 0.2	-	2 ± 0.1	4 ± 0.1
Ethyl acetate	11 ± 0.2	10 ± 0.1	5 ± 0.1	-	1 ± 0.1	9 ± 0.1
Acetone	10 ± 0.1	18 ± 0.2	1 ± 0.1	-	-	6 ± 0.2
Methanol	8 ± 0.1	15 ± 0.1	13 ± 0.1	3 ± 0.1	9 ± 0.2	8 ± 0.1
Ethanol	11 ± 0.1	10 ± 0.2	11 ± 0.1	-	-	7 ± 0.1
Aqueous	13 ± 0.1	6 ± 0.2	2 ± 0.2	6 ± 0.1	1 ± 0.1	3 ± 0.1
Streptomycin (10 mg/ml)	16 ± 0.2	14 ± 0.1	15 ± 0.1	10 ± 0.1	12 ± 0.2	13 ± 0.2

- Indicates no inhibition.

Desta (2005). A standardized inoculum 1 to 2×10^7 cfu/ml 0.5 MC Farland standards was introduced onto the surface of sterile agar plate and evenly distributed the inoculum by using a sterile glass spreader. Simultaneously 8 mm wells were cut from the plate using a sterile cork borer.

100 µl of extract at a concentration of 100 mg/ml was introduced into each well. The agar plates were incubated aerobically at 37°C. After 24 h the inhibition zones were measured with a ruler and compared with the control well containing only solvent and 10 mg/ml of streptomycin served as control.

Statistical analysis

All the tests were conducted in triplicates. The data of all the parameters were statistically analysed and expressed as mean ± S.D.

RESULTS AND DISCUSSION

The profile of uses of different parts of *C. indica* is shown in Table 1. Phytochemical screening was carried out on the methanolic extract which revealed the presence of alkaloids, steroids, tannins, saponins, ellagic acids, phenols, glycosides and triterpenoids. Lignans were absent (Table 2). Some of the phytochemicals reported are known to possess various biological activities. Oral administration of water soluble alkaloid fraction reduced fasting blood glucose of guinea pig by 29.3 and 21% in OGTT (1 gm/kg) conducted on rats (Mukherjee et al., 1972). Triterpenoids like taraxerone have exhibited anti-inflammatory analgesic and hepatoprotective activity (Rao et al., 2003). Feeding of water soluble alkaloid fraction (1 gm/kg) of *C. indica* leaves to normal fasting guinea pigs showed hypoglycaemic activity of short duration and the effect was due to the presence of betasitosterol, a phytosterol (Mukherjee et al., 1972). Proteins like glucokenin have the property of reducing the amount of sugar in the blood (Sambasivam, 1931; Murugesu, 1988; Nadkarni, 2000).

Eight different solvent extracts of *C. indica* fruit were tested against six gram negative and gram positive bacteria. Results of antibacterial activity of all the extracts and their efficacies as compared to standard and depicted are shown in Table 3. The crude extracts of *C. indica* exhibited moderate to significant antibacterial activity against all tested bacteria with inhibition zone ranging from 1 - 19 mm and comparable to standard. Petroleum ether extract was the most active and showed considerable antibacterial activity against all tested gram positive and gram negative bacteria producing a maximum inhibition zone of 19 mm against *S. aureus*. Other tested extracts also inhibited the growth of a number of test organisms but to a lesser extent and were mainly active against the gram-positive *S. aureus*. The relatively high anti-staphylococcal activity might, however, be of interest since several *Staphylococcus* strains are reported to express a multi drug resistance and natural plant extracts were shown to have good anti-staphylococcal activity (Lechner et al., 2004). The study also revealed that methanol extract was found to be active against *B. cereus* and *P. putida* producing inhibition zone 15 and 13 mm, respectively. None of the extracts showed significant antibacterial activity against *S. paratyphi A* and *B*. So these bacterial strains are considered most resistant towards all the extracts tested. Least activity was shown by chloroform extract against all the tested bacteria with inhibition zone ranging from 1 - 7 mm.

Previous studies on antibacterial and antifungal activity of leaves and stem of *C. indica* (Hugo de Boer et al., 2005; Dewanjee et al., 2007; Umbreen et al., 2008) have also detected the significant activity of methanol and ethyl acetate extracts against different bacteria and fungi and providing support to the fact that methanol is a better solvent for extraction and isolation of phytochemicals having antimicrobial activity. The present study revealed the moderate activity of water extract agreeing with earlier reports that use of organic solvents is always better

for extraction and isolation of antibacterial compounds (Varadarajan et al., 2007). In addition, the effectiveness of the extracts was not due to one main constituent, but to the combined action of other chemical compounds involved in it (Bai, 1990). Some examples include alkaloids, flavonoids, terpenoids, thymol and other compounds of phenolic nature which are classified as antimicrobial compounds (Rojas et al., 1992). These findings on antibacterial activity support the claim of the traditional healers that *C. indica* has been used to relieve pneumonia, dysentery, cough and cold.

Even though the present study on these extracts is an addition to the scientific literature, detailed investigations are needed to isolate bioactive principles from these extracts.

REFERENCES

- Afia Akhtar M, Mamunur R, Mir Imam IW, Robiul IM, Sharif MS, Ariful IMd, Shah AMd, Maruf A (2007). Comparison of Long – term Antihyperglycemic and Hypolipidemic Effects between *Coccinia cordifolia* (Linn.) and *Catharanthus roseus* (Linn.) In Alloxan - induced diabetic Rats. Res. J. Med. Med. Sci. 2: 29-34.
- Ahmed I, Mehmood Z, Mohammad F (1998) . Screening of some Indian medicinal plants for antimicrobial properties. J. Ethnopharmacol. 62: 183-193.
- Bai D (1990). Traditional Chinese material: A respect and prospect. Plant Med. 50: 502.
- Behl PN, Arora RB, Srivastava G, Malhotia (1993). Herbs useful in Dermatological therapy, CBS Publishers and Distributor, Delhi.
- Davis J (1994). Inactivation of antibiotics and the dissemination of resistance genes. Science 264: 375-382.
- Desta B (2005). Ethiopian traditional herbal drugs part II. Antimicrobial activity of 63 medicinal plants. J. Ethnopharmacol. 100: 168-175.
- Dewanjee S, Kundu M, Maiti A, Majumdar R, Majumdar A, Mandal SC (2007). *In Vitro* evaluation of Antimicrobial Activity of Crude Extract from Plants *Diospyros peregrina*, *Coccinia grandis* and *Swietenia macrophylla* . Trop. J. Pharm. Res. 6 : 773-778.
- Gibbs RD (1974). Chemotaxonomy of flowering plants. McGill – Queen's university press, Montreal and London Vol I : 523-619.
- Hugo de Boer J, Anneleen Kool, Anders Broberg , William Mziray R, Inga Hedberg, Jolanta Levenfors J (2005). Anti-fungal and antibacterial activity of some herbal remedies from Tanzania. J. Ethnopharmacol. 96: 461-469.
- Indian Materia Medica by Dr. KM Nadkarni, Volume I, pages: 300 – 302, publisher: Bombay Popular Prakashan, reprinted, 2000.
- Jayaweera DM (1980). Medicinal Plants (Indigenous and Exotic) used in Ceylon. Part 2. A Publication of the Natural Sciences Council of Srilanka, Colombo.
- Juneja D, Shrivastava PN, Guha MK, Saxena RC (2007) . PHCOG MAG.: Short communication. Preliminary Phytochemical Screening of Some Folklore Medicinal Plants for their anti-inflammatory activity. Phcog. Mag. Issue 11 (Suppl.) Jul- Sep.pp. 201-203.
- Karaman I, Sahin P, Gulluce M, Oguten H, Songul M, Adiguzed A (2003). Antimicrobial activity of aqueous and methanol extracts of *Juniperus oxycedrus* L. J. Ethnopharmacol. 2837: 1-5.
- Kusamran WR, Anong T, Piengchai K (1998). Antimutagenic and anti carcinogenic potential of some Thai vegetables. Mutat. Res. 402: 247-258.
- Lechner D, Stavri M, Oluwatuyi M, Pereda–Miranda R, Gibbons S (2004). The anti – staphylococcal activity of *Angelica dahurica* (Bai Zhi) . Phytochemistry 65: 331-335.
- Materia Medica (Vegetable Section), Volume I, by Dr. Murugesu Muthaliar, pp. 332-333, publisher, Tamilnadu Siddha Medical Council, Chennai . Fourth edition 1988.
- Perry LM (1980) Medicinal Plants of East and South East Asia, Attributed properties and Uses, MIT Press, London.
- Purintrapiban J, Keawpradub N, Jansakul C (2006). Role of the water extract from *Coccinia indica* stem on the stimulation of glucose transport in L8 myotubes . Songklanakarin J. Sci. Technol. 28: 1199-1208.
- Rajasekharan PE, Ganeshan S (2002). Conservation of medicinal plant biodiversity in Indian perspective. J. Med. Aromatic Pl. Sci. 24: 132-147.
- Rao GMM, Vijayakumar M, Sreevidya N, Rao CV, Mehrotra S, Shirwaikar A (2003). Chemical and pharmacological investigation of the fruits of *Coccinia indica*. 2nd World Congress on Biotechnological Developments of Herbal Medicine, NBRI, Lucknow, UP, India. p. 131, February 20 – 22 .
- Rojas A, Hernandez L, Pereda–Mirands R, Meta R (1992) . Screening for antimicrobial activity of crude drug extracts and pure natural products from Mexican medicinal plants. J. Ethnopharmacol. 35: 275-283.
- Sashi KJ, Ramya M, Janardhan K (2003). Antimicrobial activity of ethnomedicinal plants of Nilgiri Biosphere reserve and Western Ghats. Asian J. Microbiol. Biotechnol. Environ. Sci. 5: 183-185 .
- Scultes RE (1978). The kingdom of plants In : W.A .R Thomson (ed.), Medicines from the earth, Mc Graw – Hill Book Co., New York, p. 208.
- Tamil – English Dictionary of Medicine, Chemistry, Botany and Allied Sciences by T.V. Sambasivam Pillai, Volume II and page: 1741, publisher : The Research Institute of Siddhar's Science, Mount Road, Madras. 1931.
- Trease GE, Evans WC (1989). A textbook of pharmacognosy. Balliere Tinnall Ltd.,London. p. 53.
- Umbreen F, Huma S, Shaikat M, Syed AA, Ghazala Rizwani H (2008). Antibacterial activities of *Coccinia grandis* L. Pak. J. Bot., 40: 1259-1262.
- Varadarajan M, Guruchandran V, Nagarajan SM, Natarajan E, Punitha R, Geetha C, Ramya S (2007) . Screening for the evaluation of antibacterial activity of *Solanum nigrum* L. . Asian J. Microbiol. Biotechnol. Sci. 9: 79-81.
- Venkateshwaran S, Pari L (2003). Effect of *Coccinia indica* leaves on antioxidant status in streptozotocin - induced diabetic rats .Journal of Ethnopharmacology 84: 163-168.