

Full Length Research Paper

Soil-transmitted helminthiasis among school age children in Ethiope East Local Government Area, Delta State, Nigeria

EGWUNYENGA, O. ANDY¹ and ATAIRU, D. PALMER²

¹Department of Zoology, Delta State University, Abraka, Nigeria.

²Parasitology Laboratory, Baptist Medical Centre, Eku, Delta State, Nigeria.

Accepted 23 August, 2005

The prevalence and intensity of soil-transmitted helminth infections and their relationship to anthropometric indices of 1,351 school children from Ethiope East Local Government Area of Delta State, Nigeria were evaluated. 739 (54.70%) of the subjects were infected by soil-transmitted helminths (STH). The overall prevalences by species were *Ascaris lumbricoides* (48.41%), hookworms (29.76%), and *Trichuris trichiura* (17.39%). 174 (12.88%) were infected with two or more STHs. Males (60.81%) were generally more infected than females (43.30%), but this was only statistically significant among children aged 5 – 7 years. The mean number of eggs per gram of faeces (epg) was generally low. Analysis of epg for each species indicated that 7.8, 7.60 and 1.70% of the subjects had high intensity of infections for *Ascaris*, hookworms and *Trichuris*, respectively. There was no relationship between intensity of infections and wasting, while children with high intensity of infections were more stunted than others.

Key words: Soil-transmitted helminths, school-age, children, nigeria, prevalence.

INTRODUCTION

Soil-transmitted helminthiasis remain an important cause of morbidity and sometimes mortality in developing tropical countries, particularly among paediatric age group (WHO, 1987). It is estimated that more than one billion people in the world are infected by soil-transmitted helminths (STH), mainly *Ascaris lumbricoides*, hookworms and *Trichuris trichiura* (Crompton, 1999). Although STHs affect all age-groups, the problem is predominant among the worlds estimated 400 million school children, and is often associated with poor growth, reduced physical activity, impaired cognitive function and learning ability (Stephenson et al 1998; Nokes et al., 1992; Adams et al., 1994; Koroma et al., 1994; Stoltz et al., 1996).

Effective control of STH infections depends on improvement in sanitation and living conditions, but implementation is usually hampered by lack of resources and political will. In the short term, school based de-worming has been recommended as a highly cost-effective public health measure in less developed countries (World Bank, 1993). The World Health Organization (WHO) also recommends a baseline survey in school children to determine the prevalence and intensity of infections (Montresor et al., 1998), and develop effective treatment strategies and case management options (Andrade et al., 2001). In addition, baseline surveys provide basis for development of control programmes at national, regional and district levels.

In Nigeria various school based baseline surveys have been carried out to estimate the current status of STH infections (Ogbe et al., 2002; Adeyeba and Akinlabi, 2002; Etim et al., 2002; Nock et al., 2003; Ukpa and Ugwu, 2003). This study was undertaken to add to the store of essential baseline data on the

*Corresponding author. E-mail: aoegwunyega@yahoo.com.

Table 1. Total prevalence of soil-transmitted helminth infections by age and sex.

| Age (years) | Male | | | Female | | | Overall | | |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | No. Examined | No. Infected | % Prevalence | No. Examined | No. Infected | % Prevalence | No. Examined | No. Infected | % Prevalence |
| 5 – 7 | 246 | 127 | 51.63 | 236 | 85 | 36.02 | 482 | 212 | 43.98 |
| 8 – 11 | 231 | 160 | 69.26 | 204 | 113 | 55.39 | 435 | 273 | 62.76 |
| 12 – 15 | 263 | 163 | 61.98 | 171 | 91 | 53.22 | 434 | 254 | 58.53 |
| Total | 740 | 450 | 60.81 | 611 | 289 | 47.30 | 1,351 | 739 | 54.70 |

prevalence and intensity of STH infections in Nigeria. The relationship between intensity of infections and the anthropometric indices of subjects in the study area is also examined.

MATERIALS AND METHODS

The study area

Ethiopia East Local Government Area of Delta State, Nigeria is made up of semi-urban settlements of about 100,000 inhabitants, located between latitude 5°N - 6°S and longitude 5.5°E - 6.5°W. The area is characterized by tropical climate with rainy season lasting from March to November. The vegetation ranges from mangrove thick forest to mixed rain forest and grasslands. The inhabitants of the area are mainly indigenous farmers and significant proportion of University students and public servants who reside among the majority indigenous population. Due to rainfall for much of the year (8-9 months) and human activity the soil is porous and moist with temperature range of 28-32°C which favour the survival of STH ova.

Subjects

The subjects were 1351 primary and secondary school children aged 5-15 years, randomly selected with the lottery method (Bens et al., 1991) from three schools from each of the six main towns in Ethiopia East Local Government Area, between February and November 2004. The towns and the number of subjects surveyed were Abraka (301), Eku (255), Ovu (198), Okpara (256), Isiokolo (194) and Kokori (147). Verbal consent was obtained from headmasters of the schools before stool containers were distributed to subjects.

Stool samples were collected in the morning and examined in the afternoon by Kato-katz method to quantitate the number of eggs per gramme of faeces (WHO, 1994). The examination of the stool samples was carried out at the Parasitology Laboratory of Baptist Medical Central Eku, Delta State. To ensure consistency of the readings, second readings were performed in 20% of the slides randomly selected (Andrasw et al., 2001). Intensity of infections for each worm was classified according to the thresholds proposed by the WHO Expert Committee (WHO, 1987).

The following data were collected for each student: weight, height, intensity of infections for *A. lumbricoides*, hookworms and *T. trichiura*. Height and weight measurements were compared to a standard population of the same age by use of Tanner's growth and weight charts (Tanner and Whitehouse, 1976). Children below the third percentile for weight and height were classified as malnourished (Andrade et al., 2001).

RESULTS

Of the total of 1351 stool samples examined, 739 (54.70%) were infected by soil-transmitted helminths (Table 1). Males (60.81%) were generally more infected than females (47.30%), but this was only statistically significant among children aged 5 – 7 years, with prevalence rates of 51.63% and 36.02% among males and females, respectively ($P < 0.05$). In the three age categories STH infections were highest in 8 – 11 years group in both males and females (Table 1).

The general prevalences of STH infections by species are given in Table 2. *A. lumbricoides* had the highest overall infection rate of 48.41%. Prevalence of *A. lumbricoides* was also the highest in each of the schools sampled. Overall prevalences of other STHs were hookworm 29.76% and *T. trichiura* 17.39%. In addition 174 (12.88%) of the subjects were infected by two or more soil-transmitted helminths. Of these 170 had double infections and 4 had triple infections. Schools in Kokori recorded the highest rates of infection with *A. lumbricoides* and hookworms (57.14 and 54.64%, respectively), while infection rates were highest among children from schools located in Eku and Abraka (17.25 and 14.29%, respectively).

The mean number of eggs per gramme of faeces (epg) indicated that the intensity of infections was generally low for the three helminths, following the thresholds proposed by WHO (Table 3). However analysis of epg by species showed that 7.8, 7.60 and 1.70% of the infected subjects had high intensity of infections for *A. lumbricoides*, hookworms and *T. trichiura*, respectively.

Table 4 shows the relationship between the intensity of infections and degree of malnutrition. Anthropometric measurements indicated that of the 1351 children examined, 241 (17.84%) were below the third percentile of weight (wasted) and 382 (28.28%) were below the third percentile of height (stunted). The data further reveal that there is no relationship between intensity of infection and wasting since among the uninfected group more underweight subjects (21.03%) were recorded than moderately infected subjects (12.5%). However children with high intensity of infections were more stunted than the other

Table 2. General prevalence of soil-transmitted helminthes by species among school-age children in the study area.

| School | n* | <i>A. lumbricoides</i> | | Hookworm | | <i>T. trichiura</i> | | Mixed infection | |
|--------------|--------------|------------------------|--------------|--------------|--------------|---------------------|--------------|-----------------|--------------|
| | | No. Infected | % Prevalence | No. infected | % Prevalence | No. infected | % Prevalence | No. Infected | % Prevalence |
| Abraka | 301 | 131 | 43.51 | 75 | 24.92 | 40 | 13.29 | 43 | 14.29 |
| Eku | 255 | 120 | 47.06 | 81 | 31.76 | 48 | 18.82 | 44 | 17.25 |
| Ovu | 198 | 98 | 49.49 | 60 | 30.30 | 36 | 18.18 | 28 | 14.14 |
| Okpara | 256 | 111 | 44.92 | 72 | 28.13 | 51 | 19.92 | 34 | 13.28 |
| Isiokolo | 194 | 106 | 54.64 | 80 | 41.24 | 31 | 15.98 | 13 | 6.70 |
| Kokori | 147 | 84 | 57.14 | 34 | 23.13 | 29 | 19.73 | 12 | 8.16 |
| Total | 1,351 | 654 | 48.41 | 402 | 29.76 | 235 | 17.39 | 174 | 12.88 |

* Number of school children examined in each town.

Table 3. Prevalence and intensity of infections.

| Worm | Overall Prevalence | Mean egg (\pm SE) | High Intensity |
|------------------------|--------------------|----------------------|--|
| <i>A. lumbricoides</i> | 48.41 | 3,244 \pm 261 | ⁵¹ / ₆₅₄ (7.8%) |
| Hookworm | 29.76 | 1,074 \pm 125 | ²¹ / ₂₇₄ (7.60%) |
| <i>T. trichinura</i> | 17.39 | 608 \pm 102 | ⁴ / ₂₃₅ (1.70%) |

children.

DISCUSSION

Data obtained from this study are consistent with a direct effect of lack of sanitation on prevalence and intensity of intestinal helminth infections. The outcome of this investigation has added to the store of knowledge on the occurrence of STHs in Nigeria. The overall prevalence recorded (54.7%) is in conformity with others studies from the Niger Delta part of Nigeria (Obiamiwe, 1977; Nwosu, 1981; Obiamiwe and Nmorsi, 1991; Udonsi, 1984; Ukpai and Ugwu, 2003). These studies have attributed the relatively high prevalence of STHs in children to poor environment and personal hygiene, shortage of potable water and indiscriminate defecation.

The majority of the infections found in our survey were caused by *A. lumbricoides*, followed by hookworms and *T. trichiura*. On the basis of the prevalence rates recorded in this study, the study area may be classified as high risk for STHs, hence targeted treatment of children with anthelmintic drugs should be regular. Earlier studies in Agbon (Ogbe et al., 2002), part of the study area, and many parts of Nigeria, have demonstrated the efficacy, acceptability and cost-effectiveness of school based control of soil-helminth infections (Nworgu et al., 1998).

Data on prevalence and intensity of infections (Table 3) show that with relatively high overall

prevalence of *A. lumbricoides* (about 50%), the mean number of eggs per gramme of faeces (epg) was low and majority of subjects were lightly infected. Only 7.8% of the 48.41% subjects infected with *A. lumbricoides* had high intensity of infection. This is consistent with the findings of Anderson and May (1991) who established that intestinal parasites are neither evenly nor randomly distributed among hosts, but tend to be aggregated in a few heavily infected individuals. Although such people are in the minority, in communities where the prevalence of infection is high, significant numbers are likely to be heavily infected and may experience morbidity.

WHO guidelines recommended periodic treatment rounds for groups with high intensity infections of 10% and above, regardless of the prevalence of overall infections. Since only 7.8% was recorded in this study, a different approach of treatment of only infected persons in school-based control programmes may be more cost-effective, given the semi-urban nature of the study area with relatively high population.

While about 30% of the children were found below the third percentile of height only 17.84% were below the third percentile of weight (Table 4). This could be explained by the fact that underweight represents a state of acute malnutrition that can be corrected by food, while stunting is an index of chronic malnutrition. Lack of adequate nutrients caused by high intensity infection in a critical period can prevent the normal growth spurt in pre-pubertal and

Table 4. Relationship between intensity of infections and degree of malnutrition.

| Parameter | Intensity of soil-transmitted helminthiasis | | | Uninfected Group (409) | Overall (1,351) |
|-------------------------------------|---|----------------|------------|---------------------------|--------------------|
| | High (81)* | Moderate (731) | Low (132) | | |
| < 3 rd centile of weight | 28(34.58%) | 92(12.59%) | 35(26.52%) | 86(21.03%) | 241(17.84%) |
| < 3 rd centile of height | 36(45.57%) | 201(27.50%) | 43(32.58%) | 102(24.94%) | 382(28.28%) |

*Number of subjects in each category.

pubertal children (Andrade et al., 2001). The children sampled in this survey live in environments where they are exposed from birth to intestinal parasites. Thus nutritional status of the children reflect not only previous episodes of acute and chronic infections, but also the adequacy or inadequacy of the diet to support satisfactory rates of growth (Hall, 1993). Each child may have a different history of infections and diet. Therefore cross-sectional studies as carried out in this study may not fully address the impact of intestinal worms on growth. Prospective studies of growth after treatment are recommended to fully elucidate the effect of intestinal worms on growth.

REFERENCES

- Adams EJ, Stephenson LS, Latham MC, Kinoti SN (1994). Physical activity and growth of Kenya school children with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infections are improved after treatment with Albendazole. *J. Nutr.* 124: 1199-1206.
- Adeyeba OA, Akinlabi AM (2002). Intestinal Parasitic infections among school children in a rural community, Southwest Nigeria. *Nig. J. Parasitol.* 23: 11-18.
- Anderson RM, May RM (1991). Infections diseases of humans. Oxford University Press. Trans. R. Soc. Trop. Med. 86 (4): 461.
- Andrade C, Alava TDE, Palacio IA, Del Poggio, P Jamoletti, C Gulletta M, Montresor A (2001). Prevalence and intensity of soil-transmitted Helminthiasis in the city of Portoviejo (Ecuador). *Mem. Inst. Oswaldo Cruz.* 96(8): 1075-1079.
- Ben S, Woods T, Liyonage WM, Smith DL (1991). A simplified general method for cluster sample surveys of health in developing countries *WHO Quarterly* 44: 98-106.
- Crompton DWT (1999). How much human helminthiasis is there in the world? *J. Parasitol.* 85: 397-403.
- Etim SE, Akpan PA, Abeshi SE, Effiom OE (2002). Intestinal helminth infections in children: Implications for helminth control using school based mass chemotherapy. *Nig. J. Parasitol.* 23: 53-60.
- Hall A (1993). Intestinal parasitic worms and the growth of children. *Transactions of the Royal Society of Trop. Med. Hygiene* 87: 241-242.
- Koroma MM, William AM, DE LA Haye, RR Hodges M (1996). Effects of Albendazole on growth of primary school children and the prevalence and intensity of soil-transmitted helminths in Sierr Leone. *J. Trop. Ped.* 42: 371-372.
- Montresor A, Crompton DWT, Hall A, Bundy DAP Savioli L (1988). Guidelines for the evaluation of soil-transmitted Helminthiasis and Schistosomiasis and Community Level. A Guide for Managers of Control Programmes. World Health Org. Geneva. pp. 1-45.
- Nock IH, Duniya D Galadima M (2003). Geohelminth eggs in soil and stool of pupils of some primary schools in Samaru, Zaria, Nigeria. *Nig. J. Parasitol.* 24:115-122.
- Nokes C, Grantham-McGregor SM, Sawyer AW, Cooper ES, Bundy DAP (1992). Parasitic helminth infection and cognitive function in school children. *Proc. R. Soc. Lond.* 247: 77-81.
- Nworgu OC, Okeibunor J, Madu E, Amazigo UO, Evans O (1998). Helminthiasis Control Programme in Nigeria. Acceptability to community members. *Trop. Med. Int. Health* 3(10): 841-849.
- Nwosu ABC (1981). The community ecology of soil-transmitted helminth infections of humans in a hyperendemic area of Southern Nigeria. *Annals. Trop. Med. Parasitol.* 75:175-203.
- Obiamiwe BA (1977). The pattern of parasitic infection in human gut at the Specialist Hospital Benin City, Nigeria. *Annals Trop. Med. Parasitol.* 7:35-43.
- Obiamiwe BA Nmorsi P (1991). Human gastro-intestinal parasites in Bendel State, Nigeria. *Angrew Parasitol.* 32: 177-183.
- Ogbe MG, Edet, Isichei MN (2002). Intestinal helminth infection in primary school children in areas of operation of Shell Petroleum Development Company of Nigeria (SPDC), Western Division in Delta State. *Nig. J. Parasitol.* 23: 3-10.
- Stephenson LS, Latham MC, Kinoti SN, Kurz KM, Brigham H (1998). Improvement in physical fitness of Kenya schoolboys infected with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* following a single dose of Albendazole. *Trans. R. Soc. Trop. Med. Hyg.* 84:277-282.
- Stoltzfus RJ, Albonico M, Chwaya HM, Savioli L, Tielsch J, Schulze K (1996). Hemoquqnt determination of hookworm – related blood loss and its role in iron deficiency in African children. *Am. J. Trop. Med. Hyg.* 55:399-404.
- Tanner JM, Whitehouse RA (1976). Height and growth charts. *Arch. Dis. Child.* 51:170-180.
- Udonsi JK (1984). *Necator americanus*: a cross-sectional study of rural community in relation to some clinical signs. *Annals of Trop. Med. Parasitol.* 78: 443-448.
- Ukpai OM, Ugwu CD (2003). The prevalence of gastro-intestinal tract parasites in primary school children in Ikwuano Local Government Area of Abia State, Nigeria. *Nig. J. Parasitol.* 24: 129-136.
- Ukpai OM, Ugwu CD (2003). The prevalence of gastrointestinal tract parasites in primary school children in Ikwuano Local Government Area of Abia State, Nigeria. *Nig. J. Parasitol.* 24: 129-136.
- World Health Organization (1987). Prevention and Control of Intestinal Helminth Infections. Report of WHO Expert Committee. Technical Rep. Series 749 WHO Geneva. p. 12.
- World Bank (1993). World Development Report. Investing in Health. Oxford University Press. New York.
- World Health Org. (1994). Bench Aids for the Diagnosis of Intestinal Parasites. Laboratory Manual, WHO, Geneva. p. 114.