

*Full Length Research Paper*

# The effect of CoQ<sub>10</sub> and vitamin E on serum total sialic acid, lipid-bound sialic acid, some trace elements and minerals in rats induced with doxorubicin

Gökhan Oto<sup>1\*</sup>, Suat Ekin<sup>2</sup>, Semih Yaşar<sup>3</sup>, Hülya Özdemir<sup>1</sup> and Fatmagül Yur<sup>4</sup>

<sup>1</sup>Department of Pharmacology, Medical Faculty, Yuzuncu Yil University, Van, Turkey.

<sup>2</sup>Department of Biochemistry, Faculty of Arts and Sciences, Yuzuncu Yil University, Van, Turkey.

<sup>3</sup>Department of Biochemistry, Faculty of Arts and Sciences, Bingöl University, Bingöl, Turkey.

<sup>4</sup>Department of Biochemistry, Faculty of Veterinary, Yuzuncu Yil University, Van, Turkey.

Accepted 28 April, 2011

**This study was designed to evaluate the effect of CoQ<sub>10</sub> and vitamin E on serum total sialic acid (TSA), lipid bound sialic acid (LSA) and some elements in rat administered doxorubicin (DXR). Cu levels were increased in the group treated with DXR + vitamin E in comparison with DXR (p<0.05) and CoQ<sub>10</sub> groups (p = 0.001). Furthermore, copper levels were increased in the group treated with DXR + CoQ<sub>10</sub> in comparison with CoQ<sub>10</sub> group (p < 0.05). Zn levels were decreased in the group treated with DXR + vitamin E in comparison with CoQ<sub>10</sub> group (p < 0.05). Mg levels were decreased in subjects treated with DXR + vitamin E in comparison with the control group values (p < 0.05). Particularly, the observed increase in Cu levels in rats from DXR + vitamin E group might be due to the decrease of vitamin E. However, the oxidative damage could be as a result of DXR occurrence and may be helpful to clinicians in chemotherapy using anthracycline.**

**Key words:** Doxorubicin, total sialic acid (TSA), lipid bound sialic acid (LSA), trace elements, minerals.

## INTRODUCTION

Doxorubicin (DXR) (also known as adriamycin or 14-hydroxydaunorubicin) is an anthracycline-type antitumor (antineoplastic or cytostatic) drug that is used in chemotherapy for treatment of various cancers (Arcamone et al., 1969). DXR is an effective anti-neoplastic agent, application of which has been limited due to its cardiotoxic side effects (Balaei et al., 2010; Wold et al., 2005). The mechanism of DXR-induced cardiomyopathy remains unclear, but most of the evidence indicates that free radicals accompanied by a decrease of endogenous antioxidants and the subsequent increase in oxidants results in enhanced oxidative stress leading to a slow loss of myofibrils and vacuolization of myocardial cells which are the typical changes in the DXR-induced heart failure (Singal and Iliskovic, 1998). Furthermore, myocardial damage may be caused by an increase in tissue calcium, the inhibition of nucleic acid protein synthesis,

lipid peroxidation, the release of vasoactive amines, TNF- $\alpha$  and interleukin-2, liberation of cytokine from the tumor, changes in adrenergic function, lysosomal alterations and the inhibition of the CoQ<sub>10</sub> and the sodium-potassium-activated ATPase (Simpson et al., 2004; Iarussi et al., 2001).

CoQ<sub>10</sub> (ubiquinone) is a mitochondrial coenzyme which is essential for the production of ATP (Littarru, 1994; Kumar et al., 2009). Being at the core of cellular energy processes, it assumes importance in cells with high energy requirements like the cardiac cells which are extremely sensitive to CoQ<sub>10</sub> deficiency produced by cardiac diseases. CoQ<sub>10</sub> has thus, a potential role in the prevention and treatment of heart ailments by improving cellular bioenergetics (Kumar et al., 2009). Furthermore, CoQ acts as a potent antioxidant in the inner mitochondrial membrane. It inhibits lipid peroxidation by either scavenging free radicals directly or reducing  $\alpha$ -tocopheroxyl radical to  $\alpha$ -tocopherol (Ernster and Dallner, 1995; Lass and Sohal, 1998).

$\alpha$ -Tocopherol is known to function *in vivo* as antioxidant

\*Corresponding author. E-mail: [gokhanoto@hotmail.com](mailto:gokhanoto@hotmail.com).

and to provide cell membranes with protection against peroxidative and hydroxyl radical damage (Halliwell, 1996). CoQ<sub>10</sub>H<sub>2</sub> acts as a chain-breaking radical scavenger (Kagan et al., 1990), synergizes with  $\alpha$ -tocopherol via reduction of the  $\alpha$ -tocopherol-derived phenoxyl radical (Kagan et al., 1990<sup>2</sup>) and thus, inhibit tocopherol-mediated peroxidation (Bowry and Stocker, 1993).

Serum sialic acid levels have been used as laboratory markers in a variety of pathological conditions (Kario and Matsuo, 1993; Hakamory, 1984). Sialic acid (SA), an important component of glycoproteins and glycolipids, is found in negatively charged surface polyanions on various cell membranes and plays an important role in the antigenic characterization of cells (Lehninger et al., 1993). Total sialic acid (TSA) levels are elevated in many circumstances, such as myocardial infarction (Güngör et al., 2004; Succari et al., 1982), cancer (Seyrek et al., 2005), some auto-immune disorders (Maneva et al., 1995), diabetes mellitus (Crook et al., 1993), infections and chronic glomerulonephritis (İşler et al., 1993).

Minerals and trace elements play a dual role in the biological system through their interaction with biomolecules (Strong and Garruto, 1994). In biological and medical research, the determination of the concentration of trace elements and minerals as well as their distribution pattern is extremely important since it could provide unique information about various physiological and patho-physiological processes in the investigated tissues (Seeley et al., 1996). Some trace elements, particularly copper and zinc are in fact implied in both humoral and cellular immunity (Berger et al., 1998). These essential micronutrients can interfere directly with the propagation stage of free radical generation and scavenge free radicals (Kulikowska and Moniuszko, 2004).

So far, in literatures, there is no report related to the effect of CoQ<sub>10</sub> and vitamin E on trace elements, sialic acid and lipid-bound sialic acid, despite the fact that several studies that are related to the effect of the other parameters on doxorubicin are available. The aim this study was to evaluate the effects of vitamin E and CoQ<sub>10</sub> treatment on serum total sialic acid, lipid-bound sialic acid, some trace element and mineral levels on DXR-induced cardiotoxicity in rat model.

## MATERIALS AND METHODS

### Animals and drug administration protocols

All procedures described were studied and approved by the Local Institutional Committee for the Ethical Use of Animals. Wistar albino rats (180 to 200 g) were used in these experiments. The rats were obtained from the Department of Virology, Firat University, Veterinary Research Institute. All animals were housed in standard cages at room temperature (22 ± 2°C), with artificial light from 7.00 am to 7.00 pm, and provided with standard pelleted food and water *ad libitum*.

Rats were divided into four groups (n = 6/group). In the first group, DXR with 2.5 mg/kg normal saline were applied intraperitoneally once a week for 6 weeks period. In the second group rats

in addition to DXR, CoQ<sub>10</sub> was applied intraperitoneally with the ratio of 4 mg/kg/live weight on daily bases. In the third group rats, only CoQ<sub>10</sub> was applied with the ratio of 4 mg/kg/live weight on daily bases. In the fourth group rats, in addition to DXR, vitamin E was applied with the ratio of 10 mg/kg/subcutaneous two times a week for 6 weeks. The study period were planned as 6 weeks.

### Serum samples preparation

Blood samples were collected at the beginning of application (0th) and at the 60th day of the application period by the intra cardiac method and centrifuged in 500 *g*-force (rpm) for 15 min to obtain sera which was later stored at -70°C until all experimental procedures were carried out.

### Serum analysis

Serum TSA and serum lipid bound sialic acid (LSA) levels were measured by colorimetric method using ATI UNICAM UV/V spectrometer (Katopodis et al., 1982; Sydow, 1985). Serum trace element (Zn, Cu and Fe) and mineral (Ca, Mg and Na) concentrations were measured using a solar atomic absorption (Thermo Electron Corporation, Solar House, Cambridge England) spectrophotometer.

### Statistical analysis

Data are presented as  $\bar{X} \pm \text{SEM}$  (standard error of mean). Differences in biochemical parameters were statistically evaluated using one-way analysis of variance (Anova) followed by Tukey multiple comparison test.

## RESULTS

No death was observed in any of the experimental groups. Average serum TSA, LSA, trace element (Fe, Cu and Zn) and mineral (Ca, Mg and Na) levels between groups are shown in Table 1 ( $\bar{X} \pm \text{SEM}$ ) and Figures 1 to 3.

### Serum levels of TSA and LSA

Data on the serum levels of TSA and LSA of the animals in the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups are summarized in Table 1 and Figure 1. The levels of TSA and LSA among the different treatment groups showed no difference as compared to the control group ( $p > 0.05$ ).


### Serum levels of trace elements

Data on the serum levels of trace elements (Fe, Cu and Zn) of the animals in the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups are summarized in Table 1 and Figure 2. The serum levels of Fe among the different treatment groups showed no difference when compared to the control group ( $P > 0.05$ ). The serum levels of Cu was increased in the DXR + vitamin E treated group

**Table 1.** Average concentrations of TSA, LSA, trace element (Fe, Cu and Zn) and mineral (Ca, Mg and Na) levels in serum samples of the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups.

Group	Control ( $\bar{X} \pm SEM$ )	DXR ( $\bar{X} \pm SEM$ )	CoQ <sub>10</sub> ( $\bar{X} \pm SEM$ )	DXR + CoQ <sub>10</sub> ( $\bar{X} \pm SEM$ )	DXR + Vit E ( $\bar{X} \pm SEM$ )
TSA (mg/dl)	80.48 ± 5.29	94.43 ± 7.21	79.32 ± 10.33	93.74 ± 6.19	86.02 ± 2.39
LSA (mg/dl)	21.65 ± 0.29	23.59 ± 0.40	20.83 ± 0.83	22.39 ± 0.85	21.83 ± 0.47
Fe (mg/L)	2.80 ± 0.34	2.61 ± 0.38	2.50 ± 0.17	1.71 ± 0.26	1.66 ± 0.43
Cu (mg/L)	1.19 ± 0.06	0.97 ± 0.10 <sup>b</sup>	0.75 ± 0.05 <sup>a,b1</sup>	1.21 ± 0.12 <sup>b1</sup>	1.44 ± 0.12 <sup>a,b</sup>
Zn (mg/L)	1.03 ± 0.07	0.93 ± 0.05	1.15 ± 0.03 <sup>b</sup>	0.84 ± 0.09	0.78 ± 0.23 <sup>b</sup>
Ca (mg/dl)	11.25 ± 0.83	9.17 ± 0.60	9.96 ± 0.67	9.19 ± 1.04	11.39 ± 0.57
Mg (mg/dl)	2.68 ± 0.14 <sup>b</sup>	2.44 ± 0.20	2.47 ± 0.03	2.64 ± 0.32	1.88 ± 0.09 <sup>b</sup>
Na (mg/ml)	2.88 ± 0.03	2.77 ± 0.02	2.86 ± 0.02	2.79 ± 0.03	2.86 ± 0.05

a,  $p = 0.001$ ; b,  $p < 0.01$ ; b<sub>1</sub>,  $p < 0.05$ .

**Figure 1.** Comparison of TSA and LSA levels in serum samples of the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups.

when compared to DXR group ( $P < 0.01$ ). Furthermore, it was increased in the DXR + vitamin E and DXR + CoQ<sub>10</sub> treated groups when compared to the CoQ<sub>10</sub> group ( $P = 0.001$  and  $p < 0.05$ , respectively). The serum levels of Zn was decreased in the DXR + vitamin E treated group as compared to the CoQ<sub>10</sub> group ( $P < 0.01$ ).


### Serum levels of minerals

Data on the serum levels of minerals (Ca, Mg and Na) of the animals in the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups are summarized in Table 1 and Figure 3. The serum levels of Ca and Na among the different treatment groups showed no difference as


compared to the control group ( $p > 0.05$ ). The serum levels of Mg was decreased in the DXR + vitamin E treated group as compared to the control group ( $p < 0.01$ ).

### DISCUSSION

Cytostatic agents are known to exert their antitumour action by interacting with specific cell structures or metabolic pathways of cancer cells (Sangeetha et al., 1990). Several studies have demonstrated the protective properties of various antioxidants against the harmful action of cytostatic drugs such as doxorubicin, and cisplatin *in vitro* (Miura et al., 1994). Experimental animal studies have shown that the co-administration of free radical



**Figure 2.** Comparison of trace element (Fe, Cu and Zn) levels in serum samples of the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups.



**Figure 3.** Comparison of mineral (Ca, Mg and Na) levels in serum samples of the control, DXR, CoQ<sub>10</sub>, DXR + CoQ<sub>10</sub> and DXR + vitamin E groups.

scavengers did not reduce the antitumour effect of agents such as doxorubicin (Keizer et al., 1990; Siveski-Iliskovic et al., 1995).

Cardiomyopathy and heart failure have been observed in DXR-treated cancer patients (Singal and Iliskovic,

1998). The cardiotoxic effect of DXR are mediated via different mechanisms (Wang et al., 2004). Doxorubicin induced heart failure is associated with an increase in oxidative stress (Lou et al., 2006). Assaying sialic acid may be of value for diagnostic purposes since sialic acid

content has been reported to be altered in cardiovascular disease (Lindberg et al., 1991). It has been reported that plasma sialic acid concentrations are elevated in coronary heart disease and myocardial infarction (Allain et al., 1996; Crook et al., 1994). Deepa and Varalakshmi (2005) showed that the mean plasma sialic acid concentration in DXR-treated rats was significantly higher than those of the other treatment groups. In this study, serum TSA and LSA levels in the DXR group were increased in comparison with the control group.

The human body possesses several molecules that protect healthy tissues from the damaging action of free radicals (Halliwell and Gutteridge, 1993). Some trace elements have an important role in free radical protection. The trace elements are incorporated in the structures of the proteins, enzymes and complex carbohydrates to participate in biochemical reaction. Trace elements with enzymes, for example, are necessary for the functioning and maintenance of the immune system (Bang et al., 2002). Most enzymes contain trace elements, such as selenium in glutathione peroxidase and zinc and manganese and copper in superoxide dismutase (Weijl et al., 1997; Coudray et al., 1992). Overexpression of cardiac-specific catalase prevents injury to the heart following oxidative insults such as exposure to the anticancer agent, doxorubicin (Kang et al., 1996). Cu deficiency can lead to reduction in catalase activity in tissues such as heart and liver (Strain, 1994). Chemical studies have shown that doxorubicin is a powerful iron chelator and the resultant iron-drug complex is an efficient catalyst of the conversion of hydrogen peroxide to the highly reactive hydroxyl radical (Ortega et al., 2001).

The levels of Zn is associated with levels of Cu and Fe due to the antagonistic relationships between these metals (DiSilvestro and Blostein-Fujii, 1997). It has been suggested that an imbalance between Cu and Zn may be a factor in the aetiology of cardiovascular diseases (Tiber et al., 1986). Costa and Nepomuceno (2006) reported that minerals such as Cu and Zn protects the organism against the genotoxic effects of chemotherapeutics such as DXR. Morishima et al. (1999) reported that DXR decreased plasma zinc levels in rat. We observed a significant increase in the levels of Cu following DXR + vitamin E treatment group as compared to DXR group. Furthermore, the serum levels of Zn was decreased in the DXR + vitamin E treated group as compared to CoQ<sub>10</sub> group.

Eckenhoff and Somlyo (1998) reported that the levels of mitochondrial Ca and Mg did not change in the DXR-injected animals, even in severely symptomatic rats 5 days after DXR administration. In this study, the serum levels of Mg decreased in the DXR + vitamin E treated group as compared to the control group.

The results of this study indicated that serum TSA and LSA levels in the DXR group were increased in comparison with the control group. The levels of TSA and LSA were lower in the CoQ<sub>10</sub> and vitamin E groups when

compared with the DXR group. Particularly, the observed increase in the Cu levels in rats from DXR + vitamin E group might be due to the decrease of vitamin E. However, the oxidative damage could be as a result of DXR occurrence and may be helpful to clinicians in chemotherapy using anthracycline.

## ACKNOWLEDGEMENT

This study was supported by a grant from the Scientific Research Projects Presidency of Yuzuncu Yil University (2007-SBE-YL104).

## REFERENCES

- Allain P, Oliver E, Le-Bouil A, Benoit C, Geslin P, Tadei A (1996). Increase of sialic acid concentration in the plasma of patients with coronary disease. *Presse Med.* 25: 96-98.
- Arcamone F, Cassinelli G, Fantini G, Grein A, Orezzi P, Pol C, Spalla C (1969). Adriamycin-14-hydroxydaunomycin, a new antitumor antibiotic from *Streptomyces peuceletius* var. *caesius*. *Biotechnol Bioeng.* 11: 1101-1110.
- Balaei MR, Momeny M, Babaeikeshomi R, Mehr SE, Tavangar SM, Dehpour AR (2010). The modulatory effect of lithium on doxorubicin-induced cardiotoxicity in rat. *Eur. J. Pharmacol.* Accepted 25 May, journal homepage: [www.elsevier.com/locate/ejphar](http://www.elsevier.com/locate/ejphar)
- Bang RL, Al-Bader AL, Sharma PN, Mattapallil AB, Behbehani AI, Dahti H (2002). Trace elements content in serum, normal skin, and scar tissues of keloid and normal scar patients, *The J. Trace Elem. Exp. Med.* 15: 57-66.
- Berger MM, Spertini F, Shenkin A, Wardle C, Wiesner L, Schindler C, Chioloro RL. (1998). Trace element supplementation modulates pulmonary infection rates after major burns: a double-blind, placebo-controlled trial. *Am. J. Clin. Nutr.* 68: 365-71.
- Bowry VW, Stocker R (1993). Tocopherol-mediated peroxidation. The pro-oxidant effect of vitamin E on the radical-initiated oxidation of human low-density lipoprotein. *J. Am. Chem. Soc.* 115: 6029-44.
- Coudray C, Richard MJ, Laporte F, Faure P, Roussel AM, Favier A (1992). Superoxide dismutase activity and zinc status: A study in rats and man. *J. Nutr. Med.* 3: 13-26.
- Costa WF, Nepomuceno JC (2006) Protective effects of a mixture of antioxidant vitamins and minerals on the genotoxicity of doxorubicin in somatic cells of *Drosophila melanogaster*. *Environ. Mol. Mutagen* 47: 18-24.
- Crook M, Haq M, Haq S, Tutt P (1994). Plasma sialic acid and acute phase proteins in patients with myocardial infarction. *Angiology*, 45: 709-715.
- Crook MA, Tutt P, Simpson H, Pickup JC (1993). Serum sialic acid and acute phase proteins in type 1 and type 2 diabetes mellitus. *Clin. Chim. Acta.* 219: 131-138.
- Deepa PR, Varalakshmi P (2005). Biochemical evaluation of the inflammatory changes in cardiac, hepatic and renal tissues of adriamycin-administered rats and the modulatory role of exogenous heparin-derivative treatment. *Chemico-Biological Interactions* 156(2-3): 93-100.
- DiSilvestro RA, Blostein-Fujii A (1997). Moderate zinc deficiency in rats enhances lipoprotein oxidation in vitro. *Free Rad. Biol. Med.* 22: 739-742.
- Eckenhoff RG, Somlyo AP (1998). Cardiac mitochondrial calcium content during fatal doxorubicin toxicity. *Toxicol. Appl. Pharmacol.* 97: 167-172.
- Ernster L, Dallner G (1995). Biochemical, physiological and medical aspects of ubiquinone function. *Biochim. Biophys. Acta* 1271: 195-204.
- Güngör Ö, Sunar B, Özçelik F, Aktaş Z, Gökmen SS (2004). Serum sialic acid levels in acute myocardial infarction and its relationship to

- ceruloplasmin. *Turk. J. Biochem.* 29: 226-231.
- Hakamory S (1984). Tumor-associated carbohydrate antigens. *Ann. Rev. Immunol.* 2: 103-126.
- Halliwell B, Gutteridge JMC (1993). *Free radicals in Biology and Medicine*, 2nd ed. Oxford: Clarendon Press, pp. 86-187.
- Halliwell B (1996). Antioxidants in human health and disease. *Annu. Rev. Nutr.* 16: 33-50.
- Iarussi D, Indolfi P, Casale F, Coppolino P, Tedesco MA, DiTullio M (2001). Recent advances in the prevention of anthracycline cardiotoxicity in childhood. *Curr. Med. Chem.* 13: 1649-60.
- İşler M, Güner İ, Yılmaz H, Bayındır O, Önder R, Ve Nalbantgil İ (1993). Changes of serum total sialic acid in acute myocardial infarction. *Turk. J. Cardiol.* pp. 262-265.
- Kagan V, Serbinova E, Packer L (1990). Antioxidant effects of ubiquinones in microsomes and mitochondria are mediated by tocopherol recycling. *Biochem. Biophys. Res. Commun.* 169: 851-857.
- Kagan VE, Serbinova EA, Koynova GM, Kitanova SA, Tyurin VA, Stoytchev TS, Quinn PJ, Packer L (1990). Antioxidant action of ubiquinol homologues with different isoprenoid chain length in biomembranes. *Free Radic. Biol. Med.* 9: 117-126.
- Kang YJ, Chen Y, Epstein PN (1996). Suppression of doxorubicin cardiotoxicity by overexpression of catalase in the heart of transgenic mice. *J. Biol. Chem.* 271: 12610-12616.
- Kario K, Matsuo T (1993). Relation between sialic acid concentrations and the haemostatic system in the elderly. *BMJ.* 306: 1650-1651.
- Katopodis N, Hirshaut Y, Geller NL, Stock CC (1982). Lipid-associated sialic acid test for the detection of human cancer. *Cancer Res.* 42: 5270-5275.
- Keizer HG, Pinedo HM, Schuurhuis GJ, Joenje H (1990). Doxorubicin (adriamycin): a critical review of free radical-dependent mechanisms of cytotoxicity. *Pharmacol. Ther.* 47: 219-231.
- Kulikowska KE, Moniuszko JJ (2004). *Polish J. Environ. Stud.* 13(2): 5-13.
- Kumar A, Kaur H, Devi P, Mohan V (2009). Role of coenzyme Q10 (CoQ10) in cardiac disease, hypertension and Meniere-like syndrome. *Pharmacol. Ther.* 124: 259-268.
- Lass A, Sohal RS (1998). Electron transport-linked ubiquinone-dependent recycling of  $\alpha$ -tocopherol inhibits autoxidation of mitochondrial membranes. *Arch. Biochem. Biophys.* 352: 229-236.
- Lehninger AL, Nelson DL, Cox MM (1993). *Principles of Biochemistry* (2nd ed), Worth Publishers New York, pp. 298-324.
- Lindberg G, Eklund Ga, Gullberg B, Rastam L (1991). Serum sialic acid concentration and cardiovascular mortality. *Br Med J.* 302: 143-146.
- Littarru GP (1994). Energy and Defense. Facts and perspectives on CoenzymeQ<sub>10</sub> in biology and medicine. Casa Editrice Scientifica Internazionale, pp. 81-91.
- Lou H, Kaur K, Sharma AK, Singal PK (2006). Adriamycin-induced oxidative stress, activation of MAP kinases and apoptosis in isolated cardiomyocytes. *Pathophysiology*, 13: 103-109
- Maneva A, Michailova D, Sarbenova M, Christozova I, Taleva B. (1995). Superoxide dismutase and lipid-bound sialic acid in sera from children with cancers and juvenile chronic arthritis. *Eur. J. Cancer Prev.* 4: 429-435.
- Miura T, Muraoka S, Ogiso T (1994). Effect of ascorbate on adriamycin-Fe<sup>3+</sup>-induced lipid peroxidation and DNA damage. *Pharmacol. Toxicol.* 74: 89-94.
- Morishima I, Okumura K, Matsui H, Kaneko S, Numaguchi Y, Kawakami K, Mokuno S, Hayakawa M, Toki Y, Ito T, Hayakawa T (1999). Zinc accumulation in adriamycin-induced cardiomyopathy in rats: Effects of melatonin, a cardioprotective antioxidant. *J. Pineal Res.* 26: 204-210.
- Ortega R, Deves G, Bohic S, Simionovici A, Menez B, Bonnin-Mosbah M (2001). Iron distribution in cancer cells following doxorubicin exposure using proton and X-ray synchrotron radiation microprobes. *Nuclear Instruments and Methods in Phys. Res. B.* 181: 480-484.
- Sangeetha P, Das UN, Koratkar R, Suryaprabha P (1990). Increase in free radical generation and lipid peroxidation following chemotherapy in patients with cancer. *Free Radic. Biol. Med.* 8: 15-19.
- Seeley RR, Trent DA, Philip DA (1996). *Essentials of Anatomy and Physiology. The digestive system* 2nd ed. (Mc Graw Hill Pub), pp. 444-457.
- Seyrek K, Seyrek-Intas K, Keskin A, Kargin-Kiral F, Musal B, Toplu N (2005). Biochemical and histochemical studies on the occurrence of sialic acids in mammary tumours of bitches. *Res. Med. Vet-Toulouse*, 156: 258-263.
- Simpson C, Herr H, Courville KA (2004). Concurrent therapies that protect against doxorubicin-induced cardiomyopathy. *Clin. J. Oncol. Nurs.* 8: 497-501.
- Singal PK, Ilikskovic N (1998). Doxorubicin-induced cardiomyopathy. *N. Engl. J. Med.* 339: 900-905.
- Siveski-Ilikskovic N, Hill M, Chow DA, Singal PK (1995). Probuocol protects against adriamycin cardiomyopathy without interfering with its antitumor effect. *Circulation*, 91: 10-15.
- Strain JJ (1994). Newer aspects of micronutrients in chronic disease: copper. *Proc. Nutr. Soc.* 53: 583-598.
- Strong MJ, Garruto RM (1994). Experimental paradigms of motor neuron degeneration. In: Woodruff ML, Nonneman AJ, editors. *Toxin-induced Models of Neurological Disorders*. New York: Plenum Press: pp. 39-88.
- Succari M, Foglietti MJ, Percheron I (1982). Perchlorosoluble glycoproteins and myocardial infarct: modification of the carbohydrate moiety. *Pathol. Biol. Paris*, 30: 151-154.
- Sydow GA (1985). Simplified quick method for determination of sialic acid in serum. *Biomed. Biochem. Acta.* 44: 1721-1723.
- Tiber AM, Sakhaii M, Joffe CD and Ratnaparkhi MV (1986). Relative values of plasma copper, zinc, lipids and lipoproteins as markers for coronary artery disease. *Atherosclerosis*, 62: 105-110.
- Wang SW, Konorev EA, Kotamraju S, Joseph J, Kalivendi S, Kalyanaraman B (2004). Doxorubicin induces apoptosis in normal and tumor cells via distinctly different mechanisms-intermediacy of h2o2- and p53-dependent pathways. *J. Biol. Chem.* 279: 25535-45543.
- Weijl NI, Cleton FJ, Osanto S (1997). Free radicals and antioxidants in chemotherapy induced toxicity. *Cancer Treatment Rev.* 23: 209-40.
- Wold LE, Aberle NS, Ren J (2005). Doxorubicin induces cardiomyocyte dysfunction via a p38 MAP kinase-dependent oxidative stress mechanism cancer detection and prevention 29: 294-299.