

## Review

# Microsatellite markers: An important fingerprinting tool for characterization of crop plants

Siddra Ijaz

Centre of Agricultural Biochemistry and Biotechnology (CABB), University of Agriculture, Faisalabad, Pakistan. E-mail: [siddraijazkhan@yahoo.com](mailto:siddraijazkhan@yahoo.com). Tel: ++ 412630136.

Accepted 21 December, 2010

**Microsatellites are simple sequence repeats (SSR) of 1-6 nucleotides. They appear to be ubiquitous in higher organisms, both in animal and plant genomes and involving repetitive as well as unique sequences, although the frequency of microsatellites varies between species. They are abundant, dispersed throughout the genome and show higher levels of polymorphism than do other genetic markers. These features coupled with their ease of detection have made them useful markers. Their potential for automation and their inheritance in a co-dominant manner are additional advantages when compared with other types of molecular markers. SSRs are highly polymorphic, genome specific, abundant and co-dominant, and have recently become important genetic markers in cereals including wheat and barley.**

**Key words:** Simple sequence repeat, allelic diversity, polymorphism, co-dominance, microsatellites, genetic diversity.

## INTRODUCTION

With the advances in understanding the sequence organization of genomic deoxyribonucleic acid (DNA), a class of repetitive DNA sequences was found to be an integral part of the eukaryotic genome. Genomic simple sequence repeats (SSRs) have been used for a variety of purposes, including gene tagging, physical mapping, genome mapping and estimation of genetic diversity. SSRs/microsatellites are short sequence elements that are arranged in a simple internal repeat structure. SSR or short tandem repeats of DNA sequences (1-6 bp), is present throughout the genome of an individual, both in coding and non-coding regions. Microsatellite loci display co-dominant inheritance, high information content, reproducibility, and even distribution along chromosomes with specificity of locus (Kashi et al., 1997; Roder et al., 1998).

Microsatellite markers could characterize and discriminate all genotypes (Ijaz and Khan, 2009). The

frequency of SSR is higher in transcribed regions, especially in the untranslated portions, than in non-transcribed ones (Morgante et al., 2002). SSRs, densely and randomly distributed throughout the whole eukaryotic genome (Tautz, 1989), are tandemly repeated sets of 2 to 8 base pairs, and can vary extensively in the number of repeats, which may be due to slippage of DNA polymerase during replication and unequal crossing over (Schlotterer and Tautz, 1992).

A possible problem, which is associated with the use of simple sequence repeat markers/microsatellite markers, is the occurrence of null alleles (Callen et al., 1993). In comparison with the high polymorphism of genomic SSRs, expressed sequence tag (EST)-SSRs are less powerful in providing information on genetic variations (Eujayl et al., 2002; Thiel et al., 2003).

## MICROSATELLITE MARKERS SHOW GREAT ALLELIC DIVERSITY AND POLYMORPHISM

SSR markers are highly mutable loci, present in the eukaryotic genome at several sites, having stretches of tandemly repeated nucleotide motifs that can be as short

---

**Abbreviations:** SSR, simple sequence repeats; DNA, deoxyribonucleic acid; EST, expressed sequence tag; PCR, polymerase chain reaction; PIC, polymorphism information content; RFLP, restriction fragment length polymorphisms.

as 4, 3, 2 or 1 nucleotide. Knowledge of the sequence of these regions are used for designing specific amplifying primers, which then define sequence-tagged microsatellite sites (STMS). SSR markers/microsatellite markers are robust tools that can be exchanged between laboratories, and they are highly polymorphic (Thomas and Scott, 1993). Microsatellites and SSRs are multiallelic and hypervariable, and due to these characteristics, SSR have proven to be highly informative in studies of evolutionary relationships and genetics (Buchanan et al., 1994). Reproducibility of SSR can be efficiently utilized by different research laboratories to generate consistent data (Saghai-Marroof et al., 1994).

SSR markers are those molecular markers that prove to be useful tools for genome mapping, population and parentage analysis, individual identification, phylogenetic studies, cancer diagnostics, etc. The methods for the detection of simple sequence repeats are very quick, easy and feasible for these studies (Lefort et al., 1999). These are polymerase chain reaction (PCR)-based molecular markers and are efficient, reliable, cost-effective and common in rice. Microsatellite markers show a high degree of polymorphism in rice compared to restriction fragment length polymorphisms (RFLPs) and SSR markers are suitable molecular markers for evaluating genetic diversity among closely related rice cultivars (Temnykh et al., 2000). Microsatellite makers are easy to apply once they are developed, as the methodology for their development is complex, difficult and very costly, which are the reasons for restricting their application to important crops such as sugarcane (Scott et al., 2000).

By searching ESTs in sugarcane, 402 SSRs and various dinucleotide and trinucleotide SSRs have been found, and in saccharum species, EST-derived SSRs are useful because they are transferable and polymorphic (da Silva, 2001). Microsatellite markers have greater allelic diversity than other molecular markers (McCouch et al., 2001).

During polyploidization, gene conversion, inter-chromosomal exchange, replication slippages are involved in the evolution of microsatellite/SSR loci (Sayed et al., 2001). Microsatellite markers may be detected in the eukaryotic genome, because interspersed and simple repeats may well overlap regions transcribed by ribonucleic acid (RNA) polymerase, including ESTs in humans, tandem repeat polymorphisms in human genes being more common than generally believed with about 8% of such loci being within the coding sequence and if polymorphic, resulting in a frame shift (Wren et al., 2001).

EST-SSRs belong to that part of the genome that is a transcribed region and may be relatively well conserved, so that any polymorphism that is found and identified using EST-SSRs, may reflect a better relationship between varieties or species. Evaluation of the germplasm with SSRs derived from ESTs enhances or boosts the role of genetic or molecular markers by assaying variations in genes that are transcribed and of known function (Eujayl

et al., 2002). For application in marker assisted selection (MAS), microsatellite markers with high polymorphism information content (PIC) values were found to be more useful. The microsatellites derived from genomic sequences were found to be more successful than those SSR markers that were designed from EST (Karakousis et al., 2003). EST-SSR markers have been employed to screen different species of wheat grass, which indicate further chromosome rearrangements or variations in sequences between wheat grass genomes (Mullan et al., 2005), and SSR variations have been identified in 60 durum wheat accessions (Wang et al., 2007).

## UTILIZATION OF MICROSATELLITE MARKERS IN BREEDING PROGRAMS

The efficiency of a breeding program is dependent on the diversity of parents that are used in crosses. With the help of SSRs, diversity can be easily found or detected, that exhibits a higher level of polymorphism compared with other isozymes (Trujillo et al., 1995; Belaj, 1998). For microsatellites, heterozygosity values are 7-10 times more than RFLP. SSRs detected heterozygosity among cultivars in rice (McCouch et al., 1988; Wang and Tanksley, 1989; Virk et al., 1995) and in other crops (Kresovich et al., 1994). In plants, SSR markers have proven to be valuable tools for DNA genotyping, population genetics, genetic mapping studies, conservation and the management of genetic resources (Thomas and Scott, 1993; Bowcock et al., 1994; Jarne and Lagoda, 1996; Zane et al., 2002).

Microsatellite/SSR markers are markers of choice due to a high level of polymorphism, as well as higher reliability with the highest polymorphic content (PIC). These characteristics of microsatellites augmented the utilization of these markers as molecular markers in fingerprinting (Weising et al., 1995; Diwan and Cregan, 1997; Ashikawa et al., 1999). These markers have been widely used in soyabean due to their high level of sequence polymorphism, repeatability, co-dominance, rapid and relatively inexpensive use (Maughan et al., 1995; Mudge et al., 1997). Among various SSR loci, there is a significant variation in allelic diversity (Akagi et al., 1997; McCouch et al., 2001; Ravi et al., 2003). SSR identification in ESTs was reported for a wide range of species (Smulders et al., 1997; Cho et al., 2000; Haung et al., 2000; Lee et al., 2004). The presence of extra microsatellite/SSR motifs demonstrates that, as found in other plant genomes, there are many compound SSRs, suggesting that there may be clustering of SSRs in some genomic regions (Fisher et al., 1996). Those polymorphisms that are based on variation in the number of microsatellites have been successfully applied to plant breeding programs. Primers have been studied that are obtained from the sequence flanking the repeat unit; they are used with genomic DNA in PCR reactions and show SSR

polymorphisms that represent different alleles at that locus (Gupta et al., 1996).

Inter-microsatellite amplification (IMA) markers in the F2 population of peach, for economically important traits, including the four agronomic characters peach/nectarine, flat/round fruit, acid/non-acid fruit, and pollen sterility, have been studied and mapped to 11 linkage groups. Between pairs of markers, the average density is 4.5 cM (Dirlewanger et al., 1998). The genetic attributes of SSRs or microsatellites as DNA markers, coupled with their presence in most, if not all, sugarcane genotypes make SSR/microsatellites the preferred and reliable method for use in the construction of a reliable framework of the genetic map of sugarcane (Cordeiro et al., 1999). These markers are abundant, genome specific, co-dominant, and cover all 21 wheat chromosomes and have been successfully employed to characterize genetic diversity in seed bank collections of improved wheat germplasm (Borner et al., 2000; Huang et al., 2002) and wild relatives (Hammer, 2000). Morphologically, *Triticum urartu* and *Triticum baeticum* are very similar. It was possible to distinguish between these two species by using these markers. Therefore, SSR markers are a new powerful tool that supports the determination of critical races in diploid wild wheat species. They also allow the discussion of evolutionary pathways within *Triticum* (Hammer, 2000). So, for efficient screening of the germplasm by saturating more regions of wheat genome, more polymorphic wheat microsatellites could be used (Ijaz and Khan, 2009).

In sunflower, 980 clones from a genomic DNA library enriched for several dinucleotide and tetranucleotide SSR motifs were identified and then sequenced, of these, 35.5% were unique sequences, 71% of the amplified products displayed polymorphisms among 16 elite inbred lines; primers sets were designed for 87% of the single clones, and 74.5% of the primer sets amplified clean SSRs for genomic DNA (Yu et al., 2000). Resistance to green bug biotype E, and to map the resistance quantitative trait loci (QTLs) by using a microsatellite linkage map, it was studied and found that microsatellite markers are closely linked to the major resistance QTLs (YQ et al., 2007). The presence of locus in OS1E6 locus in various cereal genotypes have been found by analyzing these cereal genotypes. The presence of this locus in various cereal genotypes indicates its conservation across different cereal members (Daviewala et al., 2000). Microsatellite markers or SSR markers linked to the gene Pm 24 in an F2 progeny in wheat by applying bulk segregation analysis (BSA) have been identified (Huang et al., 2000). Three microsatellite markers have been mapped, namely Xgwm 106, Xgwm 337 and Xgwm 458, in the coupling phase to the Pm 24 locus. An allele of microsatellite locus Xgwm 337 was located 2.4 + -1.2 cM from Pm 24.M.

## CONCLUSION

In conclusion, microsatellite markers/SSR markers show

co-dominant expression and multiallelism, and because of these characteristics, simple sequence repeat markers/microsatellite markers exhibit high PIC values (Ferreira and Grattapaglia, 1998). These markers display high gene diversity scores, and these characteristics make them useful in distinguishing closely related genotypes (Giancola, 1998). Hence, SSRs have been found to be highly polymorphic, genome specific, abundant and co-dominant, and have recently become important genetic markers in cereals including wheat and barley.

## REFERENCES

- Akagi H, Yokozeki Y, Inagaki A, Fujimura T (1997). Highly polymorphic microsatellites of rice consist of AT repeats, and a classification of closely related cultivars with these microsatellite loci. *Theor. Appl. Genet.* 94: 61-67.
- Ashikawa I, Kurata N, Saji S, Umehara Y, Sasaki T (1999). Application of restriction fragment fingerprinting with a rice microsatellite sequence to assembling rice YAC clones. *Genome*, 42: 330-337.
- Belaj A (1998). Identificación y clasificación de variedades de olivo (*Olea europaea* L.) del Banco de Germoplasma de Cordoba con marcadores RAPDs. MSc. Universidad de Cordoba, Spain.
- Borner A, Chebotar S, Korzun V (2000). Molecular characterization of the genetic integrity of wheat (*Triticum aestivum* L.) germplasm after long-term maintenance. *Theor. Appl. Genet.* 100: 494-497.
- Bowcock AM, Ruizlinares A, Tomfohrde J, Minch E, Kidd JR, Cavallisforza LL (1994). High-resolution of human evolutionary trees with polymorphic microsatellites. *Nature (Lond.)*, 368: 455-457.
- Buchanan FC, Adams LJ, Littlejohn R.P, Maddox JF, Crawford AM (1994). Determination of evolutionary relationships among sheep breeds using microsatellites. *Genomics*, 22: 397-403.
- Callen DF, Thompson AD, Shen Y, Phillips HA, Richards RL, Mulley JC, Sutherland GR (1993). Incidence and origin of null alleles in the (AC)<sub>n</sub> microsatellite markers. *Am. J. Hum. Genet.* 52: 922-927.
- Cho YG, Ishii T, Temnykh S, Chen X, Lipovich L, McCouch SR, Park WD, Ayres N, Cartinhour S (2000). Diversity of microsatellites derived from genomic libraries and GenBank sequences in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 100: 713-722.
- Cordeiro GM, Maguire TL, Edwards KJ, Henry JR (1999). Optimization of Microsatellite Enrichment Technique in *Saccharum* spp. *Plant Mol. Biol. Reporter*, 17: 225-229.
- Da Silva JAG (2001) Preliminary analysis of microsatellite markers derived from sugarcane expressed sequence tags (ESTs). *Genet. Mol. Biol.* 24: 1-4.
- Daviewala AP, Ramakrishna W, Ranjekar PK, Gupta VS (2000). Sequence variations at a complex microsatellite locus in rice and its conservation in cereals. *Theor. Appl. Genet.* 101(8): 1291-1298.
- Dirlewanger E, Pronier V, Parvery C, Rothan C, Guye A, Monet R (1998). Genetic linkage map of peach (*Prunus persica* L. Batsch) using morphological and molecular markers. *Theor. Appl. Genet.* 97(5-6): 888-895.
- Diwan N, Cregan PB (1997). Automated sizing of fluorescent-labeled simple sequence repeat (SSR) markers to assay genetic variation in soybean. *Theor. Appl. Genet.* 95: 723-733.
- Eujayl I, Sorrells M, Baum M, Wolters P, Powell W (2002). Isolation of EST-derived microsatellite markers for genotyping the A and B genomes of wheat. *Theor Appl Genet* 104: 399-407.
- Ferreira ME, Grattapaglia D (1998). Introdução ao Uso de Marcadores Moleculares em Análise Genética. 3. ed. Embrapa, Brasília: p. 220.
- Fisher PJ, Gardner RC, Richardson TE (1996). Single locus microsatellites isolated using 5'-anchored PCR. *Nucleic Acids. Res.* 24: 4369-4371.
- Giancola SM (1998). Caracterización y diferenciación de germoplasma de soja (*Glycyne max* L. Merr.) mediante marcadores moleculares. PhD. Thesis. Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina.
- Gupta PK, Balyan HS, Sharma PC, Ramesh B (1996). Microsatellites in

- Plants: A new class of molecular markers. *Curr. Sci.* 70(1): 45-54.
- Hammer K (2000). Microsatellite markers-A new tool for distinguishing diploid wheat species. *Genet. Resour. Crop Evol.* 47: 497-505.
- Huang XQ, Hsam SLK, Zeller FJ, Wenzel G, Mohler V (2000). Molecular mapping of the wheat powdery mildew resistance gene Pm24 and marker validation for molecular breeding. *Theor. Appl. Genet.* 101(3): 407-414.
- Huang XQ, Bo Arner MS, Roder, Ganai MW (2002). Assessing genetic diversity of wheat (*Triticum aestivum* L.) germplasm using microsatellite markers. *Theor. Appl. Genet.* 105: 699-707.
- Ijaz S, Khan IA (2009). Molecular characterization of wheat germplasm using microsatellite markers. *Genet. Mol. Res.* 8(3): 809-815.
- Jarne P, Lagoda JL (1996). Microsatellites, from molecules to populations and back. *Trends Ecol. Evol.* 11: 424-429.
- Karakousis A, Barr RA, Chalmers JK, Ablett AG, Holton AT, Henry JR, Lim P, Langridge P (2003). Potential of SSR markers for plant breeding and variety identification in Australian barley germplasm. *Aust. J. Agric. Res.* 54(12): 1197-1210.
- Kashi Y, King D, Soller M (1997). Simple sequence repeats as a source of quantitative genetic variation. *Trends Genet.* 13: 74-78.
- Kresovich S, Lamboy WF, Li R, Ren J, Szewc-McFadden AK, Blick SM (1994). Application of molecular methods and statistical analysis for discrimination of accessions and clones of vetiver grass. *Crop Sci.* 34: 804-809.
- Lee JM, Nahm SH, Kim YM, Kim BD (2004). Characterization and molecular genetic mapping of microsatellite loci in pepper. *Theor. Appl. Genet.* 108: 619-627.
- Lefort F, Douglas GC (1999). An efficient micromethod of DNA isolation from mature leaves of four hardwood tree species *Acer*, *Fraxinus*, *Prunus* and *Quercus*. *Ann. Sci. For.* 56: 259-263.
- Maughan PJ, Saghai-Marooif MA, Buss GR (1995). Microsatellite and amplified sequence length polymorphisms in cultivated and wild soybean. *Genome*, 38: 715-772
- McCouch SR, Kochert G, Yu ZH, Wang ZY, Khush GS, Coffman WR, Tanksley SD (1988). Molecular mapping of rice chromosomes. *Theor. Appl. Genet.* 76: 815-829.
- McCouch SR, Temnykh S, Lukashova A, Coburn J, Declerck G, Cartinhour S, Harrington S, Thomson M, Septinginsi E, Semon M, Moncada P, Jiming L (2001). Microsatellite markers in rice: Abundance, diversity and applications. In: *Rice Genetics IV*. IRRRI, Manila, Philippines. pp: 117-135.
- Morgante M, Hanafey M, Powell W (2002). Microsatellites are preferentially associated with non-repetitive DNA in plant genomes. *Nat. Genet.* 30: 194-200.
- Mudge J, Cregan PB, Kenworthy JP, Orf JH, Young ND (1997). Two microsatellite markers that flank the major soybean cyst nematode resistance locus. *Crop Sci.* 37: 1611-1615.
- Mullan JD, Amanda P, Natasha L, Rudi TA, Timothy DC, Joseph MA, Michael GF (2005). EST-derived SSR markers from defined regions of wheat to identify *Lophopyrum elongatum* specific loci. *Genome*, 48(12): 811-822.
- Ravi M, Geethanjali S, Sameeyafarheen F, Maheswaran M (2003). Molecular marker based genetic diversity analysis in rice (*Oryza sativa* L.) using RAPD and SSR markers. *Euphytica*, 133: 243-252.
- Roder MS, Korzun V, Wendehake K, Plaschke J, Tixier MH, Leroy P, Ganai MW (1998). A microsatellite map of wheat. *Genetics*, 149: 2007-2023.
- Saghai Marooif MA, Biyashev RM, Yang GP, Zhang Q, Allard RW (1994). Extraordinarily polymorphic microsatellite DNA in barley: species diversity, chromosomal locations, and population dynamics. *Proc. Natl. Acad. Sci. USA*, 91: 5466-5470.
- Sayed NH, Hyeon SL, Minghua M, Peggy T, David MS, Feffrey Z (2001). Variability and evolution of microsatellite loci in cotton (*Gossypium*) diploid and polyploidy genomes. *Chem. Plant and Animal Genome IX Conference*, Town and Country Hotel, San Diego, C.A., January 13-17.
- Schlotterer C, Tautz D (1992). Slippage synthesis of simple sequence DNA. *Nucleic Acids Res.* 20: 211-215.
- Scott KD, Eggler P, Seaton G, Rossetto M, Ablet EM, Lee LS, Henry RJ (2000). Analysis of SSRs derived from grape ESTs. *Theor. Appl. Genet.* 100: 723-726.
- Smulders MJM, Bredemeijer G, Kortekaas WR, Arens P, Vosman B (1997). Use of short microsatellites from data-base sequences to generate polymorphisms among *Lycopersicon esculentum* cultivars and accessions of other *Lycopersicon* species. *Theor. Appl. Genet.* 94: 264-272.
- Tautz D (1989). Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Res.* 17: 6463-6471.
- Temnykh S, Park WD, Ayres N, Cartinhour S, Hauck N, Lipovtievich L (2000). Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 100: 697-712.
- Thiel T, Michalek W, Varshney RK, Graner A (2003). Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.* 106: 411-422.
- Thomas MR, Scott NS (1993). Microsatellite repeats in grapevine reveal DNA polymorphisms when analyzed as sequence-tagged sites (STSs). *Theor. Appl. Genet.* 86: 985-990.
- Trujillo I, Rallo L, Arus P (1995). Identifying olive cultivars by isozyme analysis. *J. Am. Soc. Hort. Sci.* 120: 318-324.
- Virk PS, Fordlloyd BV, Jackson MT, Newbury HJ (1995). Use of RAPD for the study of diversity within plant germplasm collections. *Heredity*, 97(2): 170-179.
- Wang HY, Wei YM, Yan ZH, Zheng YL (2007). ETS-SSR DNA Polymorphism in durum wheat (*Triticum aestivum* L.) Collections. *Appl. Genet.* 48: 35-42.
- Wang ZY, Tanksley SD (1989). Restriction Fragment Length Polymorphism in *Oryza sativa* L. *Genome*, 32: 1113-1118.
- Weising K, Atkinson RG, Gardner RC (1995). Genomic fingerprinting by microsatellite-primed PCR: a critical evaluation. *PCR Methods Appl.* 4: 249-255.
- Wren JD, Forgacs E, Fondon JW, Pertsemidis A, Cheng S, Gallardo T, Williams RS, Sorret RV, Minna JD, Garner HR (2000). Repeat Polymorphisms within Gene Regions - Phenotypic and Evolutionary Implications. *Am. J. Hum. Genet.* 67(2): 345-356.
- YQ-Wu Y, Huang DR, Porter CG, Tauer L, Hollaway (2007). Identification of major quantitative trait locus conditioning resistance to green bug biotype E in Sorghum PI 550610 using simple sequence repeat markers. *J Econ Entomol.* 100(5): 1672-1678.
- Yu JK, Mangor J, Edwards E, Thompson L, Knapp SJ (2000). Simple sequence repeats in sunflower: length polymorphisms among elite inbred lines for several motifs. In *Abstracts of Plant and Animal Genome Conference VIII*. San Diego, Calif. 9-12 January.
- Zane L, Bargelloni L, Patarnello T (2002). Strategies for microsatellite isolation: a review. *Mol. Ecol.* 11: 1-16.