

Full Length Research Paper

Non-sugar matter as an indicator of technological value in different sugar beet genotypes

Vladimir Filipović^{1*}, Stevan Radivojević², Jean Andrei³, Jonel Subić⁴ and Vladan Ugrenović¹

¹Institute "Tamiš", Pančevo, Serbia.

²Institute for Food Technology, Novi Sad, Serbia.

³Petroleum and Gas University of Ploiesti, Romania.

⁴Institute of Agricultural Economics, Belgrade, Serbia.

Accepted 15 September, 2011

This work represents a research of non-sugar matter (the content of α - amino nitrogen, potassium and sodium) in the root, and their effect on the recoverable sucrose quantity of three genotypes (Esprit, Belinda and Chiara) of sugar beet from vegetation areas of different sizes. They were grown by standard agrotechnical measures for sugar beet in the densities of 80.000, 100.000 and 120.000 plants per hectare. In 2005, the highest sugar utilization coefficient (12.94%) was achieved in the variant where plant density was 120.000 per hectare (0.083 m²) and the genotype was Esprit (N-type), which was 5.48% more than that in the variant with the lowest utilization coefficient (12.23%), where density was 100.000 per hectare (0.1 m²) and the genotype was Chiara (E-type). In 2006, the highest sugar utilization coefficient of 14.03% was achieved in the variant where plant density was 120.000 per hectare (0.083%) and the genotype Belinda (Z-type), which was 6.84% more than that in the variant with the lowest utilization coefficient (13.07%), with the density of 100.000 per hectare (0.1m²) and genotype Esprit (N-type). In 2007, the highest sugar utilization coefficient of 14.27% was recorded in the variant with the plant density of 80.000 per hectare (0.125 m²) and the genotype Belinda (Z-type), which was higher by 8.26% as compared to the variant with the lowest utilization coefficient (13.09%), where density was 100.000 per hectare (0.1m²) and the genotype was Chiara (E-type).

Key words: Sucrose content, recoverable sucrose, alpha amino nitrogen, potassium, sodium, genotype, sugar beet.

INTRODUCTION

Sugar beet technological value is determined by sucrose and non-sugar matter content in the storage root. Non-sugar components (α -amino-nitrogen or harmful nitrogen, potassium, sodium, magnesium, calcium) in the root reduces sugar crystallization from molasses during sugar beet processing (Sinobad and Brdar, 1996; Bashir et al., 2000; Sklenar et al., 2000; Radivojević and Kabić, 2000; Leilah et al., 2005; Pytlarz-Kozicka, 2005; Jaćimović et al., 2006; Mahmood et al., 2007; Filipović, 2009). The percentage of crystallized sugar from molasses is calculated with Reinefeld's formula. The obtained value is

the percentage of total sugar content utilization from the storage root. The percentage of crystal sugar content is affected by a multitude of factors. Significant differences between total and crystal sugar content are caused by the size of plant's vegetative site as well as by sugar beet genotypes (Filipović, 2009).

The selection of the best suited genotype should involve the testing of large number of genotypes differing in sugar synthesis intensity and sugar accumulation in the storage root. Genotypes of a shorter vegetative period have faster initial growth and earlier sugar synthesis hence they achieve earlier technological maturity, that is, the period of sugar optimal utilization from the root starts earlier. These genotypes are referred to as sugary genotypes or Z-type genotypes, and they are suitable for early harvest dates. E-type genotypes, on the other hand, have the longest sugar accumulation period and they

*Corresponding author. E-mail: vladimirfilipovic@gmail.com.
Tel: +381 (0)13 313092 or +381 (0)62 522642. Fax: +381 (0)13 2520991.

Table 1. Analysis of variance for indicators of technological value of sugar beet root (2005).

Source of variation	df	Mean squares				
		Sucrose content	Potassium content	Sodium content	α -Amino nitrogen content	Recoverable sugar content
Blocks	3	0.30 ^{NS}	0.07 ^{NS}	0.20*	0.13 ^{NS}	0.30 ^{NS}
Plant density	2	0.61*	0.09 ^{NS}	0.69**	0.03 ^{NS}	0.61*
Genotype	2	0.11 ^{NS}	0.06 ^{NS}	0.03 ^{NS}	0.03 ^{NS}	0.11 ^{NS}
Plant density x Genotype	4	0.14 ^{NS}	0.02 ^{NS}	0.06 ^{NS}	0.07 ^{NS}	0.14 ^{NS}
Error	24	0.13	0.09	0.05	0.05	0.13

NS = Non-significant; * = significant at 0.05 probability level; ** = significant at 0.01 probability level.

are suitable for late harvest dates (Ramadan, 1999). Total sugar content in the root and percentage of sugar utilization is considerably affected by nitrogen rates and arrangement of fertilizer application and plant nutrition (Halvorson and Hartman, 1975).

This paper presents, firstly, the studies of non-sugar matter content in the root of three sugar beet genotypes (Belinda Z-type, Esprit N-type and Chiara E-type) grown at different crop densities and, secondly, the effect of non-sugar matter on percentage of crystal sugar sucrose.

MATERIALS AND METHODS

Three-year trials were conducted during 2005 to 2007 period on the experimental field of the Institute of "Tamiš" Pančevo (N 44° 56', E 20° 43') as a randomized complete block design with four replications. The size of vegetative site for individual plants was 0.125 m² (80.000 plants per hectare), 0.1 m² (100.000 plants per hectare) and 0.083 m² (120.000 plants per hectare). It was formed by crop hand thinning during a sprouting stage. The studies comprised genotypes Belinda (KWS – 2004. Z-type), Esprit (Strube Dieckmann – 2004. N-type) and Chiara (KWS – 2004. E-type). Sugar beet was hand harvested and analysis was conducted on the sugar beet roots from middle rows of the elementary site. About 150 g of pulp from each plot was prepared by Venema apparatus and kept in a freezer until analysis. Frozen sugar beet pulp samples were analyzed in sugar technology laboratory in Sugar Factory "Jedinstvo" from Kovačica of Serbia for purity parameters with Betalyser. Betalyser is a computer-controlled system for automated routine analysis of sugar beet on sugar content and impurities including Na⁺, K⁺ and α -amino-N. Sugar content (S⁰) was measured by polarimetry, Na⁺ and K⁺ by flame-emission photometry and α -amino-N by double beam filter photometry using the blue number method (Sheikh-Aleslami, 1997).

The values of potassium, sodium and harmful nitrogen were expressed in mmol/100 g of sugar beet, while content of total sugar sucrose and crystal sugar content are given in percentages (%). Based on the results obtained, computations were done for crystal sugar percentage on Reinefeld formula (1974):

$$SU = S^0 - [0.343 \times (K + Na) + 0.094 \times \alpha\text{-amino-N} + 0.29]\%$$

Where, SU is sugar utilization and S⁰ is sugar content.

The obtained data were analyzed by mathematical-statistic methods (Snedecor and Cochran, 1971). The statistical analysis was performed according to the design established in each field using the SYSTAT 7.0 statistical package (SYSTAT, 1997). Analysis of variance was applied to estimate significance derived by

F-test and LSD-test for 5 and 1% significance.

RESULTS AND DISCUSSION

In the first study year, the impact of crop density on sodium content ($p < 0.01$) was highly statistically significant. In the same year, the recorded statistical significance of the effect of crop density on sucrose content and sugar utilization coefficient was at the significance level of 0.05 (Tables 1 and 2).

The highest content of total sugar was found in genotype Esprit grown on the vegetative site of 0.083 m² (13.25%) and the lowest was found in genotype Chiara, with vegetative site of 0.1 m² (12.54%). In genotype Esprit, the content of harmful nitrogen was highest in the crop of the lowest density (2.69 mmol/100 g of sugar beet) and lowest when the size of vegetative site was 0.1 m² (2.40 mmol/100 g of sugar beet). Many researchers have also suggested that choice of genotypes increased the yield and quality of sugar beet (Lauer, 1997; Wyszynski et al., 1998; Sohrabi and Heidari, 2008). Genotype Chiara had the highest potassium and sodium content in the root. On the vegetative site of 0.125 m², the content of potassium was 3.38 mmol/100 g of sugar beet and in genotype Belinda (0.1 m²) it amounted to 3.03 mmol/100 g of sugar beet. According to Kessel and Schladen (1984), the values of potassium content lower than 4 mmol/100 g of sugar beet are low, while those that are higher than 5 mmol/100 g of sugar beet are high. The highest difference in sodium content per genotype was found between genotype Chiara (1.80 mmol/100 g of sugar beet) and genotype Esprit (1.14 mmol/100 g of sugar beet) with identical size of the vegetative site (0.1 m²). The amounts of non-sugar matter which varied per genotype and crop density influenced the percentage of crystal sugar sucrose too. The lowest content of crystal sugar was in genotype Chiara on the vegetative site of 0.1 m² (12.23%) and it was highest in genotype Esprit on the vegetative site of 0.083 m² (12.94%). These results conform to the reports by Radivojević and Kabić (2000).

Interaction of genotypes and plant density produced significant effects on difference in sodium content potassium ($P < 0.05$). Variations per genotype were not

Table 2. Effect of sugar beet genotypes and plant density treatments combination on sucrose content and technological value of sugar beet root (2005).

Genotype (A) and plant density (B)	Sucrose content (%)	Potassium content (mmol/100 g of root)	Sodium content (mmol/100 g of root)	α -Amino nitrogen content (mmol/100 g of root)	Recoverable sugar content (%)
Genotype (A)					
Esprit	12.95 ^{at}	3.28 ^a	1.47 ^a	2.61 ^a	12.64 ^a
Chiara	12.95 ^a	3.22 ^a	1.43 ^a	2.58 ^a	12.64 ^a
Belinda	12.78 ^a	3.13 ^a	1.37 ^a	2.52 ^a	12.48 ^a
Plant density (B)					
80000	13.13 ^a	3.30 ^a	1.66 ^a	2.62 ^a	12.82 ^a
100000	12.88 ^{ab}	3.16 ^a	1.44 ^b	2.54 ^a	12.57 ^{ab}
120000	12.68 ^b	3.16 ^a	1.18 ^c	2.53 ^a	12.37 ^b

[†]Means not sharing a common letter in a column differ significantly at 0.05 probability.

Table 3. Analysis of variance for indicators of technological value of sugar beet root (2006).

Source of variation	df	Mean squares				
		Sucrose content	Potassium content	Sodium content	α -Amino nitrogen content	Recoverable sugar content
Blocks	3	0.69 ^{NS}	0.09 ^{NS}	0.08 ^{NS}	0.23 ^{NS}	0.69 ^{NS}
Plant density	2	0.67 ^{NS}	0.05 ^{NS}	0.09 ^{NS}	0.30 ^{NS}	0.67 ^{NS}
Genotype	2	0.13 ^{NS}	0.01 ^{NS}	0.02 ^{NS}	0.04 ^{NS}	0.13 ^{NS}
Plant density x Genotype	4	0.29 ^{NS}	0.14 ^{NS}	0.51 [*]	0.11 ^{NS}	0.29 ^{NS}
Error	24	0.29	0.10	0.13	0.35	0.29

NS = Non-significant; * = significant at 0.05 probability level; ** = significant at 0.01 probability level.

significant. In the second study year, individual variations in total sugar and non-sugar matter content as well as sugar utilization from juice were not significant (Tables 3 and 4).

The highest total sugar content (14.34%) was found in genotype Belinda at crop density of 120.000 plants per hectare, and the lowest was in genotype Esprit (13.38%) at the density of 100.000 plants per hectare. In average, the lowest values of potassium, sodium and harmful α -amino nitrogen content were found in the variant genotype Esprit \times 100.000 plants per hectare. This automatically resulted to the lowest sugar utilization in this variant, while the highest sugar utilization was found in genotype Belinda at crop density of 120.000 plants per hectare.

In the third study year, the effect of crop density showed extremely high statistical significance for sugar content, sodium content and utilization coefficient ($p < 0.01$). However, the coefficient of sugar utilization was at the same level; over 97.6%. According to the results reported by Sinobad and Brdar (1996), non-sugar components significantly reduced total sucrose content in the root but did not significantly reduce the coefficient of

sugar utilization. Sklenar et al. (2000), on the other hand, pointed out that non-sugar matter content significantly influences the coefficient of sugar utilization. Technological value of sugar beet roots, at total average, was highest in the third study year (Tables 5 and 6).

The size of vegetative site influenced total sugar content in the root. As crop density was increased, total sugar content was decreased; 14.09 to 13.89%. This difference was significant. The coefficient of nitrogen utilization in the soil depends on the number of plants per unit area. In more dense crops, the competition is great, which causes reduced crop nitrogen supply and lesser production of sugar (Winter, 1990). Variations in total sugar content per genotype were not significant. Non-sugar matter content was significantly influenced by crop densities. As vegetative site was increased, sodium content was also increased; 1.25 to 1.42 mmol/100 g of sugar beet, the obtained difference was very significant. Sugar utilization from juice was higher in the roots that were developed on a larger vegetative site, and variations from crops of highest density to crops of lowest density were very significant. Increased content of non-sugar matter did not significantly influence sugar

Table 4. Effect of sugar beet genotypes and plant density treatments combination on sucrose content and technological value of sugar beet root (2006).

Genotype (A) and Plant density (B)	Sucrose content (%)	Potassium content (mmol/100 g of root)	Sodium content (mmol/100 g of root)	α -amino nitrogen content (mmol/100 g of root)	Recoverable sugar content (%)
Genotype (A)					
Esprit	13.77 ^{at}	2.58 ^a	1.64 ^a	2.92 ^a	13.46 ^a
Chiara	13.76 ^a	2.54 ^a	1.59 ^a	2.88 ^a	13.45 ^a
Belinda	13.59 ^a	2.52 ^a	1.57 ^a	2.81 ^a	13.28 ^a
Plant density (B)					
80000	13.93 ^a	2.61 ^a	1.69 ^a	3.05 ^a	13.62 ^a
100000	13.73 ^a	2.54 ^a	1.59 ^a	2.84 ^a	13.42 ^a
120000	13.46 ^a	2.49 ^a	1.52 ^a	2.74 ^a	13.15 ^a

[†]Means not sharing a common letter in a column differ significantly at 0.05 probability.

Table 5. Analysis of variance for indicators of technological value of sugar beet root (2007).

Source of variation	df	Mean squares				
		Sucrose content	Potassium content	Sodium content	α -Amino nitrogen content	Recoverable sugar content
Blocks	3	0.23 ^{NS}	0.06 ^{NS}	0.04 ^{NS}	0.52 ^{NS}	0.23 ^{NS}
Plant density	2	2.39 ^{**}	0.33 ^{NS}	1.28 ^{**}	0.56 ^{NS}	2.39 ^{**}
Genotype	2	0.13 ^{NS}	0.07 ^{NS}	0.02 ^{NS}	0.16 ^{NS}	0.12 ^{NS}
Plant density x Genotype	4	0.12 ^{NS}	0.04 ^{NS}	0.06 ^{NS}	0.47 ^{NS}	0.11 ^{NS}
Error	24	0.24	0.22	0.03	0.28	0.25

NS = Non-significant; * = significant at 0.05 probability level; ** = significant at 0.01 probability level.

Table 6. Effect of sugar beet genotypes and plant density treatments combination on sucrose content and technological value of sugar beet root (2007).

Genotype (A) and Plant density (B)	Sucrose content (%)	Potassium content (mmol/100 g of root)	Sodium content (mmol/100 g of root)	α -Amino nitrogen content (mmol/100 g of root)	Recoverable sugar content (%)
Genotype (A)					
Esprit	14.09 ^{at}	3.26 ^a	1.46 ^a	2.32 ^a	13.78 ^a
Chiara	14.00 ^a	3.19 ^a	1.42 ^a	2.23 ^a	13.70 ^a
Belinda	13.88 ^a	3.11 ^a	1.37 ^a	2.10 ^a	13.58 ^a
Plant density (B)					
80000	14.30 ^a	3.28 ^a	1.64 ^a	2.43 ^a	14.00 ^a
100000	14.19 ^a	3.28 ^a	1.56 ^a	2.24 ^a	13.89 ^a
120000	13.48 ^b	2.99 ^a	1.04 ^b	2.00 ^a	13.17 ^b

[†]Means not sharing a common letter in a column differ significantly at 0.05 probability.

crystallization, so in the third study year, its coefficient of utilization was very high; over 97.7%. Variations in the amount of total sugar and non-sugar matter per genotype

were not significant and depended individually on the size of vegetative site and environmental conditions throughout vegetative period in sugar beet. These results are in

agreement with the findings of Rother (2000), Rosso and Candolo (2001) and Hoffman et al. (2009).

Conclusions

Three sugar beet genotypes were studied for the effects of “non-sugar” matter on sugar utilization of different technological value at different crop densities. Genotype Esprit (N – type) gave on average, the highest sucrose and recoverable sugar content. Plant density significantly influenced the quality parameters (sugar content, K, Na and alpha-amino-nitrogen). The highest recoverable sugar content was achieved at plant densities of 100.000 plants per hectare. Higher densities led to decrease in quality parameters. According to the study, genotype Belinda (Z – type) is more related to medium and low plant density.

ACKNOWLEDGEMENTS

This paper represents a part of the research results of the Project of Integral and Interdisciplinary Research (No. 046006) “Sustainable Agriculture and Rural Development in the Function of the Accomplishment of Strategic Objectives of the Republic of Serbia in the Danube Region”, funded by the Ministry of Education and Science of the Republic of Serbia.

REFERENCES

- Filipović V (2009). The Effect of Crop Density on the Growth, Yield and Quality of the Root in Different Sugar Beet Varieties. Doctoral Dissertation. Faculty of Agriculture. Belgrade. Zemun.
- Bashir S, Saeed M, Ghaffar A, Ali Z, and Zuffarullah RMK (2000). Analysis of Economic Aspects of Raising Autumn Sugarcane at Different Planting Patterns and Seeding Densities, *International J. of Agric. & Biol.* 1560–8530/2000/02–4–322–325.
- [Halvorson AD, Hartman GP \(1975\). Long-term nitrogen rates and sources influence sugar beet yield and quality. *Agron. J.* 67: 389-393.](#)
- Hoffman CM, Huijbregts T, Swaaji NV, Jansen R (2009). Impact of different environments in Europe on yield and quality of sugar beet genotypes. *European J. Agron.* 30: 17-26.
- Jačimović G, Marinković B, Crnobarac J (2006). The Yield of Refined Sugar Depending on the Level of Sugar Beet Fertilization. *Contemporary Agric.* 55(5): 139-145.
- Kessel W, Schladen Ch (1984). Qualität und Zuckerertrag—die Weichen Werdun bereits im Fruhjaer Gesreilit. *Die Zuckerrube.* 33(2): 68-73.
- [Lauer J \(1997\). Sugar Beet Performance and Interactions with Planting Date, Genotype, and Harvest Date. *Agron. J.* 89: 469-475.](#)
- Leilah AA, Badawi MA, Said EM, Ghonema MH, Abdou MAE (2005). Effect of Planting Dates, Plant Population and Nitrogen Fertilization on Sugar Beet Productivity Under the Newly Reclaimed Sandy Soils in Egypt. *Scientific J. King Faisal Univ.* 6(1): 525-529.
- Mahmood M, Ishfaq M, Iqbal J, Nazir, MS (2007). Agronomic Performance and Juice Quality of Autumn Planted Sugarcane (*Saccharum officinarum* L.) as Affected by Flat, Ditch and Pit Planting Under Different Spatial Arrangements. *International J. of Agric. & Biol.* 1560–8530/2007/09–1–167–169.
- Pytlarz-Kozicka M (2005). The effect of nitrogen fertilization and anti-fungal plant protection on sugar beet yielding. *Plant, Soil Environ.* 51(5): 232-236.
- Radivojević S, Kabić D (2000). Biological and Technological Properties of the Varieties in Sugar Beet Production in 1998. *Acta Periodica Tehnologica. Novi Sad.* 31(A). APTEFF. 31. 2000: 333-338.
- Ramadan BSH (1999). Differential response of some sugar beet varieties to plant density and harvesting dates. *J. Agric. Sci. Mansoura University,* 24(2): 413-423.
- Reinefeld von E (1974). Rubenqualität und Zucker aus beute. *Die Zuckerrube.* 2(23): 13-15.
- Rosso F, Candolo G (2001). Evaluation of sugar beet main quality features through the analysis of the diffusion juices produced by a pilot plant. 64. IIRB Congress, Bruges, Belgium, 26-27 Jun 2001. 64: 437-442.
- Rother B (2000). The internal quality of sugar beet influenced by different growing factors. 63. IIRB Congress, Interlaken, Switzerland, 9-10 Feb 2000: pp. 279–294.
- Sheikh Aleslami R (1997). Laboratorial Methods and their Application to Control Food and Sugar Industries Process. Mersa Publications. Tehran. Iran.
- Sinobad V, Brdar P (1996). Analysis of the Effect of the Percentages of Sucrose, Potassium, Sodium and Alpha-amino Nitrogen in Sugar Beet on the Value of Sucrose Content. *Chem. Ind.* 50. (3-4): 83-87.
- Sklenar P, Kovačev L, Čačić N, Željka S (2000). The Effect of Potassium, Sodium and Alpha-amino Nitrogen on Sugar Utilization in Sugar Beet Hybrids. *Acta Periodica Tehnol. Novi Sad.* 31(A). APTEFF. 31. 2000: 339-344.
- Snedecor J, Cochran V (1971). *Statistical Methods.* “Vuk Karadzic”. Belgrade.
- Sohrabi Y, Heidari G (2008). Investigating the influence of withholding irrigation and harvest times on yield and quality of sugar beet (*Beta vulgaris*). *Int. J. Agric. Biol.* 10: 427-431.
- SYSTAT (1997). SYSTAT for Windows, Version 7.0. SPSS Inc. Chicago. Illinois. USA.
- [Winter SR \(1990\). Sugar beet Response to Nitrogen as Affected by Seasonal Irrigation. *Agron. J.* 82: 984-988.](#)
- Wyszynski Z, Kalinowska-Zdun M, Rozbicki J, Madry W, Laudanski Z, Roszkowska B (1998). Agronomical and environmental factors affecting sugar beet yielding in Central and Eastern Poland. *Sci. Agric. Boh.* 29(3-4): 183-192.