

Full Length Research Paper

Occurrence, isolation and DNA identification of *Streptococcus thermophilus* involved in Algerian traditional butter ‘Smen’

Farid Bensalah^{1*}, Asmaa Labtar¹, Christine Delorme² and Pierre Renault²

¹Faculté des Sciences, Département de Biologie, LGM (Laboratoire de Génétique Microbienne), Université Es-Sénia, Oran 31000, Algérie.

²INRA-CRJ, Institut National de la Recherche Agronomique, Micalis Institute, Département de Microbiologie et Chaîne Alimentaire. 78352 Jouy-En- Josas, cedex, France.

Accepted 19 October, 2011

Streptococcus thermophilus isolates from traditional butter ‘Smen’, a fermented product from cow’s and ewe’s milk in arid area was subjected to taxonomical investigations. The identification procedure included phenotypic approaches, molecular characterization by using genus polymerase chain reaction (PCR) amplifications for *sodA* gene encoding the manganese-dependant superoxide dismutase A, and species-specific primers from gene encoding glucose kinase (*glcK*), gene encoding DNA polymerase III (*dnaE*) and gene encoding threonyl-tRNA synthetase (*thrS*) housekeeping genes in order to distinguish among reference, and wild strains of *S. thermophilus* and for their differentiation from *Enterococcus* spp. A total of 12 strains were tested by DNA identification analysis and these indigenous isolates were unambiguously characterised by their housekeeping gene profiles. Finally, four genotypes were recognised. To our knowledge, this is the first report on the isolation and molecular characterization of *S. thermophilus* strains from fermented milk transformed into home-production butter provided from Southern West Algeria. These results are of particular interest as they favour the selection of future cultures starters as *S. thermophilus* from traditional dairy products. It is, therefore, of great importance to preserve the genetic pool of the wild strains.

Key words: *Streptococcus thermophilus*, traditional butter, PCR, *Enterococcus*, DNA, species-specific primer.

INTRODUCTION

The microbial composition of traditional dairy products and, particularly, of European artisanal cow’s and ewe’s fermented milk and cheeses, have been subjected to microbiological investigation at genus and species level (Tornadjio et al., 1995; Cogan et al., 1997; Mannu et al., 2000; Mannu and Paba, 2002; Alegría et al., 2009; Secchi et al., 2011). Many indigenous fermented foods from Algeria are based on lactic fermentation from cow’s and ewe’s milk (Bensalah et al., 2009). Among these,

‘Smen’ is a traditional butter, which, so far has not been investigated for microbiological genotypic strains characterization. The nature of this fermented product is different from one region to another and depends on the local indigenous microflora; which in turn reflects the climatic conditions of the area. However, there is very little information on strain composition within species, and on the specific contribution of each wild strain to the lactic fermentation. Moreover, the study of strain heterogeneity in African traditional butter (El-Baradei et al., 2008; Ongol and Asano, 2009), and cheese starters is believed to be of great importance to the dairy industry (Girrafa et al., 1998).

Fermented milk ‘leben’ or ‘rayeb’ (Elotmani et al., 2002; Hegazi, 1988; Bensalah et al., 2009) can be eaten fresh or suffers a churning and creaming in goatskin called

*Corresponding author. E-mail: bensalahfarid@yahoo.fr.

Abbreviation: PCR, Polymerase chain reaction; *glcK*, gene encoding glucose kinase; *dnaE*, gene encoding DNA polymerase III; *thrS*, gene encoding threonyl-tRNA synthetase.

'Chekoua'. Skimming is usually done in the morning; gourd is half filled with milk fermented, then strained by swelling, and then the 'Chekoua' is tied and shaken vigorously for half an hour. The formation of fat globules is considered by the sound change that occurs within the skin. To assist the agglomeration of particles of butter, water is usually added, hot or cold depending on the temperature of milk. The fresh butter is removed manually in a single lump called 'Zebda' which is transformed into 'Smen' by washing with salt water. 'Smen' has a very strong smell, easily recognizable during ripening. Implication of *Streptococcus thermophilus* in Egyptian artisanal butter 'zabady' was reported recently (El-Baradei et al., 2008). This strain is an essential micro-organism in the manufacture of many types of fermented dairy products, occurring in natural as well as commercial starter cultures (Coppola et al., 1998; Delorme, 2008), and the increasing number of commercial strains available used as starters requires rapid and reliable methods to accurately differentiate strains at both species and strain levels in pure and mixed cultures in order to protect patent rights and eliminate risks of use confusion (Ramos and Harlander, 1990; Moschetti et al., 1998). *Enterococcus* strains can harbour specific biochemical traits that are essential in the manufacture of various fermented milk products, and some strains are technologically exploited as functional starters or probiotics (Giraffa et al., 1997; Franz et al., 1999; Bensalah et al., 2006). Enterococci has also been implicated in the spoilage of processed meats (Franz and Holy, 1996) and include strains that have been recognized as emerging human pathogens mostly in nosocomial urinary tract infections and bacteremias (Giraffa et al., 2000; Martinez and Baquero, 2002).

In the last few years, the development of new methods involving various DNA-based typing techniques (Farber, 1996), widely used in epidemiological studies for the identification of medically important bacterial strains (Kuhn et al., 1995; Descheemaeker et al., 1997), has opened up new perspectives for typing strains from raw milk and from traditional cheeses (Bouton et al., 1998; Desmasures et al., 1998). It was previously reported that a genus-specific PCR assay based on the use of degenerate primers enabled amplification of an internal fragment representing approximately 85% of the *sodA* gene encoding a manganese-dependent superoxide dismutase (Mn-SOD) in various gram-positive bacteria including genera of *Streptococcus* (Poyart et al., 1998) and *Enterococcus* (Poyart et al., 1995). The *sodA* gene has been identified as a target for the identification of mycobacteria (Zolg and Philippi-Schulz, 1994) and staphylococci at the species level by PCR (Poyart et al., 2001). Another *Streptococcus* species-specific, PCR-based identification method was reported recently that utilises the housekeeping gene encoding glucose kinase (*glcK*), gene encoding DNA polymerase III (*dnaE*) and gene encoding threonyl-tRNA synthetase (*thrS*) (Delorme et al., 2007), and the oligonucleotide primers from the

intergenic sequences 16S-23S rRNA (Tissala-Timisjarvi and Alatossava, 1997).

Moreover, since little is known about the involvement of thermophilic cocci with Algerian artisanal foods, this study was aimed at characterizing phenotypically and genotypically, the streptococci associated with the traditional fermented cow's and ewe's raw milk transformed into home-production butter provided from southern west Algeria. We applied the developed methodology of genus and species-specific primers from *sodA*, *glcK*, *dnaE* and *thrS* housekeeping genes in order to distinguish among reference, and wild strains of *S. thermophilus* and for their differentiation from *Enterococcus* spp.

MATERIALS AND METHODS

Sampling and bacterial strains

Six samples of cow's and ewe's traditional butter 'Smen' were obtained from individual households in rural arid area. Samples were collected in sterile small bottles and stored in Laboratoire de Génétique Microbienne, Es-Sénia, Algeria (LGM) under refrigeration at 4°C until use. Reference strains representing four Gram-positive species were used in this study and were provided from INRA collection (Jouy-En-Josas, France). These included *S. thermophilus* CNRZ 1066, *Enterococcus faecalis* JH2-2, *Enterococcus faecium* 64/3 and *Corynebacterium glutamicum* ATCC 13032.

Isolation and phenotypic characterization of selected isolates

Serial dilutions of homogenized 10 g of 'Smen' samples in 90 ml sterile solution saline 0.85% NaCl were used for microbial isolation with M17 agar (Terzaghi and Sandine, 1975) containing 1% lactose (LM17). Plates were incubated anaerobically at 37, 45 or 50°C for 48 to 72 h for isolation of thermophilic streptococci. Identification was carried out according to previously published characterization schemes (Devriese et al., 1993; Moschetti et al., 1998; Muller et al., 2001; Michaylova et al., 2007). Gram staining, catalase activity, ammonia production from arginine and gas production from glucose were determined. Tests for growth on Slanetz-Bartley (SB) medium, in M17 broth at 10°C, pH value adjusted to 9.6 and with NaCl 6.5% were performed for all the isolates. Colonies morphology were determined for overnight cultures by using phase-contrast microscopy and all isolates were stored in liquid cultures with 15% glycerol at -80°C.

Oligonucleotides

The first preliminary phenotypic investigation was followed by using the nucleotide sequences of internal fragment with PCR identification analysis. *Streptococcus* /*Enterococcus* genus-specific primers *sodAd1/sodAd2* (encoding superoxide dismutase) as previously described (Poyart et al., 1998, 2000) and housekeeping genes of *Streptococcus* species-specific primers from *glcK*, *dnaE* and *thrS*, (Delorme et al., 2007) were used in this study.

DNA template preparation

Template DNA was prepared by a modification of the method of

Table 1. PCR primers.

Primer	Sequence	Product size (bp)
<i>sodAd1</i> (universel-F)	5'-CITAYICITAYGAYGCIYTIGARCC-3'	438; Enterococcus
<i>sodAd2</i> (universel-R)	5'-ARRTARTAIGCRTGYTCCCAIACRTC-3'	480; Streptococcus
<i>Streptococcus thermophilus</i> (<i>glcK</i>)	5'-TGGGCAGAAACTCAAGA-3' 5'-AACACCACCACCGATAAC-3'	470
<i>Streptococcus thermophilus</i> (<i>dnaE</i>)	5'-GGACTGGGAGCCTGGGAT-3' 5'-ACTCCCTGCAGCAGACCC-3'	480
<i>Streptococcus thermophilus</i> (<i>thrS</i>)	5'-ATCACTGAAGATGGAAGC-3' 5'-CCAAGTTTACGGTGGTCA-3'	497

I = A, T, G or C; Y = C or T; R = A or G.

Moschetti et al. (1998). Total DNA was extracted from 100 µl of overnight cultures which was picked from one colony and resuspended in M17 broth. Cells were pelleted in a microcentrifuge at 8000 × *g* for 5 min, resuspended in 20 µl of GeneReleaser™ (Bio Ventures, Inc.) and lysed using PCR program as follows: 65°C for 30 s, 8°C for 30 s, 65°C for 1 min 30 s, 97°C for 3 min, 8°C for 1 min, 65°C for 3 min, 97°C for 1 min, 65°C for 1 min and 80°C for 5 min.

Amplification of internal part from *sodA* gene by using degenerate primers

PCR procedure was performed in 25 µl reaction mixtures containing: 2.5 µl 10X PCR buffer, 2 µl dNTP, 0.5 µl of each primer, 0.25 µl of *Taq* DNA polymerase (Takara) and 5 µl of DNA from cellular lysates. Amplification was carried out in a Thermal cycler (Applied Biosystem), as follows: an initial denaturation step of 95°C for 3 min, 30 cycles of amplification (60 s of annealing at 37°C, 60 s of elongation at 72°C, and 30 s of denaturation at 95°C) and 7 min at 72°C for the last elongation. PCR primers pairs used in this study are shown in Table 1. An amount of 5 µl of the PCR product was electrophoresed at 100 V on a 1% agarose gel. The documented sizes of the amplicons were 480 and 438 bp from *sodA* gene in *Streptococci* and *Enterococci* respectively (Poyart et al., 1998, 2000). 200 bp DNA Smartladder (Eurogentec) was used as molecular size marker and DNA of references strains was included.

Streptococcus-specific PCR

DNA identification analysis by PCR was followed with *Streptococcus* species-specific primers from *S. thermophilus* CNRZ1066 used as the positive control. The cellular lysates were obtained with the generereleaser™ (bio Ventures, inc.) from overnight cultures as described above and the following three primers pairs were used for this study (Table 1). The amplification profile included a denaturation step of 94°C for 5 min, and then subjected to 30 cycles of 94°C for 30 s, 50°C for 30 s, 72°C for 1 min and a final soak at 4°C. DNA fragments were determined by comparing the positive control bands from reference strains with amplified fragments of wild members within indigenous flora isolates. The documented bands of the amplicons were 470, 480 and 497 from *glcK*, *dnaE* and *thrS* respectively (Delorme et al., 2007). The presence of PCR products was determined by gel electrophoresis in 1% agarose gel containing ethidium bromide. Electrophoresis in

Tris-borate-EDTA was performed at 100 V, and DNA Smartladder (Eurogentec) of 200 bp was used as the molecular size marker.

RESULTS

Biochemical and morphological characteristics

12 strains were selected and presumptively identified as cocci thermophilic lactic acid bacteria (LAB). All the strains were Gram-positive, catalase negative and produced no gas from glucose. Eight strains did not grow at all or poorly in 6.5% NaCl, three of them were able to grow at pH 9.6 and showed no or only weak growth on SB medium. Morphology cells were spherical, ovoid in short or means and long chain. Enterococci are reported to grow at 10°C, but six strains presumptively identified in these genera by classical methods failed to do so. The contradictions were established in strains which grew well with pink colonies in SB agar but failed to grow in broth with 6.5% NaCl and at pH 9.6. Therefore, another identification approach was necessary to clarify the taxonomic positions of these atypical isolates.

Analysis of DNA amplicons products from *sodA* gene

Identification by using the universal primers of *sodA* database of Gram-positive cocci is frequently reported particularly from genus and species of streptococci and enterococci. In this work, the specificity of the genus was determined by testing all isolates presumptively identified as thermophilic lactic acid bacteria in short and long chain. Only five strains reacted with *sodAd1/sodAd2* genus universal primers, and occurred as a single product. One band of weak intensity at 438 bp in the references and three isolates (LGM5, LGM6 and LGM7) and another band of strong intensity located approximately between 438-480 bp was revealed in only the reference and two selected strains (LGM1 and LGM2).




Figure 1. Genus and species-specific PCR amplifications for *sodA* gene. **(a)** Genus PCR with primer pair d1/d2. Lane M, 200-pb DNA molecular mass marker; lane 1, *S. thermophilus* CNRZ 1066 as positive control; lane 2, LGM1 isolate; lane 3, LGM2 isolate; lane 4, *C. glutamicum* ATCC 13032 as negative control; lane 5, LGM5 isolate; lane 6, LGM6 isolate; lane 7, LGM7 isolate; lane 8, *E. faecalis* JH2-2 as positive control; lane 9, *E. faecium* 64/3 as positive control. **(b)** Species-specific PCR with primer pair *glcK*-up/*glcK*-dn. Lane M, 200-pb DNA molecular mass marker; lane 1, LGM5 isolate; lane 2, isolate 6; lane 3, LGM6 isolate; lane 4, LGM7 isolate; lane 5, isolate 7; lane 6, isolate 8; lane 7, LGM1; lane 8, isolate 9; lane 9, LGM2; lane 10, *E. faecalis* JH2-2 as negative control; lane 11, isolate 10; lane 12, *E. faecium* 64/3 as negative control; lane 13, isolate 11; lane 14, isolate 12; lane 15, *St. thermophilus* CNRZ 1066 as positive control. **(c)** Species-specific PCR with primer pair *dnaE*-up/*dnaE*-dn. Lane M, 200-pb DNA molecular mass marker; lane 1, LGM5 isolate; lane 2, LGM6 isolate; lane 3, LGM7 isolate; lane 4, isolate 6; lane 5, LGM1; lane 6, LGM2; lane 7, isolate 7; lane 8, isolate 9; lane 9, isolate 9; lane 10, LGM3; lane 11, LGM4; lane 12, isolate 10; lane 13, isolate 11; lane 14, isolate 12; lane 15, *E. faecalis* JH2-2 as negative control. **(d)** Species-specific PCR with primer pair *thrS*-up/*thrS*-dn. Lane M, 200-pb DNA molecular mass marker; lane 1, *Ent. faecium* 64/3 as negative control; lane 2, *E. faecalis* JH2-2 as negative control; lane 3, LGM1; lane 4, LGM2; lane 5, isolate 6; lane 6, LGM 3; lane 7, LGM5; lane 8, LGM6; lane 9, LGM4; lane 10, LGM7, lane 11, isolate 7, lane 12, isolate 8; lane 13, isolate 9; lane 14, isolate 10; lane 15, isolate 11; lane 16, isolate 12.

No significant common bands were found from other isolates, and from the control-negative strain *C. glutamicum* (Figure 1a). These results represent a similar profile and do not indicate the discriminating power of the universal primers of *sodA* gene PCR described here.

Amplification using streptococcal species

On the basis of the unclear results of phenotypic identification and the weak discriminating power of *sodA* gene between *Streptococcus* and *Enterococcus* genus,

species-specific PCR of *glcK*, *dnaE* and *thrS* from streptococci were carried out to determine the taxonomic identification and to compare the isolates with reference strains. In one hand, the expected 470-pb band from *glcK* gene was revealed in the streptococci reference strain and two isolates (LGM1 and LGM2). However, a double product band was visualised also at approximately 600 bp in two reference enterococci strains and in three traditional butter isolates (LGM5, LGM6, and LGM7), while no significant bands were found from the other strains tested (Figure 1b). Two other strains suspected to be streptococci were confirmed by using primers of *glcK*,




Figure 2. Microscopic cell morphology of wild *Streptococcus thermophilus* LGM1, LGM2, LGM3 and LGM4 isolated from traditional butter 'Smen' ($\times 1000$).

LGM3 and LGM4, (results not shown).

On the other hand, the specific 480 and 497 bp streptococci bands from *dnaE* and *thrS* respectively were obtained in four isolates which reacted also with *glcK* from all isolates tested, (Figure 1c, d). Finally, in this study, we demonstrated that four isolates reacted with the three species-specific *glcK*, *dnaE* and *thrS* amplicons used in this work, indicating that they are members of *Streptococcus thermophilus* which are provided from Algerian traditional butter 'Smen'. These results indicate the discriminating power of the *glcK*-up/*glcK*-dn, *dnaE*-up/*dnaE*-dn and *thrS*-up/*thrS*-dn primers described in this work. This evidence allowed us to clearly differentiate *Streptococcus thermophilus* (Figure 2) from *Enterococcus* spp.

DISCUSSION

While the association of Streptococci and Enterococci

with European fermented foods is well known (Moschetti et al., 1998; Cogan et al., 1997; Sengun et al., 2009; Secchi et al., 2011), little is known about the involvement of these bacteria in African fermented foods (Youssif et al., 2005). Streptococci were shown in this study to form a predominantly inhabitants of the microflora of fermented milk product 'Smen', as these bacteria could be isolated throughout the fermentation process. Enterococci were also present in 'Smen', they appear to be often associated with fermented products from Africa, although at a high incidence compared with other cocci lactic acid bacteria (LAB) (Mathara et al., 2004; Bensalah et al., 2006, 2009).

Streptococcus and *Enterococcus* are Gram-positive cocci, commonly recognized by their growth on Rogosa agar plates. These genera differ phenotypically; however, biochemical and physiological tests often give imprecise specifications. Due to complicated and uncertain identification, efforts have been made to develop simple, fast and discriminating methods for identification of

Streptococcus from *Enterococcus* species (Moshetti et al., 1998). The result of this study demonstrates the necessity for use of molecular methods which is crucial for successful identification of unknown thermophilic cocci strains. PCR analysis is useful for clearly discriminating a multitude of species of LAB (like *Streptococcus*, *Lactococcus* and *Enterococcus* species) (Mannu and Paba, 2002; Bensalah et al., 2006, 2009). Species-specific PCR was previously used for the identification of clinical and food isolates of Enterococci and Streptococci (Poyart et al., 1998; Delorme et al., 2007).

The primers used for the *sodA* gene in this study gave a DNA profile almost similar to Streptococci and Enterococci; however, the use of three other primer pairs *glcK*, *dnaE* and *thrE* helped highlight different profiles and amplified the expected DNA bands. The loci chosen formed a suitable basis for DNA identification procedures and the housekeeping gene provided evidence and demonstrated that species-specific *glcK*, *dnaE* and *thrE* discriminate among streptococci and enterococci isolates. PCR methods in this work proved to be useful for species identification and to reveal different patterns throughout this genus. We also attempted to show that phenotypic characterisation and amplified DNA amplicons using PCR analyses were quick, reliable and convenient methods to differentiate Streptococci from Enterococci. PCR assays developed here are technically affordable for laboratories that do not have easy access to more sophisticated or expensive procedures. Also, the genotypic and phenotypic analysis carried out on wild Streptococci and Enterococci isolated from traditional fermented 'Smen' highlighted diversity among strains. This study enabled the dominance biotypes in 'Smen' to be defined at each period of maturation and, it was found that the strains that dominated the first stage of ripening were not necessarily predominant in the later periods (Mannu et al., 2000).

To our knowledge, this is the first report on the occurrence, isolation and molecular identification of *S. thermophilus* in 'Smen' in Algeria, and further investigation into strain variability within species should be extended not only to more old samples but also, to dairy artisanal product from spontaneously fermented milk. The analysis of bacterial properties as synthesis of exopolysaccharides (Minic et al., 2007), proteolysis, aroma, acidifying power and antagonistic activity must be realised. These studies are of particular interest as they favour the isolation and selection of future cultures starters as *S. thermophilus* from traditional dairy products. It is, therefore, of great importance to preserve the genetic pool of strains of lactic acid bacteria in Africa.

REFERENCES

Alegria A, Alvarez-Martín P, Sacristán N, Fernández E, Delgado S, Mayo B (2009). Diversity and evolution of the microbial populations

- during manufacture and ripening of Casin, a traditional Spanish, starter-free cheese made from cow's milk. *Int. J. Food Microbiol.* 136: 44-51.
- Bensalah F, Maria-José F, Mouats A (2006). A rapid PCR based method to distinguish between *Enterococcus* species by using degenerate and species-specific *sodA* gene primers. *Afr. J. Biotechnol.* 5: 697-702.
- Bensalah F, Delorme C, Renault P (2009). Characterisation of thermotolerant cocci from indigenous flora of 'leben' in Algerian arid area and DNA identification of atypical *Lactococcus lactis* strains. *Curr. Microbiol.* 59: 139-46.
- Bouton Y, Guyot P, Grappin R (1998). Preliminary characterization of microflora of Comté cheese. *J. Appl. Microbiol.* 85: 123-31.
- Cogan TM, Barbosa M, Beuvier E, Bianchi-Salvadore B, Cocconcelli PH, Fernandez PS, Gomez I, Kalantzopoulos G, Ledda A, Medina M, Rea MC, Rodriguez E (1997). Characterisation of the lactic acid bacteria in artisanal dairy products. *J. Dairy Res.* 64: 409-421.
- Coppola S, Parente E, Dumontet S, La Peccerella A (1998). The microflora of natural whey cultures utilized as starters in the manufacture of Mazzarella cheese from water-buffalo milk. *Le Lait.* 68: 295-313.
- Descheemaeker P, Lammens C, Pot B, Vandamme P, Goossens H (1997). Evaluation of arbitrarily primed PCR analysis and pulsed-field gel electrophoresis of large genomic DNA fragments for identification of enterococci important in human medicine. *Int. J. Syst. Bacteriol.* 47: 555-561.
- Desmaures N, Mangin I, Corrolier D, Guéguen M (1998). Characterization of lactococci isolated from milk produced in the Camembert region of Normandy. *J. Appl. Microbiol.* 85: 999-1005.
- Delorme C, Poyart C, Ehrlich SD, Renault P (2007). Extent of horizontal gene transfer in evolution of Streptococci of the *Salivarius* group. *J. Bacteriol.* 189: 1330-1341.
- Delorme C (2008). Safety assessment of dairy microorganisms: *Streptococcus thermophilus*. *Int. J. Food Microbiol.* 126: 274-277.
- Devriese LA, Pot B, Collins MD (1993). Phenotypic identification of the genus *Enterococcus* and differentiation of phylogenetically distinct enterococcal species and species groups. *J. Appl. Bacteriol.* 75: 399-408.
- El-Baradei G, Delacroix-Buchet A, Ogier JC (2008). Bacterial biodiversity of traditional Zabady fermented milk. *Int. J. Food Microbiol.* 121: 295-301.
- Elotmani F, Revol-Junelles AM, Assobhei O, Millière JB (2002). Characterisation of anti-*Listeria monocytogenes* bacteriocins from *Enterococcus faecalis*, *Enterococcus faecium*, and *Lactococcus lactis* strains isolated from Raib, a Moroccan traditional fermented milk. *Curr. Microbiol.* 44: 10-17.
- Farber JM (1996). An introduction to the hows and whys of molecular typing. *J. Food Prot.* 59: 1091-1101.
- Franz CM, von Holy A (1996). Thermotolerance of meat spoilage lactic acid bacteria and their inactivation in vacuum-packaged vienna sausages. *Int. J. Food Microbiol.* 29: 59-73.
- Franz CMAP, Holzappel WH, Stiles ME (1999). Enterococci at the crossroads of food safety? *Int. J. Food Microbiol.* 47: 1-24.
- Giraffa G, Carminati D, Neviani E (1997). Enterococci isolated from dairy products: a review of risks and potential technological use. *J. Food Prot.* 60: 732-738.
- Giraffa G, De Vecchi P, Rossi P, Nicastro G, Fortina MG (1998). Genotypic heterogeneity among *Lactobacillus helveticus* strains isolated from natural cheese starters. *J. Appl. Microbiol.* 85: 411-416.
- Giraffa G, Olivari AM, Neviani E (2000). Isolation of vancomycin-resistant *Enterococcus faecium* from Italian cheeses. *Food Microbiol.* 17: 671-677.
- Hegazi FZ (1988). Proteolysis and electrophoretic pattern of casein of some fermented milks. *Nahrung.* 32: 539-543.
- Kühn I, Burman LG, Hæggen S, Tullus K, Murray BE (1995). Biochemical fingerprinting compared with ribotyping and pulsed-field gel electrophoresis of DNA for epidemiological typing of enterococci. *J. Clin. Microbiol.* 33: 2812-2817.
- Mannu L, Paba A, Pes M, Scintu MF (2000). Genotypic and phenotypic heterogeneity among lactococci isolated from traditional Pecorino Sardo cheese. *J. Appl. Microbiol.* 89: 191-197.
- Mannu L, Paba A (2002). A Genetic diversity of lactococci and

- enterococci isolated from home-made Pecorino Sardo ewes' milk cheese. *J. Appl. Microbiol.* 92: 55-62.
- Martinez JL, Baquero F (2002). Interactions among strategies associated with bacterial infection: pathogenicity, epidemicity, and antibiotic resistance. *Clin. Microbiol. Rev.* 15: 647-679.
- Mathara JM, Schillinger U, Kutima PM, Mbugua SK, Holzapfel WH (2004). Isolation, identification and characterisation of the dominant microorganisms of kule naoto: the Maasai traditional fermented milk in Kenya. *Int. J. Food Microbiol.* 94: 269-278.
- Michaylova M, Minkova S, Kimura K, Sasaki T, Isawa K (2007). Isolation and characterization of *Lactobacillus delbrueckii* ssp. *bulgaricus* and *Streptococcus thermophilus* from plants in Bulgaria. *FEMS Microbiol. Lett.* 269: 160-169.
- Minic Z, Marie C, Delorme C, Faurie JM, Mercier G, Ehrlich D, Renault P (2007). Control of EpsE, the phosphoglycosyltransferase initiating exopolysaccharide synthesis in *Streptococcus thermophilus*, by EpsD tyrosine kinase. *J. Bacteriol.* 189: 1351-1357.
- Moschetti G, Blaiotta G, Aponte M, Catzeddu P, Villani F, Deina P, Coppola S (1998). Random amplified polymorphic DNA and amplified ribosomal DNA spacer polymorphism: powerful methods to differentiate *Streptococcus thermophilus* strains. *J. Appl. Microbiol.* 85: 25-36.
- Müller T, Ulrich A, Ott EM, Müller M (2001). Identification of plant-associated enterococci. *J. Appl. Microbiol.* 91: 268-278.
- Ongol MP, Asano K (2009). Main microorganisms involved in the fermentation of Ugandan ghee. *Int. J. Food Microbiol.* 133: 286-291.
- Poyart C, Berche P, Trieu-Cuot P (1995). Characterization of superoxide dismutase genes from Gram-positive bacteria by polymerase chain reaction using degenerate primers. *FEMS Microbiol. Lett.* 131: 41-45.
- Poyart C, Quesne G, Coulon S, Berche P, Trieu-Cuot P (1998). Identification of streptococci to species level by sequencing the gene encoding the manganese-dependent superoxide dismutase. *J. Clin. Microbiol.* 36: 41-47.
- Poyart C, Quesnes G, Trieu-Cuot P (2000). Sequencing the gene encoding manganese-dependent superoxide dismutase for rapid species identification of enterococci. *J. Clin. Microbiol.* 38: 415-418.
- Poyart C, Quesne G, Boumaila C, Trieu-Cuot P (2001). Rapid and accurate species-level identification of coagulase-negative staphylococci by using the *sodA* gene as a target. *J. Clin. Microbiol.* 39:4296-4301.
- Ramos MS, Harlander SK (1990). DNA fingerprinting of lactococci and streptococci used in dairy fermentations. *Appl. Microbiol. Biotechnol.* 34: 368-374.
- Secchi N, Giunta D, Pretti L, García MR, Roggio T, Mannazzu I, Catzeddu P (2011). Bioconversion of ovine scotta into lactic acid with pure and mixed cultures of lactic acid bacteria. *J. Ind. Microbiol. Biotechnol.* [Epub ahead of print]. 10 : 1013-1019
- Sengun IY, Nielsen DS, Karapinar M, Jakobsen M (2009). Identification of lactic acid bacteria isolated from Tarhana, a traditional Turkish fermented food. *Int. J. Food Microbiol.* 135: 105-111.
- Terzaghi BE, Sandine WE (1975). Improved medium for lactic streptococci and their bacteriophages. *Appl. Microbiol.* 29: 807-813.
- Tissala-Timisjarvi A, Alatossava T (1997). Development of oligonucleotide primers from the 16S-23S rRNA intergenic sequences for identifying different dairy and probiotic lactic acid bacteria by PCR. *Int. J. Food Microbiol.* 35: 49-56.
- Tornadijo ME, Fresno JM, Bernardo A, Sarmiento RM, Carballo J (1995). Microbiological changes throughout the manufacturing and ripening of a Spanish goats' raw milk cheese (Armada variety). *Le Lait.* 75: 551-570.
- Youssif NM, Dawyndt P, Abriouel H, Wijaya A, Schillinger U, Vancanneyt M, Swings J, Dirar HA, Holzapfel WH, Franz CM (2005). Molecular characterization, technological properties and safety aspects of enterococci from 'Hussuwa', an African fermented sorghum product. *J. Appl. Microbiol.* 98: 216-228.
- Zolg J, Philippi-Schulz S (1994). The superoxide dismutase gene, a target for detection and identification of mycobacteria by PCR. *J. Clin. Microbiol.* 32: 2801-2812.