

Full Length Research Paper

Biochemical and kinetic characterization of geranylgeraniol 18-hydroxylase (CYP97C27) from *Croton stellatopilosus* Ohba

Manee Chanama¹ and Suchart Chanama^{2*}¹Department of Microbiology, Faculty of Public Health, Mahidol University, Bangkok 10400, Thailand.²Department of Biochemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.

Received 18 April, 2015; Accepted 13 July, 2015

Geranylgeraniol 18-hydroxylase (EC 1.14.13.110) that exists solely in *Croton stellatopilosus* Ohba catalyses the last committed step of plaunotol biosynthetic pathways by conversion of geranylgeraniol (GGOH) to plaunotol. This enzyme and its gene are an attractive target for development of plaunotol production and its detailed biochemical properties need to be understood. Recently, even though the gene (CYP97C27) coding for GGOH 18-hydroxylase has been identified, cloned, and expressed in *Escherichia coli* system, the enzyme activity has been detected mainly in the insoluble fraction (20,000 g). This means that biochemical and kinetic studies could not be undertaken. However, our previous study indicated that this enzyme activity was easily and specifically detected in the microsomal fraction (100,000 g) of a crude enzyme extract. Therefore, in this report we describe a comprehensive biochemical characterization of GGOH 18-hydroxylase activity in the microsomal fraction from *C. stellatopilosus* Ohba. The oxygen-dependent enzyme activity of GGOH 18-hydroxylase was inhibited by carbon monoxide and the inhibition was partially reversible upon illumination with white light. Kinetic studies of the GGOH 18-hydroxylase showed high affinity to GGOH and NADPH with apparent K_m values of 0.8 and 53 μM , respectively. Furthermore, the enzyme activity was inhibited by P450 inhibitors, including ancymidol, metyrapone, miconazole, potassium cyanide and cytochrome c, with the IC_{50} values of 428, 65, 75, 66 and 8 μM , respectively. Based on the biochemical and kinetic characteristics, the GGOH 18-hydroxylase in the microsomal fraction is likely a P450 encoded by CYP97C27 gene as previously described.

Key words: Plaunotol, cytochrome P450, enzyme activity, enzyme inhibitor, microsome.

INTRODUCTION

Plaunotol, an acyclic diterpenoid compound, has important chemotherapeutic activities; it exhibits an anti-inflammatory action in association with peptic ulcers and

antimicrobial activities against *Helicobacter pylori* and *Staphylococcus aureus* (Koga et al., 2002; Inoue et al., 2004; Premprasert et al., 2013). It was originally found in

*Corresponding author. E-mail: suchart.c@chula.ac.th. Tel: +662 218 5433. Fax: +662 218 5418.

Author(s) agree that this article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Abbreviations: GGPP, Geranylgeranyl diphosphate; GGOH, geranylgeraniol; PMSF, phenylmethylsulfonyl fluoride; KCN, potassium cyanide; CO, carbon monoxide.




Figure 1. The biosynthetic pathway of plaunotol in *C. stellatopilosus* via deoxyxylulose phosphate pathway (modified from Nualkaew et al., 2006).

leaves and stems of *Croton stellatopilosus* Ohba (Euphorbiaceae), a medicinal plant growing in tropical Southeast Asian countries, especially Thailand. It appears to be accumulated mainly in the chloroplasts (Wungsintaweekul and De-Eknamkul, 2005; Sithithaworn et al., 2006). Previous study on plaunotol biosynthesis in *C. stellatopilosus* has revealed that biosynthesis of plaunotol was carried out via the deoxyxylulose phosphate pathway in which geranylgeranyl diphosphate (GGPP) and geranylgeraniol (GGOH) were the intermediate precursors (Nualkaew et al., 2005; Wungsintaweekul and De-Eknamkul, 2005). As shown in Figure 1, GGPP is dephosphorylated by phosphatase to form GGOH, then the GGOH is subsequently hydroxylated at the C-18 position by GGOH 18-hydroxylase (EC 1.14.13.110) to form plaunotol, where the hydroxylation of GGOH exhibits the requirement of NADPH as a reducing equivalent of the reaction (Tansakul and De-Eknamkul, 1998; Nualkaew et al., 2005; Nualkaew et al., 2006). This suggested that the GGOH 18-hydroxylase is a member of cytochrome P450

monooxygenases (CYP), similar to other P450 hydroxylases involved in plant secondary metabolisms (Collu et al., 2001; Jennewein et al., 2003). As yet, there are no more details with regard to the biochemical and kinetic properties of GGOH 18-hydroxylase although the corresponding gene (*CYP97C27*) has been identified, cloned and expressed in an *Escherichia coli* system (Sintupachee et al., 2014). The expressed GGOH 18-hydroxylase protein was highly detectable in the insoluble fraction of the crude enzyme, however it was difficult to obtain in highly purified solubilized form and thus it was not amenable to detailed characterization. However, all P450 enzymes in plants have been reported to be localized to the microsomal fraction (Schuler, 1996) and the activity of our enzyme of interest (GGOH 18-hydroxylase) has been shown to exist in the 100,000 g microsomal pellet fraction of crude enzyme extract. Therefore, in this work, we investigated the biochemical and kinetic properties of GGOH 18-hydroxylase in the 100,000 g microsomal fraction.

MATERIALS AND METHODS

Plant materials and chemicals

Fresh mature leaves of *C. stellatopilosus* Ohba were kindly provided by the Institute of Biotechnology and Genetic Engineering, Chulalongkorn University, Bangkok, Thailand, and maintained at -20°C until used. All chemicals were of the highest purity available. Geranylgeraniol (GGOH), β -nicotinamide adenine dinucleotide phosphase (reduced form) tetrasodium salt (NADPH), β -mercaptoethanol, ancymidol, metyrapone, miconazole and cytochrome c were purchased from Sigma Chemical Co., (USA). Tricine and dithiothreitol (DTT) from USB (USA). Phenylmethylsulfonyl fluoride (PMSF) and potassium cyanide (KCN) from Fluka (Switzerland). Ethyl acetate and absolute ethanol were of HPLC grade and purchased from Lab-Scan Asia (Thailand). Plaunotol was obtained from Kelnac[®] soft gelatin capsules (Sankyo, Co., Japan).

Preparation of microsomal fraction containing GGOH 18-hydroxylase activity

All enzyme preparation steps were performed at 0 to 4°C . The GGOH 18-hydroxylase was prepared and purified according to the method as previously described (Chanama et al., 2009). Briefly, 30 g of frozen mature leaves were ground rapidly to a powder in liquid nitrogen using a prechilled mortar and pestle. Then, the fine powder was extracted in 60 ml of extraction buffer containing 83 mM Tricine-NaOH (pH 7.8), 0.4 M sucrose, 10 mM EDTA, 10 mM MgCl_2 , 10 mg/ml BSA, 1 mM DTT, 0.8 mM PMSF and 5 mM β -mercaptoethanol, and the mixture was stirred for 10 min. The homogenate was filtered through several layers of cheesecloth, centrifuged at 3,000 g for 10 min and the supernatant collected prior to spinning for 20 min at 20,000 g. Resulting supernatant was further ultracentrifuged at 100,000 g for 60 min and the microsomal precipitate was suspended in 3 to 5 ml of 0.1 mM Tricine-NaOH (pH 7.8) containing 0.2 M sucrose, 1 mM EDTA, 1 mM DTT, 15% glycerol and 5 mM β -mercaptoethanol. The final enzyme preparation containing the GGOH 18-hydroxylase was stored at -80°C , and protein concentration of the enzyme was measured using the Bio-Rad protein assay (Bradford, 1976).

GGOH 18-hydroxylase activity assays

Enzyme activity of GGOH 18-hydroxylase was assayed as previously described (Chanama et al., 2009). The reaction mixture consisted of 83 mM Tricine-NaOH (pH 7.8), 0.8 mM NADPH and 57 μ M GGOH (substrate). The reaction was initiated by the addition of 100 μ l of enzyme preparation (~ 250 μ g of protein) to the mixture. The reaction mixture was incubated at 30°C for 30 min and then stopped by extraction twice with an equal volume of ethyl acetate. The extracts were pooled and dried, and the residue was redissolved in ethyl acetate for analysis of plaunotol. Plaunotol content in the extract was determined using a TLC-densitometric technique as previously described. Briefly, the extract of enzyme product was applied onto silica gel₆₀ F₂₅₄ plates (CAMAG) using Linomat IV (CAMAG). The plate was developed by ethyl acetate and scanned at wavelength of 210 nm to obtain chromatogram of the sample. The plaunotol content was then estimated on the basis of the standard calibration curve of pure plaunotol compound. The assays were performed in triplicate. The enzyme activity is expressed as katal unit (kat). One katal is the amount of enzyme required to convert GGOH substrate to one mole of plaunotol product per second. For kinetic studies of the GGOH 18-hydroxylase, the maximum velocity (V_{max}) and the Michaelis constant (K_m) values were determined under the standard assay conditions with substrate concentrations ranging from 0.2 to 10 μ M for GGOH and from 0.02 to 1.0 mM for NADPH.

Inhibition experiments

The inhibition by gaseous carbon monoxide (CO) and reversal of CO inhibition by white light were performed in 2 ml septum-capped glass vials containing all the reaction components except GGOH and NADPH. The vials were placed on ice, and each gas (CO, N₂ or air) was bubbled through the reaction mixtures. The reactions were initiated by the addition of GGOH and NADPH, and then incubated at 30°C for 30 min in the dark or under white light (for light reversal of CO inhibition). After incubation, the reaction products were extracted and analysed as described above. The control vials without the addition of inhibitors were carried out in the dark. The cytochrome P450 inhibitors: ancymidol, metyrapone, miconazole, KCN and cytochrome c were tested. The concentrations of ancymidol, metyrapone, miconazole and KCN were adjusted from 0.01 to 6.0 mM and cytochrome c from 0.003 to 0.3 mM. The enzyme inhibition tests were carried out by addition of the inhibitors to the standard assay. After completion of the reactions, the enzyme products (plaunotol) were extracted and analysed by TLC. The inhibition is expressed as IC₅₀ value, that is, the concentration of an inhibitor where the enzyme activity is reduced by 50%.

RESULTS AND DISCUSSION

The enzyme GGOH 18-hydroxylase found in *C. stellatopilosus* catalyses the conversion of GGOH to plaunotol by addition of hydroxyl group (-OH) to acyclic diterpenoid GGOH at position of C-18 and the hydroxylation reaction is highly specific to this acyclic diterpene substrate and not for other terpenoid compounds (C-10: geraniol, C-15: farnesol) (Tansakul and De-Eknamkul, 1998). In addition, the activity was determined 1.5-fold higher in 100,000 g microsomal fraction than in the 20,000 g insoluble fraction under the presence of NADPH and aeration (Chanama et al., 2009)

and also observed in other plant sources (Bolwell et al., 1994, Pierrel et al., 1994). In this work, we studied the kinetic properties of the GGOH 18-hydroxylase enzyme in the 100,000 g microsomal fraction with its substrate and cofactor. The results showed that the GGOH 18-hydroxylase was saturated at approximately 7 μ M GGOH and 500 μ M NADPH, and apparent K_m values for the hydroxylation of GGOH and for NADPH were 0.8 and 53 μ M, respectively (Figure 2). The very low K_m value of the hydroxylase for GGOH implies the high affinity of the GGOH 18-hydroxylase for the GGOH substrate, and this result is in good agreement with the K_m values obtained for the hydroxylation of laurate by lauric acid (P450) monooxygenase from Jerusalem-Artichoke (*Helianthus tuberosus*) (0.97 μ M) (Salaun et al., 1978) and from wheat (*Triticum aestivum*) (8 μ M) (Zimmerlin et al., 1992). Interestingly, the apparent K_m value of this enzyme for diterpenoid GGOH was 20 to 60 fold lower than those for the hydroxylations of other fatty acid (16-hydroxypalmitic acid, 50 μ M), acyclic monoterpene (geraniol, 15 μ M), cyclic monoterpene (limonene, 18 to 21 μ M), cyclic diterpene (taxoid, 50 μ M) and phenolic compound (cinnamic acid, 35 μ M) from other plant species (Soliday and Kolattukudy, 1978; Karp et al., 1990; Hallahan et al., 1992; Petersen, 1997; Jennewein et al., 2003). In addition, consumption of NADPH in the hydroxylation is essential for the enzyme activity and in a good agreement with the result reported by Tansakul and De-Eknamkul (1998). The K_m value observed in this study certainly supports the view that GGOH and NADPH are actual substrate and cofactor in the enzyme system.

To investigate whether the GGOH 18-hydroxylase present in the microsomal fraction belongs to a member of cytochrome P450 monooxygenase, several of the following criteria, that is, a requirement for molecular oxygen (O₂), inhibition by CO and reversal of the CO inhibition by light, and inhibition by specific cytochrome P450 inhibitors were examined (Krochko et al., 1998; Kim et al., 2004). To address whether molecular oxygen (O₂) was essential for GGOH 18-hydroxylase activity, replacement of dissolved oxygen in the reaction mixture by nitrogen gas prior to initiating reaction was performed. The reaction exhibited a dramatic decrease in enzyme activity (85% inhibition) (Table 1). Moreover, carbon monoxide, which is known to bind effectively to Fe(II)-heme region of the cytochrome P450 (Krochko et al., 1998), also exhibited the strongest inhibition when it was introduced into the enzyme system in the dark. The relative activity of GGOH 18-hydroxylase inhibited by CO was reduced to as low as 5.87% of control (94% inhibition). This inhibition was partially reversible upon illumination with visible light (relative activity of 35%) as shown in Table 1. This partial reversion of CO-inhibition by white light (particularly at wavelength of 450 nm) could be due to the photolysis of Fe (II)-CO complex which exists in the GGOH 18-hydroxylase. The effects of specific cytochrome P450 inhibitors (ancymidol,




Figure 2. Michaelis-Menten plots showing the variation of initial velocity of GGOH 18-hydroxylase reaction as a function of GGOH (A) and NADPH (B) concentrations. The double reciprocal plots $1/v$ as a function of $1/[GGOH]$ and of $1/[NADPH]$ are shown in the insets. The values are means (\pm SEM) of three separate experiments.

Table 1. Effect of oxygen (air), nitrogen (N_2) and carbon monoxide (CO), and white light on the activity of GGOH 18-hydroxylase from *C. stellatopilosus*.

Treatment	Relative activity ^a (% of control)	Inhibition (%)
Control (air, dark)	100.0	00.00
N_2	14.47	85.53
CO (dark)	05.87	94.13
CO (white light)	35.56	64.44

^aThe enzyme activity of the control was 1.77 pkat/ mg protein.

metryrapone, miconazole, cytochrome c and KCN) were used in this study. All cytochrome P450 inhibitors inhibited the enzyme activity of GGOH 18-hydroxylase completely or to a lesser extent and with variable concentrations for half-maximal activity (IC_{50} values) (Figure 3). Ancymidol, metyrapone, miconazole and KCN exhibited 100% inhibitory effects on the hydroxylation reaction at concentrations ranging from 0.01 to 6 mM with IC_{50} values of 428, 65, 75 and 66 μ M, respectively.

On the other hand, cytochrome c which is known to remove electrons competitively from NADPH-cytochrome




Figure 3. Inhibitions of GGOH 18-hydroxylase activity by cytochrome P450 inhibitors. The values are means (\pm SEM) of three separate experiments.

P450 reductase complex (Petersen, 1997), showed about 75% inhibition with the lowest IC_{50} of 8 μ M. Among these inhibitors tested, cytochrome c seemed to be the most potent inhibitor against GGOH 18-hydroxylase (IC_{50} of 8 μ M). These results strongly support a typical feature of plant cytochrome P450s of the enzyme (Friederich et al., 1999; Yamamoto et al., 2000; Katano et al., 2001). According to the biochemical and kinetic properties of the

GGOH 18-hydroxylase, that is, high affinity to GGOH substrate, requirement for molecular oxygen and NADPH, inhibition by P450 inhibitors, inhibition by CO, and reversal of the CO inhibition by light, it is likely that the GGOH 18-hydroxylase in the 100,000 g microsomal fraction of *C. stellatopilosus* is the CYP97C27 previously identified by RT-PCR methodology from leaves of *C. stellatopilosus* (Sintupachee et al., 2014).

Conflict of interests

The authors did not declare any conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by Chulalongkorn University and Mahidol University. The authors would like to thank the Institute of Biotechnology and Genetic Engineering, Chulalongkorn University for leaves of *C. stellatopilosus* Ohba. This study was supported for publication by the China Medical Board (CMB), Faculty of Public Health, Mahidol University, Bangkok, Thailand.

REFERENCES

- Bolwell GP, Bozak K, Zimmerlin A (1994). Plant cytochrome P450. *Phytochemistry* 37:1491-1506.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72:248-254.
- Chanama M, Wunnakup T, De-Eknamkul W, Chanama S (2009). Improvement of thin-layer chromatography for enzyme assay of geranylgeraniol 18-hydroxylase from *Croton stellatopilosus* Ohba. *JPC-J. Planar Chromat.* 22:49-53.
- Collu G, Unver N, Peltenburg-Looman AMG, van der Heijden R, Verpoorte R, Memelink J (2001). Geraniol 10-hydroxylase, a cytochrome P450 enzyme involved in terpenoid indole alkaloid biosynthesis. *FEBS Lett.* 508:215-220.
- Friederich S, Rueffer M, Asakawa Y, Zenk MH (1999). Cytochromes P-450 catalyze the formation of marchantins A and C in *Marchantia polymorpha*. *Phytochemistry* 52:1195-1202.
- Hallahan DL, Dawson GW, West JM, Wallsgrove RM (1992). Cytochrome P-450 catalysed monoterpene hydroxylation in *Nepeta mussinii*. *Plant Physiol. Biochem.* 30:435-443.
- Inoue Y, Shiraiishi A, Hada T, Hirose K, Hamashima H, Shimada J (2004). The antibacterial effects of terpene alcohols on *Staphylococcus aureus* and their mode of action. *FEMS Microbiol.* 237:325-331.
- Jennewein S, Rithner CD, Williams RM, Croteau R (2003). Taxoid metabolism: Taxoid 14 beta-hydroxylase is a cytochrome P450-dependent monooxygenase. *Arch. Biochem. Biophys.* 413:262-270.
- Karp F, Mihaliak CA, Harris JL, Croteau R (1990). Monoterpene biosynthesis: specificity of the hydroxylations of (-)-limonene by enzyme preparations from peppermint (*Mentha piperita*), spearmint (*Mentha spicata*), and perilla (*Perilla frutescens*) leaves. *Arch. Biochem. Biophys.* 276:219-226.
- Katano N, Yamamoto H, Iio R, Inoue K (2001). 7-Deoxyloganin 7-hydroxylase in *Lonicera japonica* cell cultures. *Phytochemistry* 58:53-58.
- Kim T-W, Chang SC, Lee JS, Hwang B, Takatsuto S, Yokota T, Kim S-K (2004). Cytochrome P450-catalyzed brassinosteroid pathway activation through synthesis of castasterone and brassinolide in *Phaseolus vulgaris*. *Phytochemistry* 65:679-689.
- Koga T, Inoue H, Ishii C, Okazaki Y, Domon H, Utsui Y (2002). Effect of plaunotol in combination with clarithromycin or amoxicillin on *Helicobacter pylori* in vitro and in vivo. *J. Antimicrob. Chemother.* 50:133-136.
- Krochko JE, Abrams GD, Loewen MK, Abrams SR, Cutler AJ (1998). (+)-abscisic acid 8'-hydroxylase is a cytochrome P450 monooxygenase. *Plant Physiol.* 118:849-860.
- Nualkaew N, De-Eknamkul W, Kutchan TM, Zenk MH (2005). Geranylgeraniol formation in *Croton stellatopilosus* proceeds via successive monodephosphorylations of geranylgeranyl diphosphate. *Tetrahedron Lett.* 46:8727-8731.
- Nualkaew N, De-Eknamkul W, Kutchan TM, Zenk MH (2006). Membrane-bound geranylgeranyl diphosphate phosphatases: purification and characterization from *Croton stellatopilosus* leaves. *Phytochemistry* 67:1613-1620.
- Petersen M (1997). Cytochrome P450-dependent hydroxylation in the biosynthesis of rosmarinic acid in coleus. *Phytochemistry* 45:1165-1172.
- Pierrel MA, Batard Y, Kazmaier M, Mignotte-Vieux C, Durst F, Werck-Reichhart D (1994). Catalytic properties of the plant cytochrome P450 CYP73 expressed in yeast. *Eur. J. Biochem.* 224:835-844.
- Premprasert C, Tewtrakul S, Plubrukarn A, Wungsintaweekul J (2013). Anti-inflammatory activity of diterpenes from *Croton stellatopilosus* on LPS-induced RAW264.7 cells. *J. Nat. Med.* 67:174-181.
- Salaun JP, Benveniste I, Reichhart D, Durst F (1978). A microsomal (cytochrome P450)-linked lauric-acid-monooxygenase from aged jerusalem-artichoke-tuber tissues. *Eur. J. Biochem.* 90:155-159.
- Schuler MA (1996). Plant cytochrome P450 monooxygenases. *Crit. Rev. Plant Sci.* 15:235-284.
- Sintupachee S, Ngamrojanayanich N, Sitthithaworn W, De-Eknamkul W (2014). Molecular cloning, bacterial expression and functional characterisation of cytochrome P450 monooxygenase, CYP97C27, and NADPH-cytochrome P450 reductase, CPR I, from *Croton stellatopilosus* Ohba. *Plant Sci.* 229:131-141.
- Sitthithaworn W, Potduang B, De-Eknamkul W (2006). Localization of plaunotol in the leaf of *Croton stellatopilosus* Ohba. *Science Asia* 32:17-20.
- Soliday CL, Kolattukudy PE (1978). Midchain hydroxylation of 16-hydroxypalmitic acid by the endoplasmic reticulum fraction from germinating *Vicia faba*. *Arch. Biochem. Biophys.* 188:338-347.
- Tansakul P, De-Eknamkul W (1998). Geranylgeraniol-18-hydroxylase: the last enzyme on the plaunotol biosynthetic pathway in *Croton sublyratus*. *Phytochemistry* 47:1241-1246.
- Wungsintaweekul J, De-Eknamkul W (2005). Biosynthesis of plaunotol in *Croton stellatopilosus* proceeds via the deoxyxylulose phosphate pathway. *Tetrahedron Lett.* 46:2125-2128.
- Yamamoto H, Katano N, Ooi A, Inoue K (2000). Secologanin synthase which catalyzes the oxidative cleavage of loganin into secologanin is a cytochrome P450. *Phytochemistry* 53:7-12.
- Zimmerlin A, Salaun J-P, Durst F, Mioskowski C (1992). Cytochrome P-450-dependent hydroxylation of lauric acid at the subterminal position and oxidation of unsaturated analogs in wheat microsomes. *Plant Physiol.* 100:868-873.