

*Full Length Research Paper*

# Impact of bee pollen as feed supplements on the body weight of broiler Ross 308

Peter Haščík<sup>1</sup>, Ibrahim Elimam<sup>1\*</sup>, Jozef Garlík<sup>1</sup>, Miroslava Kačániová<sup>2</sup>, Juraj Čuboň<sup>1</sup>,  
Marek Bobko<sup>1</sup> and Hasan Abdulla<sup>3</sup>

<sup>1</sup>Department of Animal Products Evaluation and Processing, Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic.

<sup>2</sup>Department of Microbiology, Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic.

<sup>3</sup>Department of Animal Nutrition, Faculty of Agrobiological and Food Science, Slovak University of Agriculture, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic.

Accepted 5 October, 2012

**This study was aimed at investigating the meat performance of broiler Ross 308 after the application of bee pollen extract in their diet. The experiment group was fed with bee pollen (400 mg.kg<sup>-1</sup>) added in the feed. After analysis, it was found that the female chicken group's average live body weight were higher in control group (2246.60 g) than that of experimental group (2194.40 g). Also, the carcass weight, giblet weight and carcass yield percentage were high in the control group than in the experimental group. However, in case of male group, the average live body weight in experimental group (2354.60 g) were higher than control group (2299.20 g). In addition, the weights (g) of carcass, giblets, and carcass yield (%) were higher in the experimental group than control group, and there were no significant differences ( $P \geq 0.05$ ) among the experimental groups. Hence, it was concluded that bee pollen has positive effect on the growth of male chicken in terms of increasing the body weight, whereas it has negative effect on the female chickens.**

**Key words:** Broiler, pollen, carcass weight, live body weight, carcass yield.

## INTRODUCTION

The prominence of poultry production today is primarily due to the short generation interval and relatively quick turn over on investment and high quality protein from poultry products (Adeyemo et al., 2010). Production of poultry meat for the rapidly growing human population is an important system for supplying high-quality protein and provides an interesting source of finance (Gueye, 2009). The ratio of the composition of feed mixtures for chickens is important in terms of the required nutrients and energy. The increasing energy and nutrients in chickens' feed mixtures are likely to increase their body weight without changing the quality of the carcasses of

chickens (Donaldson et al., 1957; Combs and Nicholson, 1964; Saleh et al., 2004; Haščík et al., 2010). The growth promoters and feed additives in chicken's diets have been used for many years (USDA, 2008).

The bees are among the beneficial insects that produce mainly the honey, and also many by-products such as royal jelly, beeswax, propolis, pollen and bee stings. Bee pollen represents a rich source of proteins (25%), essential amino acids, oils (6%), containing more than 51% of polyunsaturated fatty acids of which 39% represent linolenic acid, 20% represent palmitic acid and 13% linoleic acid. Bee pollen also represents a source of more than 12 vitamins, 28 minerals, 11 enzymes or co-enzymes, 11 carbohydrates (35 - 61%; mainly glucose, fructose and sucrose), free amino acids, flavonoids, carotenoids and phytosterols (Crane, 1990; Abreu, 1992;

\*Corresponding author. E-mail: [alkrshola@yahoo.com](mailto:alkrshola@yahoo.com).

**Table 1.** Ingredients and nutrient composition of experimental feed mixture.

Ingredient (%)	Starter HYD-01 (1 to 21 days of age)	Grower HYD-02 (22 to 42 days of age)
Wheat	35.00	35.00
Maize	35.00	40.00
Soybean meal (48% N)	21.30	18.70
Fish meal (71% N)	3.80	2.00
Dried blood	1.25	1.25
Ground limestone	1.00	1.05
Monocalcium phosphate	1.00	0.70
Fodder salt	0.10	0.15
Sodium bicarbonate	0.15	0.20
Lysine	0.05	0.07
Methionine	0.15	0.22
Palm kernel oil Bergafat	0.70	0.16
Premix Euromix BR 0,5 %	0.50	0.50
<b>Nutrient composition (g.kg<sup>-1</sup>)</b>		
Crude protein	210.76	190.42
Fibre	30.19	29.93
Ash	24.24	19.94
Ca	8.16	7.28
P	6.76	5.71
Mg	1.41	1.36
Linoleic acid	13.51	14.19
ME <sub>N</sub> (MJ.kg <sup>-1</sup> ), calculated	12.02	12.03

Xu et al., 2009). There are great amount and variability of bee pollen phenolic constituents (total phenols, phenylpropanoids, flavonoids and anthocyanins) and its antioxidant activity (Broadhurst, 1999; Leja et al., 2007; Šaric et al., 2009). Its use in the human diet is very highly appreciated (Serra Bonvehi et al., 1991; Block et al., 1994).

Bee pollen has recently received an increased attention for its antibacterial (Garcia et al., 2001; Proestos et al., 2005; Carpes et al., 2007) and anti-fungicidal effects (Garcia et al., 2001). In mice, bee pollen constituents can be found in the blood, in cerebrospinal fluids, and in the urine within 2 h after ingestion (Markham and Campos, 1996). Diets supplemented with 1.5% bee pollen could boost the early development of thymus and *Fabrici bursa*, retard the bursa degeneration and promote the immune response of spleen chickens (Wang et al., 2005). Crane (1990) reported that in the liver and brain of mice, bee pollen affected several gene expressions that are important in the apoptosis pathway and chemiotaxis (Šaric et al., 2009).

One of the most widely used natural supplements is the bee pollen because it contains most of the essential nutritional elements needed for growth and development in humans and animals (Orzaez et al., 2002; Bell et al., 1983), and it could also promote the early development of

the digestive system, and therefore is a potentially beneficial food supplement (Wang et al., 2007). The aim of this study was to verify the influence of the addition of bee pollen extract in the feeding mixture diet of broiler chickens Ross 308. This effect will be considered in terms of changes in body weights of the chicken.

## MATERIALS AND METHODS

The experiment was implemented in test poultry station of Slovak University of Agriculture in Nitra. The tested chickens were Ross 308. The experiment included 180 one day-old chicks, which were divided into 2 groups: control (C) and experimental (E). Each of the group was also segregated according to the gender. The fattening duration was 42 days. The chickens were bred in a cage conditions; each cage was equipped with feed disperser and water intake was ensured *ad libitum* through a self feed-pump. The temperature was controlled during fattening period and it was 33°C at the first day and every week it was reduced about 2°C. The lighting during the feeding period was continuous. Each group was fed by same starter complete feed mixture (CFM) HYD-02 (loose structure) until to 21<sup>st</sup> day of their age. From the 22<sup>nd</sup> to 42<sup>nd</sup> day of their age, chickens were fed by grower fattening (CFM) HYD-02 (loose structure), in all investigated groups of experiments (Table 1.).

The feed mixture, HYD-01 and HYD-02, had been produced without antibiotic preparations and coccidiostatics. During the fattening, all groups were fed by the same complete feed mixture; however, to chickens of the experimental group, pollen extract in

**Table 2.** Effects of bee pollen on growth performance and yield of broiler Ross 308 (female).

Data	Live body weight (g)		Carcass weight (g)		Gilblets weight (g)		Carcass yield (%)	
	Group C	Group E	Group C	Group E	Group C	Group E	Group C	Group E
n	30	30	30	30	30	30	30	30
x	2246.60	2194.40	1573.20	1510.40	179.36	167.27	78.01	76.49
S.D.	115.02	73.79	83.63	29.913	15.81	15.46	1.56	1.52
Min.	2157.00	2115.00	1484.00	1488.00	161.89	143.89	76.40	74.40
Max.	2375.00	2285.00	1690.00	1556.00	197.01	186.19	79.88	78.46
CV%	5.12	3.36	5.32	1.98	8.82	9.24	2.00	1.99
SS	P ≥ 0.05		P ≥ 0.05		P ≥ 0.05		P ≥ 0.05	

n, Number of chicken; x, mean; S.D., standard deviation; CV, coefficient of variation; SS, statistical significance. P ≥ 0.05: not significant.

**Table 3.** Effects of bee pollen on growth performance and yield of broiler Ross 308 (male)

Data	Live body weight (g)		Carcass weight (g)		Gilblets weight (g)		Carcass yield (%)	
	Group C	Group E	Group C	Group E	Group C	Group E	Group C	Group E
n	30	30	30	30	30	30	30	30
x	2299.20	2354.60	1605.40	1646.40	168.53	172.62	77.17	77.25
S.D.	105.74	23.17	91.41	46.75	15.73	10.04	3.11	1.42
min.	2194.00	2330.00	1478.00	1581.00	150.52	160.77	74.45	75.25
max.	2472.00	2382.00	1698.00	1700.00	193.44	186.30	82.24	78.56
CV%	4.60	0.98a	5.69	2.84a	9.33	5.81	4.03	1.83
SS	P ≥ 0.05		P ≥ 0.05		P ≥ 0.05		P ≥ 0.05	

n, Number of chicken; x, mean; S.D, standard deviation; CV, coefficient of variation; SS, statistical significance. P ≥ 0.05: not significant.

amount 400 mg.kg<sup>-1</sup> were added to the feed mixtures (HYD-01 and HYD-02). The bee pollen extract was prepared from minced bee pollen (150 g) in the conditions of the 80% ethanol in the 500 cm<sup>3</sup> flask (Krell, 1996). Extraction was carried out in a water bath at 80°C for 1 h. Consequently, the extract was cooled and centrifuged. The obtained supernatant was evaporated in a rotary vacuum evaporator at bath temperature of 40 - 50°C and then weighed. At the end of the fattening (42 days), from each group were chosen 60 chickens for slaughter analysis (30 hens and 30 cocks) and then the meat performance of chickens was determined (slaughter weight, carcass weight, giblets weight, carcass yield). The experimental analysis was evaluated at the Department for Evaluation and Processing of Animal Products in the Faculty of Biotechnology and Food Sciences SPU Nitra, Slovakia. The results of meat performance (arithmetic mean, standard deviation, coefficient of variation) were processed by the statistic program Statgraphics 5.0. For the determination of significant differences between the tested groups, F-test was used followed by t-test.

## RESULTS AND DISCUSSION

In the present study, the results obtained showed that there were no significant differences (P ≥ 0.05) between the control and experimental groups. Table 2 shows the results for female group; the body weight (g) for control group (2246.60 g) was higher than experimental group (2194.40 g), the carcass weight (g) was higher in control group (1573.20 g) than experimental group (1510.40 g),

the giblets' weight(g) was higher in control group (179.36 g) than experimental group (167.27 g) and the carcass yield (%) was higher in control group (78.01%) than the experimental group (76.49%). Since the bee pollen stimulates the reproductive female hormones (Kolesarová et al., 2011), some energy are therefore channeled to the reproductive system.

Table 3 shows the results for male group; where the live body weight (g) was higher in experimental group (2354.60 g) than control group (2299.20 g), the carcass weight (g) was higher in experimental group (1646.40 g) than control group (1605.40 g), the giblets weight (g) in experimental group (172.62 g) was higher than control group (168.53 g) and carcass yield were higher in experimental group (77.25%) than control group (77.17%). The present results confirmed the reports of Angelovičová et al. (2010) who found that the body weight of experimental (1773.53 g) broiler Ross 308 was higher than in control group (1708.48 g) by about 65.05 g, with the addition of bee pollen (0.10%). Similar results were also reported by Wang et al. (2007) who found that the body weight of chickens fed with bee pollen addition (1585.67 ± 68.27 g) was higher than in control group (1173.33 ± 44.13 g). They also found that the size of the small intestine in experimental group was longer than in the control group.

The study by Attia et al. (2011) confirmed our results with rabbits fattening; they added bee pollen in different concentrates (T1-100, T2-200 and T3-300 g) to rabbits feed mixture and found that the body weight of the experimental group (T1-3094.00 ± 190.10, T2-2999.00 ± 175.20 and T4-3015.00 ± 155.00 g) was higher than the control group (2990.00 ± 186.40 g). Our study was also in agreement with that of Haro et al. (2000) who reported that at the end of their experiment, the group of rats fed with only bee pollen and water in the nutrition during 12 weeks was healthier with increased weight. On the contrary, Eman (2010) found that the different doses of bee pollens (2.5, 5 and 10 g. kg<sup>-1</sup> of body weight. day<sup>-1</sup>) as nutritional supplement for pregnant rats had harmful effect on the mother's and also to fetus' life.

## Conclusion

From the present results, it was concluded that the use of bee pollen as a dietary supplement in feed mixture of broiler Ross 308 in amount of 400 mg.kg<sup>-1</sup> led to an increase in the live body weight, carcass weight, giblet weight and carcass yield in males, but it had negative effect on the females, as it decreased the body weight of the hens.

## ACKNOWLEDGEMENT

This work was supported by KEGA 053 SPU-4/2011.

## REFERENCES

- Abreu M (1992). Food use of pollen in relation to human nutrition. *Alimentaria* 235:45-46.
- Adeyemo GO, Ologhobo AD, Adebisi OA (2010). The effect of graded levels of dietary methionine on the haematology and serum biochemistry of broilers. *Int. J. Poult. Sci.* 9:158-161.
- Angelovičová M, Štofán D, Močár K, Liptaiová D (2010). Biological effects of oilseed rape bee pollen and broiler's chickens performance. International conference on food innovation-foodinnova, 2010, Universidad politecnica de Valencia, Spain, pp. 246-247.
- Attia YA, AL-Hanoun A, Bovera F (2011). Effect of different levels of bee pollen on performance and blood profile of New Zealand White bucks and growth performance of their offspring during summer and winter months. *J. Anim. Physiol. Anim. Nutr.* 95:17-26.
- Bell RR, Thornber EJ, Seet JL, Groves MT, Ho NP, Bell DT (1983). Composition and protein quality of honey bee collected pollen of *Eucalyptus marginata* and *Eucalyptus calophylla*. *J. Nutr.* 113:2479-2484.
- Block G, Sinha R, Gridley G (1994). Collection of dietary-supplement data and implications for analysis. *Am. J. Clin. Nutr.* 59:232-239.
- Broadhurst CL (1999). Bee products: medicine from the hive. *Nutr. Sci.* 4:366-368.
- Carpes ST, Begnini R, De alencar SM, Masson ML (2007). Study of preparations of bee pollen extracts antioxidant and antibacterial activity. *Ciência e Agrotecnologia* 31:1818-1825.
- Combs GF, Nicholson JL (1964). Testing energy, amino acid and protein level specifications for linear programming of broiler rations. *Feed Stuffs* 36:17-19.
- Donaldson WE, Combs GF, Romoser GL, Supplee WC (1957). Studies on energy levels in poultry rations. 2. Tolerance of growing chicks to dietary fat. *Poult. Sci.* 36:807-815.
- Eman Abdel-gawad I (2010). Potential impact of bee pollen administration during pregnancy in rats. *J. Am. Sci.* 6:44-53.
- García M, Pérez arquillue C, Juan T, Juan MI, Herrera A (2001). A note: pollen analysis and antibacterial activity of Spanish honeys. *Food Sci. Technol. Int. Camb.* 7:55-158.
- Gueye EF (2009). The role of networks in information dissemination to family poultry farmers. *World's Poult. Sci. J.* 65:115-123.
- Haro A, Lopez-aliaga I, Lisbona F, Barrionuevo M, Alférez MJM, Campos MS (2000). Beneficial effect of pollen and/or propolis on the metabolism of iron, calcium, phosphorus, and magnesium in rats with nutritional ferropenic anemia. *J. Agric. Food Chem.* 48:5715-5722.
- Haščík P, Kačániová M, Mihok M, Pochop J, Benczová E (2010). Performance of various broiler chicken hybrids fed with commercially produced feed mixtures. *Int. J. Poult. Sci.* 9:1076-1082.
- Kolesárová A, Capcarová M, Baková Z, Gálik B, Juráček, M, Šimko M, Sirotkin A (2011). The effect of bee pollen on secretion activity, markers of proliferation and apoptosis of porcine ovarian granulosa cells *in vitro*. *J. Environ. Sci. Health Part B: Pesticides, Food Contaminants and Agricultural Wastes.* 46:207-212.
- Krell R (1996). Value-Added products from bee keeping. Milan, FAO Publications, p. 395. ISBN 92-5-103819-8.
- Leja M, Mareczek A, Wyzgolik G, Klepacz-baniak J, Czekonska K (2007). Antioxidative properties of bee pollen in selected plant species. *Food Chem.* 100:237-240.
- Markham KR, Campos M (1996). 7- and 8-O-methylherbacetin-3-O-sophorosides from bee pollens and some structure/activity observation. *Phytochemistry* 43:762-767.
- Orzaez villanueva MT, Diaz marquina A, Bravo serrano R, Blazquez abellán G (2002). The importance of bee-collected pollen in the diet: a study of its composition. *J. Food Sci. Nutr.* 53:217-224.
- Proestos C, Chorianopoulos N, Nichas GJE, Komaitis M (2005). RP-HPLC analysis of the phenolic compounds of plant extracts: investigation of their antioxidant capacity and antimicrobial activity. *J. Agric. Food Chem.* 53:1190-1195.
- Saleh EA, Watkins SE, Waldroup AL, Waldroup PW (2004). Effects of dietary nutrient density on performance and carcass quality of male broilers grown for further processing. *Int. J. Poult. Sci.* 3:1-10.
- Šarić A, Balog T, Sobocanec S, Kusic B, Sverko V, Rusak G, Likic S, Bubalo D, Pinto B, Reali D, Marott T (2009). Antioxidant effects of flavonoid from Croatian *Cystus incanus* L. rich bee pollen. *Food Chem. Toxicol.* 47:547-554.
- Serra bonvehí J, Escura pseudo F, Giner pallare's J (1991). Quantitative determination of free amino acids in honeybee collected pollen using gas chromatography and spectrophotometry. *Annales des Falsifications et de l'Expertise Chimique.* 897:153-166.
- USDA (2008). Additives in Meat and Poultry Products. [Showed.202-04-10]. Available at <[http://www.fsis.usda.gov/factsheets/Additives\\_in\\_Meat\\_&\\_Poultry\\_Products/index.asp](http://www.fsis.usda.gov/factsheets/Additives_in_Meat_&_Poultry_Products/index.asp)>.
- Wang J, Jin G, Zheng Y, Li S, Wang H (2005). Effect of bee pollen on development of immune organ of animal. *Zhongguo Zhong Yao Za Zhi.* 30:1532-1536.
- Wang J, Li S, Wang Q, Xin B, Wang H (2007). Trophic effect of bee pollen on small intestine in broiler chickens. *J. Med. Food.* 10:276-280.
- Xu X, Sun L, Dong J, Zhang H (2009). Breaking the cells of rape bee pollen and consecutive extraction of functional oil with supercritical carbon dioxide. *Innov. Food Sci. Emerg. Technol.* 10:42-46.