

Full Length Research Paper

Determination of effective nodulation in early juvenile soybean plants for genetic and biotechnology studies

E. T. Gwata^{1*}, D. S. Wofford¹, K. J. Boote¹, and H. Mushoriwa²

¹Department of Agronomy, University of Florida, P. O. Box 110300, Gainesville, FL – 32611 – 0300, USA.

²Crop Breeding Institute, P. O. Box CY 550, Causeway, Harare, Zimbabwe.

Accepted 6 October 2003

Symbiotic fixation of atmospheric nitrogen (N₂) is a complex physiological process influenced by the interaction of genetic elements in the higher plant species and rhizobia. No standardized, efficient method is available to critically examine the effect of altering the genetic elements in either component by selection and/or genetic engineering. At planting, seeds of a tropical ('TGX-4E') and non-tropical ('Soma') soybean cultivar were inoculated individually in sand-filled Conetainers® in a greenhouse with each of two strains within two rhizobial types (*Bradyrhizobium japonicum* and cowpea). Six weeks after inoculation, each plant was classified into one of two categories; vigorous plant with dark green leaves indicating effective nodulation and N₂-fixation (+), and stunted plant with chlorotic yellow leaves indicating ineffective nodulation and no N₂-fixation (-). The results indicated that this non-destructive method could be used to identify major genetic differences in the soybean and inoculant. Therefore, this method could be used to rapidly identify genetic segregants resulting from selection in plant breeding programs and/or genetic engineering.

Key words: effective nodulation, rhizobia, tropical soybean type, symbiosis.

INTRODUCTION

The formation of effective (functional) nodules in soybean [*Glycine max* (L.) Merr.] when inoculated with compatible rhizobia leads to fixation of atmospheric nitrogen (N₂) making nitrogenous fertilization of the soybean unnecessary. Symbiotic N₂-fixation is a complex physiological process influenced by the interaction of

genetic elements in the higher plant species and rhizobia.

Genetic variation for N₂-fixation ability has been reported involving both the soybean and rhizobium components of the symbiotic association (Hungria and Bohrer, 2000; Sanginga et al., 2000; Sinclair et al., 1991; Pulver et al., 1982). Currently, there is no reliable, simple and nondestructive method to identify N₂-fixation. Various methods, including acetylene reduction (Denison et al., 1983) and xylem ureide assay (McClure et al., 1980) that involve sophisticated and expensive equipment have been used for determining N₂-fixation. Destructive

*Corresponding author. E-mail: egwata@ufl.edu.

methods for the determination of shoot dry weight, which has a positive relationship to N₂-fixation ability (Neuhausen et al., 1988), have been reported.

This study reports a rapid, inexpensive and non-destructive method to screen soybean-rhizobium combinations which have improved N₂-fixation ability and have been generated by either plant breeding selection programs and/or genetic engineering.

MATERIALS AND METHODS

Genetic material

Pathogen-free seeds of two soybean cultivars ('Tropical G. max - 4E' = TGX -4E and non-tropical 'Soma') were obtained from the International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria and the Crop Breeding Institute (CBI), Harare, Zimbabwe, respectively.

Two rhizobial strains of *Bradyrhizobium japonicum* (designated I and II) and cowpea type (designated III and IV), were used in the study. The *B. japonicum* strains were obtained from a commercial outlet in Gainesville, Florida and the cowpea-type strains were obtained from Dr. P. Singleton at NifTAL, Hawaii, USA. Each strain was maintained in the laboratory by sub-culturing periodically (every 60 days) on yeast-mannitol agar (YMA) growth medium described by Vincent (1970). The cultures were streaked directly on YMA plates (15.0 x 100.0 mm) using the method described by Somasegaran and Hoben (1994) and incubated at room temperature.

Inoculum was prepared by excising (under sterile conditions in the laboratory) two pieces of agar (20 x 20 mm) supporting rhizobial colonies from the subculturing dishes that were then placed in bottles containing 300 ml sterile distilled water. Two drops of the surfactant Tween® 80 (polyoxyethylene sorbitan monooleate) were added to each bottle before shaking vigorously to disperse the rhizobial cells. Each bottle was wrapped with aluminum foil to protect rhizobia from inactivation by UV light.

Planting procedure and evaluation of N₂-fixation

The planting procedure involved placing one seed in a hole (1 cm deep) in the center of a Conetainer® (Stuewe and Son, Inc., USA) measuring 4 cm in diameter x 20.5 cm in length filled with moist sterile washed sand. After adding inoculum (0.1 ml), each seed was immediately covered with the sand. The resulting plants were grown in the greenhouse with day/night temperatures of 28/20°C without supplemental light in Gainesville, Florida, USA.

One week after emergence 0.1 g of microelement fertilizer (3.72% Fe, 9.28% Mg, 0.002% Mo and 2.32% Mn) was applied to each plant. Two weeks after emergence and at weekly intervals thereafter, 0.1 g of nitrogen-free (0-10-20) fertilizer was applied to each plant.

Six weeks after emergence, each plant was classified as either vigorous with dark green leaves indicating effective nodulation and N₂-fixation (+) or stunted with chlorotic yellow leaves indicating ineffective nodulation and no N₂-fixation (-). In each cultivar x strain combination, five plants (replications) were inoculated. One uninoculated (control) plant was included to assess effective nodulation.

Nodules from plants with green leaves and yellow leaves were harvested. Each nodule was washed and carefully rinsed with water before excision into halves in order to determine the color of the nodular tissue and relate it to the leaf color of the plant.

RESULTS AND DISCUSSION

After six weeks of growth under nitrogen-free conditions, all the uninoculated (control) plants were distinctly chlorotic, yellow and stunted. Rhizobial strain I induced effective nodules on both cultivars (Table 1). However, 'Soma' responded differentially to the strains. The cowpea-type rhizobial strain IV, was compatible with the cultivar TGX-4E but not with cultivar Soma (Table 1). This indicated that major cultivar and rhizobial differences were present and could be detected by this sensitive greenhouse conetainer method.

Table 1. The effectiveness of nodulation in early juvenile plants of two soybean cultivars grown in nitrogen-free medium inoculated individually with rhizobial strains. (+ = dark green leaves on normal, N₂-fixing plant; - = yellow leaves on chlorotic, nonfixing plant).

Cultivar	Rhizobial type			
	<i>B. japonicum</i> strain		Cowpea strain	
	I	II	III	IV
TGX-4E	+	+	+	+
Soma	+	+	-	-

Chlorotic plants with yellow leaves, were visually distinguishable from the vigorous plants with dark green leaves (Figure 1). Because the plants were grown in a nitrogen-free medium, the available nitrogen as indicated by the dark green leaves, was derived from the N₂-fixation process. Nitrogenous compounds resulting from N₂-fixation are exported from root nodules in the form of ureides (allantoin and allantoic acids) and translocated to the leaves where they are catabolized (Winkler et al. 1987) and used for the biosynthesis of chlorophyll and other proteins essential for photosynthesis.

All the inoculated plants had nodules regardless of cultivar or rhizobial strain. The morphology, size and exterior color of intact nodules were generally indistinguishable. However, the cross-sections of nodules indicated clear differences in the color of nodular tissue between effective and ineffective nodules (Figure 2). The ineffective nodules were white to light green inside while the effective nodules were characteristically pinkish-brown indicating differences in N₂-fixation capabilities between the two nodule types.

In summary, our results demonstrated a simple, inexpensive and reproducible method for identifying early juvenile soybean genotypes effective in N₂-fixation under nitrogen-free growth conditions. The method would be useful in plant breeding programs concerned with rapid screening of soybean for N₂-fixation effectiveness.



Figure 1. Differences in leaf color indicating differences in N_2 -fixation effectiveness among early juvenile soybean plants grown in nitrogen-free medium inoculated with a cowpea type rhizobial strain IV.




Figure 2. Intact and cross-section of root nodules from early juvenile soybean plants grown in a nitrogen-free medium and inoculated with a cowpea-type rhizobial strain IV. a = root nodules from plants with dark green leaves indicating effective nodulation and b = root nodules from plants with light yellow leaves indicating ineffective nodulation.

Moreover, the method is non-destructive and therefore allows for the development of advanced filial generations necessary for plant breeding, selection and genetic studies of variation in N_2 -fixation in soybean and possibly other leguminous species. In addition, this method can

also be useful in screening of transgenic plants as well as in biotechnology studies requiring identification of molecular markers using DNA extracted from early juvenile plant populations segregating for specific genetic traits such N_2 -fixation effectiveness.

ACKNOWLEDGEMENTS

For supplying soybean seed, we thank Mrs. P. N. Nyakanda and staff at CBI (Zimbabwe) and IITA (Nigeria). We are grateful to Dr. P. L. Pfahler for making improvements to this manuscript and Mr. E. Ostmak for technical assistance. We are grateful to the Rockefeller Foundation for financial support. (The use of trade names in this publication does not imply endorsement by the authors or Rockefeller Foundation of the products named, nor criticism of similar ones not mentioned).

REFERENCES

- Denison RF, Sinclair TR, Zobel RW, Johnson MN, Drake GM (1983). A non-destructive field assay for soybean nitrogen fixation by acetylene reduction. *Plant Soil* 70: 173-182.
- Hungria M, Bohrer TRJ (2000). Variability of nodulation and dinitrogen fixation capacity among soybean cultivars. *Biol. Fertil. Soils* 31: 45-52.
- McClure PR, Israel DW, Volk RJ (1980). Evaluation of the relative ureide content of xylem sap as an indicator of N₂ fixation. *Plant Physiol.* 66: 720-725.
- Neuhausen SL, Graham PH, Orf JH (1988). Genetic variation for dinitrogen fixation in soybean of maturity group 00 and 0. *Crop Sci.* 28: 769-772.
- Pulver EL, Bockman F, Wien HC (1982). Nodulation of soybean cultivars with *Rhizobium* spp. and their response to inoculation with *R. japonicum*. *Crop Sci.* 22:1065-1070.
- Sanginga N, Thottappilly G, Dashiell K (2000). Effectiveness of rhizobia nodulating recent promiscuous soybean selections in the moist savanna of Nigeria. *Soil Biol. Biochem.* 32: 127-133.
- Sinclair TR, Soffes AR, Hinson K, Albrecht SL, Pfahler PL (1991). Genotypic variation in soybean nodule number and weight. *Crop Sci.* 31:301-304.
- Somasegaran P, Boben HJ (1994). General microbiology of rhizobia In: Handbook for rhizobia methods in legume-rhizobium technology. Springer-Verlag, Berlin. pp 1-77.
- Vincent JM (1970). A manual for the practical study of root nodule bacteria. IBP Handbook No. 15. Oxford, Blackwell Scientific Publications, Oxford, England.
- Winkler RD, Blevins DG, Pollacco JC, Randall DD (1987). Ureide catabolism in soybeans. II. Pathway of catabolism in intact leaf tissue. *Plant Physiol.* 83: 585-591.