

Full Length Research Paper

# Behavioral response of resistant and sensitive *Pseudomonas aeruginosa* S22 isolated from Sohag Governorate, Egypt to cadmium stress

El-Sayed, M. Soltan<sup>1</sup>, Rehab, M. Mohamed<sup>1</sup> and Ahmed, A. Shoreit<sup>2\*</sup>

<sup>1</sup>Sohag University, Faculty of Science, Botany Department, Egypt.

<sup>2</sup>Assiut University, Faculty of Science, Botany Department, Egypt.

Accepted 21 March, 2008

A total of 62 *Pseudomonas* isolates were selected from a contaminated site at Sohag district and tested for their resistance to cadmium metal. Isolate no. 22 was found to be the most resistant one and has a plasmid with molecular wt 27.491 Kbp. The minimal inhibition concentrations (MIC) for the resistant (R) and adopted sensitive (S) isolates were 1.0 and 0.119 mmol L<sup>-1</sup>, respectively. The growth, protein content and maximum specific growth rate (k) decreased with increasing concentrations of heavy metal. The generation time (T) increased with the addition of divalent cations (Fe<sup>2+</sup>, Mg<sup>2+</sup> and Ca<sup>2+</sup>) plus cadmium. Under cadmium stress, three proteins of high molecular weights (208, 78 and 33.5 KD) were lost while one protein with low molecular weight (2.5 KD) was induced. LPS profile was modified under stress of CdCl<sub>2</sub>. Absence of bands in region I, which represents the O-specific antigen, and region II (core-oligosaccharide linked to lipid A) were recorded at high concentrations of CdCl<sub>2</sub>.

**Key words:** *Pseudomonas aeruginosa*, cadmium stress, heavy metal resistance.

## INTRODUCTION


The release of heavy metals into our environment is still large and causes an environmental pollution problem because of their unique characteristics (Banat et al., 2005). The sources of heavy-metal pollution are mining, milling and surface finishing industries, discharging a variety of toxic metals such as Cd, Cu, Ni, Co, Zn, and Pb into the environment (Malik, 2004). Our air, food and water often contain heavy metals (Nies, 2000). Heavy metal cations play an important role in many biochemical reactions due to their ability to form complex compounds (Roszbach et al., 2000). Toxicity occurs through the displacement of essential metals from their native binding sites, ligand interactions, alterations in the conformational structure of nucleic acids and proteins or interference with oxidative phosphorylation and osmotic balance (Bruins et al., 2000; Poole and Gadd, 1989).

A microorganism may survive toxic effects of heavy metals by means of a detoxification mechanism and these

are designated as resistant. However, the microorganism which has the ability to survive toxic effect of heavy metal because of intrinsic properties and/or environmental modification of toxicity are known as tolerant microorganism (Gadd, 1992). Resistance to heavy metals is observed in a wide variety of bacteria, especially in gram negative bacteria (Poole, 2002) such as *Pseudomonas* sp., *Ralstonia metallidurans*, *Enterobacter cloacae*, *Thiobacillus ferrooxidans*, and mucilage producing cyanobacteria (Malik, 2004; Ünalı et al., 2003, Piotrowska-Seget et al., 2005).

Bacteria of the genus *Pseudomonas* are well-studied and are of great interest not only because of their high resistance to heavy metals and other toxic substances, but also for their simple nutritional requirements and rapid growth on standard laboratory media (Pardo et al., 2003). *P. aeruginosa* was isolated from different heavy metal-polluted environments such as sewage, irrigation and agricultural drainage canals, and soil (Shoreit and Soltan, 1992; Soltan, 2001). Therefore, it has been considered as a water quality indicator organism. In the laboratory, *P. aeruginosa* shows resistance to high concentrations of heavy metals, for example, Zn, Cu, Ni, Pb, Cd and Hg

\*Corresponding author. E-mail: ashoreit@yahoo.com. Fax: +2 088 2342708.



**Figure 1.** A map showing the different places of Sohag Governorate from which soil and water samples were collected

(Wang et al., 1997; Teitzel and Parsek 2003; Hussein et al., 2005). This demonstrates that this strain could be a good candidate for heavy metal removal from polluted sites (Filali et al., 2000).

Heavy metal accumulation is markedly affected by the presence of other ions. Cations such as magnesium and calcium can often reduce heavy metal inhibition (Issa et al., 1998). Calcium is a strong competitor for cadmium and zinc binding (Plette et al., 1996). In addition, selective permeability and ion uptake through membranes is regulated by calcium; externally added calcium acts as a membrane stabilizer and performs this function even in heavy metal-treated cells (Pessoa-de-Franca et al., 2002).

The main objective of the present study is to select a resistant and sensitive *P. aeruginosa* strain from contaminated sites at Sohag district and study their growth parameters, morphology, and the behavioral response of protein and LPS profiles to cadmium stress.

## MATERIALS AND METHODS

### Collection of samples

Eighteen samples (9 sewage, 5 soils and 4 River Nile) from different places at Sohag Governorate (Egypt) (Figure 1) were collected and transferred immediately to the laboratory. The samples were stored at 4 - 5°C. Physicochemical characteristics of water and soil samples were carried out according to standard method (APHA, 1980).

### Screening for heavy metal toxicity

The isolates grown on CNA medium (Havelaar et al., 1985) were tested for heavy metals toxicity by using different concentrations (100 - 2000 ppm) of CdCl<sub>2</sub>.

### Toxic effect of CdCl<sub>2</sub> on isolate S22

Isolate S22 was tested to CdCl<sub>2</sub> by using the disk-method. The plates containing 20 ml of tris minimal agar were inoculated with 100 µl of previously grown culture in the same medium (O.D. ≈ 0.8) by pour plate method. The discs impregnated with 40 ppm CdCl<sub>2</sub> was applied to the surface and incubated for 48 h at 37°C.

### Plasmid isolation and curing

Isolation and curing of plasmid DNA were carried out according to Ghosh et al., 2000.

### Determination of minimal inhibitory concentrations (MICs)

The MICs were determined by allowing isolate 22 (R, S) to grow in tris minimal broth containing 0 - 40 ppm CdCl<sub>2</sub> for S strain and 0 - 200 ppm for R strains. It was incubated at 37°C in a rotatory shaker (150 rpm). Growth was observed after 12 h. The minimal inhibitory concentration expressed in mmol L<sup>-1</sup> is the concentration at which the growth was not observed (Mergeay, 1995; Yilmaz, 2003).

### Identification of isolate S22

The isolate no. 22 was identified on the basis of morphological, physiological and biochemical characteristics according to Bergey's manual 1994).

### Effect of CdCl<sub>2</sub> on the growth of *P. aeruginosa* S22

*P. aeruginosa* S22 (R strain) was grown in Tris minimal broth containing 0, 40, 80, 120, 160 and 200 ppm of CdCl<sub>2</sub>. The S-strain was grown in the same medium containing 0 and 40 ppm CdCl<sub>2</sub>. All treatments were agitated at 150 rpm and incubated at 37°C for 12 h. CdCl<sub>2</sub> was added after 3 h of incubation. Optical density at λ = 600 nm of each treatment were monitored using a Spekol 11 spectrophotometer every 3 h over 12 h of incubation (Nepple et al., 1999). Growth parameters were determined according to Mahapatra and Banerjee, 1996) after 12 h of incubation.

### Effect of divalent cations on growth of *P. aeruginosa* S22 (R-strain) in the presence of CdCl<sub>2</sub>

To study the effect of divalent cations on living cells of *P. aeruginosa* S22 (R-strain), it was grown in Tris minimal cations-free

**Table 1.** Physicochemical characteristics of soil samples collected from different sites at Sohag Governorate.

Sample No.	Physicochemical characteristics												
	Temp (°C)	pH	E.C (μhoms/cm)	OM (%)	NO <sub>3</sub> <sup>+</sup> (μg/L)	PO <sub>4</sub> <sup>3-</sup> (ppm)	Cl <sup>-</sup> (ppm)	K <sup>+</sup> (ppm)	Na <sup>+</sup> (ppm)	Pb <sup>2+</sup> (ppm)	Cu <sup>2+</sup> (ppm)	Cd <sup>2+</sup> (ppm)	Mg <sup>2+</sup> (ppm)
2	18	8.5	200	5.2	15	1.5	.06	45	285	0.5	0.09	0.1	32
7	18	8.2	220	6.8	35	1.5	0.03	35	200	0.5	0.09	0.1	35
17	20	8.5	200	8.2	41	1.65	0.028	28	187	0.4	0.08	0.02	26
5	18	7.8	170	10.6	59	1.8	0.029	20	137	0.43	0.05	0.03	28
6	18	7.7	170	14.5	82	2.1	0.025	18	126	0.2	0.04	0.05	31

WHO (2005) for cd=0.003mg/l

medium (control), or in presence of 40 ppm Cd<sup>2+</sup> only or in addition to each of Fe<sup>2+</sup>, Mg<sup>2+</sup>, or Ca<sup>2+</sup>. All treatments were incubated at 37°C with shaking at 150 rpm. Heavy metals were added after 3 h of incubation. Growth curve and growth parameters were measured as stated before.

#### Effect of CdCl<sub>2</sub> on morphology

The Effect of CdCl<sub>2</sub> on morphology was observed by using scanning and transmission electron microscope (SEM and TEM) according to Suh et al. (1998a, b).

#### Effect of CdCl<sub>2</sub> on protein content

The effect of CdCl<sub>2</sub> on protein content of *P. aeruginosa* S22 (R-strain) was detected according to the method of Lowery et al (1951). Protein and lipopolysaccharides (LPS) analysis was by polyacrylamide gel electrophoresis (SDS-PAGE; Laemmli, 1970).

## RESULTS AND DISCUSSION

### Physicochemical characteristics of *Pseudomonas* sp.

Tables (1 and 2) show the different physicochemical characteristics of the samples. The data indicate high contamination with Cd<sup>2+</sup> and Pb<sup>2+</sup> (Table 2). The bacterial counts from River Nile samples ranged from 10-45 for controls and 3 - 21

cfu/g dry wt or ml/ water for CdCl<sub>2</sub> (Table 3). The highest counts were obtained from sewage samples (215 - 780 cfu/ml) followed by soil samples (60 - 155 cfu/g dry wt), whereas, the lowest counts were obtained from the River Nile samples. This was explained by high total nitrogen and biological oxygen demand values of sewage samples (116 - 190 and 263 - 340 ppm, respectively). Our data show that the higher the heavy metal content of the samples, the lower the bacterial total counts (Figure 2). This is in agreement with Kandeler et al. (2000), Kozdroj and van Elsas (2001), Suhadolc et al. (2004) and Piotrowska-Seget et al. (2005) who reported significant decreases in microbial numbers, diversity and activity due to the toxic effects of high concentrations of metals. Soltan (2001) observed that the widespread occurrence of *P. aeruginosa* reflects its role as a contaminant, as an indicator of pollution and as a member of the bacterial community in a highly polluted area.

### Screening for heavy metal toxicity

Figure 2 shows that most isolates were resistant to CdCl<sub>2</sub> at low concentrations while few are resistant to higher concentrations up to 2000 ppm. These results are similar to that of Hassen et al. (1998) who reported that three strains of *P.*

*aeruginosa* (S6, S7 and S8) isolated from natural polluted environments on nutrient agar were tolerant to CuSO<sub>4</sub>, Cr<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> and CoSO<sub>4</sub> up to concentrations of 1.6, 1.2 and 0.8 mM, respectively. Similarly, Ünaldi et al. (2003) isolated strains of *Pseudomonas* sp. on glutamine-starch-phenol (GSP) agar which have tolerance to CuSO<sub>4</sub>, NiCl<sub>2</sub>, CdCl<sub>2</sub>, and K<sub>2</sub>Cr<sub>2</sub>O up to 5250, 2594.2, 2010 and 343.44 ppm, respectively.

Among the 62 *Pseudomonas* spp., isolate no. 22 (soil sample no.7) was the most heavy metal-resistant one. It was resistant to CdCl<sub>2</sub> (1800 - 2000 ppm) and was selected for further investigations.

### Isolation and curing of plasmid

A single plasmid was detected of about 27.491 kb. This suggests that the cadmium resistance of *Pseudomonas* 22 is plasmid mediated (Figure 3). This closely agrees with Malik and Jaiswal (2000). Ünaldi et al. (2003) reported that *Pseudomonas* strain WP19 which was resistant to Cu, Ni, Cd and Cr had 4 plasmids of approximately 20.8, 19.6, 8.0 and 4.7 kb. On the other hand, Malik et al. (2002) reported that the molecular size of the plasmids located in the bacterial strains isolated from agricultural and industrial soils which are re-

**Table 2.** Physicochemical characteristics of sewage and River Nile samples collected from different sites at Sohag Governorate.

Sample type	Sample No.	Physicochemical characteristics															
		Temp (°C)	pH	E.C (µhoms/cm)	TSS (mg/l)	TDS (mg/l)	TN (ppm)	BOD (mg/l)	NO <sub>3</sub> <sup>+</sup> (µg/L)	PO <sub>4</sub> <sup>3-</sup> (ppm)	Cl <sup>-</sup> (ppm)	K <sup>+</sup> (ppm)	Na <sup>+</sup> (ppm)	Pb <sup>2+</sup> (ppm)	Cu <sup>2+</sup> (ppm)	Cd <sup>2+</sup> (ppm)	Mg <sup>2+</sup> (ppm)
Sewage	3	17	8.1	220	959	304	116	263	103	2.2	80	35	65	1.2	0.28	0.1	55
	15	16	7.6	200	975	300	110	260	125	2.7	56	34	35	0.8	0.28	0.09	47
	13	18	8.2	200	976	311	121	266	168	3.2	58	34	46	0.7	0.2	0.07	49
	18	23	8.5	210	984	315	115	278	275	4.1	42	20	24	0.7	0.12	0.06	32
	14	25	8.1	200	965	312	136	280	315	5.7	47	21	67	0.6	0.1	0.06	54
	8	17	8.2	180	986	410	127	300	414	5.7	51	26	54	0.6	0.1	0.05	46
	10	16	7.8	190	1050	406	150	305	495	5.8	38	15	35	0.5	0.09	0.04	68
	9	26	8.0	160	1220	414	188	311	668	6.9	65	14	45	0.5	0.07	0.03	75
	1	23	8.1	150	1300	417	190	340	705	7.5	55	14	51	0.4	0.06	0.02	58
River Nile	11	18	7.5	180	215	109	3.0	4.0	2.7	0.3	22	14	77	0.4	0.06	0.04	23
	12	19	7.6	200	221	112	5.2	4.2	4.2	0.4	23	9	59	0.3	0.05	0.02	34
	16	17	8.4	190	283	200	6.5	4.5	3.4	0.4	25	8	56	0.2	0.04	0.02	32
	4	22	8.1	160	314	215	7.4	5.5	10.8	1.3	23	1.8	45	0.2	0.01	0.014	33

Temp = Temperature; E.C = electric conductivity; TSS = total suspended salts; TDS = total dissolved salts; TN = total nitrogen; BOD = biological oxygen demand.

sistant to Hg, Cd, Cu, Cr, Pb, Zn and Ni were in the range of 7 - 70 kb. Similarly, Mahapatra et al. (2002) reported that *Acidiphilium symbioticum* KM2 is highly resistant to several metals and harbors three plasmids of 3.8, 7.1, and 56 kb in size. Moreover, Piotrowska-Seget et al (2005) found that 10 bacterial isolates out of 28 isolated from arable soil contained the plasmid of roughly 20 kb in size

#### Toxic effects of CdCl<sub>2</sub> on isolate S22 and its plasmid-cured strain

To confirm the previous results, the toxic effect of CdCl<sub>2</sub> (40 ppm) on the two strains of isolate 22 (plasmid-containing and plasmid-cured) was obtained. Figure 4 shows that the plasmid-cured

strain formed a clear zone while the former was not affected. Thus CdCl<sub>2</sub> resistant (R) and sensitive (S) strains were obtained. Sabry et al. (1997) indicated that the resistant strains are those which were not inhibited by 1 mmol L<sup>-1</sup> Cd. Mahapatra et al. (2002) reported that *Acidiphilium symbioticum* KM2 is highly resistant to several metals and harbors several plasmids. It becomes extremely sensitive to metals when it is cured of its plasmids.

#### Growth of *P. aeruginosa* S22 (R, S) under CdCl<sub>2</sub> stress

Figure 5 and Table 4 Show that optical density and maximum specific growth rate (k) for R, S strains decreased with increasing the concentra-


tions of CdCl<sub>2</sub>, while the generation time (T) increased. Cd<sup>2+</sup> had a potent inhibitory action on growth. Mahapatra and Banerjee (1996) reported that ZnSO<sub>4</sub>, CdSO<sub>4</sub>, CuSO<sub>4</sub> and NiSO<sub>4</sub> decreased the growth rate (k) and increased the generation time (T) of different species of *Acidocella facilis*.

#### Determination of MICs of CdCl<sub>2</sub> for isolate S22 (R, S)

The obtained data revealed that the MIC for isolate no 22 R and S strains were 1.0 and 0.119 mmol L<sup>-1</sup>, respectively. Wang et al. (1997) isolated a cadmium-resistant *P. aeruginosa* which tolerated cadmium up to 5 mM in artificial sea water medium. Hassen et al. (1998) and Hussein et al. (2005) isolated a Cd-resistant *P. aeruginosa* with


**Table 3.** Total counts of *Pseudomonas* sp. of different samples collected from different contaminated sites at Sohag district on CNA medium containing 40 ppm of CdCl<sub>2</sub>.

Sample type	Sample no.	Heavy metal concentrations (cfu /g ry wt)	
		Control	CdCl <sub>2</sub> (40 ppm)
Soil	2	60	47
	7	72	54
	17	120	89
	5	145	107
	6	155	111
	Sewage	3	215
15		235	128
13		260	156
18		330	213
14		335	224
8		350	243
10		355	224
9		400	320
River Nile	11	10	3
	12	20	9
	16	32	18
	4	45	21




**Figure 2.** The percentage of the resistant *Pseudomonas* isolates to CdCl<sub>2</sub> at low (100-700), moderate (800-1400) and high (1500-2000) concentration levels out of 18, 32 and 12 isolates isolated from soil, sewage and River Nile, respectively.


MIC of 1.5 mM in nutrient broth. On the other hand, Malik (2004) reported that *Pseudomonas* strain H1 isolated from soil was resistant to 225 ppm Cd<sup>2+</sup>. Moreover, Piotrowska-Seget et al. (2005) reported that the MICs of Pb, Cd, Cu and Zn for a soil-borne *Pseudomonas gla-*



**Figure 3.** Lambda DNA cleaved with HindIII (A, molecular weight markers), the absence of plasmid in cured "sensitive" cells (B), and presence of plasmid in "resistant cells" (C) of *P. aeruginosa* S22 isolate.



**Figure 4.** Toxic effect of CdCl<sub>2</sub> on isolate S22 (A) and the plasmid-cured strain (B) grown on tris minimal agar medium by using disc-method. The discs were impregnated with 40 ppm CdCl<sub>2</sub>.




**Figure 5.** Growth curves of *P. aeruginosa* S22 grown in Tris minimal broth containing 0 ppm (●), 40 ppm CdCl<sub>2</sub> (◆) for S-strain (A) ; 0 ppm (●), 40 ppm (◆), 80 ppm (▲), 120 (■), 160 ppm (▼) and 200 ppm (◆) of CdCl<sub>2</sub> (A'), for R-strain incubated at 37 C with agitation at 150 rpm for 12 h. CdCl<sub>2</sub> was added after 3 h of incubation (arrow).

**Table 4.** Growth parameters of *Pseudomonas aeruginosa* S22 (R, S) grown in Tris minimal broth containing 0, 40 ppm of CdCl<sub>2</sub> for S-strain and 0, 40, 80, 120, 160 and 200 ppm for R-strain of CdCl<sub>2</sub> agitated at 150 rpm and incubated at 37°C for 12 h. Heavy metals were added after 3 h of incubation.

CdCl <sub>2</sub> conc. (ppm)	T <sub>λ</sub> (h)	T <sub>α</sub> (h)	a R (S)	k R (S)	T (h) R (S)
Control	1.5	9.0	1.61	0.117	3.45
<b>CdCl<sub>2</sub></b>					
40	1.5	9.0	1.57 (0.60)	0.086 (0.01)	3.9 (>12)
80	1.5	9.0	1.52	0.085	4.35
120	1.5	9.0	1.23	0.07	5.4
160	1.5	9.0	1.16	0.04	8.75
200	1.5	9.0	1.01	0.03	12.0

T<sub>λ</sub> (h) = Lag period; T<sub>α</sub>(h) = the end of the exponential phase; a = asymptotic level = ln (OD<sub>α</sub>/OD<sub>0</sub>); k = the maximum specific growth rate = a / T<sub>α</sub> - T<sub>0</sub>; T = generation time; and R = resistant strain.



**Figure 6.** Effect of divalent cations (Ca<sup>2+</sup>, Mg<sup>2+</sup> and Fe<sup>2+</sup>) on *P. aeruginosa* S22 (R-strain) grown in tris minimal medium containing 40 ppm Cd<sup>2+</sup> in addition to 40 ppm of each of Mg<sup>2+</sup>, Fe<sup>2+</sup> or Ca<sup>2+</sup> and incubated at 37°C with agitation at 150 rpm for 12 h. Time at which Cd<sup>2+</sup> and divalent cations were added (3 h).

*diolus* were 0.1, 3, 5 and 10 mmol<sup>-1</sup>, respectively.

### Identification of isolate S22

Isolate was identified as *P. aeruginosa* S22 based on phenotypic methods. Shoreit and Soltan (1992) isolated 394 isolates on selective isolation media from different soil and water sites at Sohag district and they were

identified as *Pseudomonas*; 9% of them were identified as *P. aeruginosa*. Soltan (2001) isolated 240 strains of *P. aeruginosa* from a wastewater treatment plant, agricultural drainage canals, the River Nile and irrigation canals at Sohag area 7.15, 12.9, 25.4, and 53.7% of the isolates were resistant to lead, cadmium, mercury and zinc, respectively.

### Effect of divalent cations on growth of *P. aeruginosa* S22 in the presence of Cd<sup>2+</sup>

Figure 6 and Table 5 depict that growth and the maximum specific growth rate (k) increased by addition Fe<sup>2+</sup>, Mg<sup>2+</sup> and Ca<sup>2+</sup>, while the generation time (T) decreased compared to Cd<sup>2+</sup> only. These results are closely related to the data obtained by Adam et al. (1997) who observed that Ca<sup>2+</sup> ions increased cell multiplication of *Scenedesmus obliquus* in comparison with heavy metal treatment alone. Also, magnesium found to reduce toxicity of nickel to bacteria and yeast (Hettiarachchi et al., 2000; Sandrin and Maier, 2003).

### The effect of cadmium on morphology


Scanning and transmission electron micrographs (Figure 7) of *P. aeruginosa* S22 cells grown in tris minimal broth containing 120 ppm CdCl<sub>2</sub> show malformations such as filament formation where some cells are attached together (Figure 7B) in addition to roughened cell surface (Figure 7C) in comparison with control (Figure 7A). Ultrastructural modifications such as thinner and ruptured cell walls and large intracellular vacuoles appeared within the cells (Figure 7F).

Filament formation by *P. aeruginosa* S22 under cadmium stress may be a result of cadmium interference with the division mechanisms. This agrees with Nepple et al. (1999) who suggested that heavy metals act strongly on cell division. Morita (1997) reported that cell surfaces of cultures treated with cadmium chloride tended to be rough, suggesting that the cell increased its surface to

**Table 5.** Effect of divalent cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Fe}^{2+}$ ) on the growth parameters of *P. aeruginosa* S22 (R-strain) grown in tris minimal broth containing 40 ppm  $\text{Cd}^{2+}$  in addition to 40 ppm of each of  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$  or  $\text{Ca}^{2+}$ , incubated at 37°C with agitation at 150 rpm for 12 h.

Conc. of $\text{Cd}^{2+}$ and divalent cations (40 ppm of each)	$T_{\lambda}$ (h)	$T_{\alpha}$ (h)	a	K	T (h)
$\text{Cd}^{2+} + \text{Ca}^{2+}$	1.5	9	1.84	0.30	3.4
$\text{Cd}^{2+} + \text{Mg}^{2+}$	1.5	9	1.75	0.29	3.57
$\text{Cd}^{2+} + \text{Fe}^{2+}$	1.5	9	1.64	0.27	3.8
$\text{Cd}^{2+}$	1.5	9	1.61	0.26	3.87
Control	1.5	9	1.89	0.32	3.3

$T_{\lambda}$  (h) = Lag period;  $T_{\alpha}$ (h) = the end of the exponential phase; a = asymptotic level =  $\ln(\text{OD}_{\alpha}/\text{OD}_0)$ ; k = the maximum specific growth rate =  $a / T_{\alpha} - T_0$ ; T = generation time; and R = resistant strain.




**Figure 7.** Scanning (A, B, C) and transmission (D, E, F) electron micrographs of *P. aeruginosa* S22 grown in Tris minimal broth containing 0 (control) (A, D) and 120 ppm  $\text{CdCl}_2$  (B, C, E and F) agitated at 150 rpm and incubated at 37°C (Bar = 1 $\mu$ ).

improve the interaction of toxic substances with the cell surface.


#### Effect of $\text{CdCl}_2$ on protein content

Figure 8 shows that different concentrations of  $\text{CdCl}_2$



**Figure 8.** Toxic effect of  $\text{CdCl}_2$  on the total protein content of *P. aeruginosa* S22 grown in Tris minimal broth containing 0, 40, 80, 120, 160 and 200 ppm of  $\text{CdCl}_2$ , agitated at 150 rpm and incubated at 37°C for 12 h.

inhibit protein synthesis of *P. aeruginosa* S22 compared to the respective control. The protein inhibition ratio caused by  $\text{CdCl}_2$  reached about 27, 36, 48, 56 and 62% at 40, 80, 120, 160 and 200 ppm  $\text{CdCl}_2$ , respectively. Our obtained data indicate that  $\text{Cd}^{2+}$  decreases the total protein content of *P. aeruginosa* S22. Fahmy (2001) stated a decrease in the total protein content of two strains of photosynthetic bacteria *Rhodobacter capsulatus* B10 (capsulated strain) and 37b4 (non-capsulated strain) treated with different concentrations of  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$ . Lima Ana et al. (2005) reported a drastic decrease in total soluble protein content of *Rhizobium leguminosarum* bv. *Viciae* treated with high concentrations of  $\text{Cd}^{2+}$  which suggests that intracellular  $\text{Cd}^{2+}$  imposed on cellular metabolism.



**Figure 9.** Soluble protein patterns (Coomassie blue stained 12% SDS-PAGE) of whole cell lysates extracted with lysing buffer (see materials and methods) of S-strain (lane 2) and R-strain (lanes 3 - 9) of *P. aeruginosa* S22 grown in Tris minimal broth containing 0, 40, 80, 120, 160 and 200 ppm and incubated with shaking at 150 rpm at 37°C for 12 h. Lane 1: migration distance of molecular mass standards (in kilodaltons) are indicated in the left; lanes 2, 3 : 0 ppm CdCl<sub>2</sub> of both strains; lanes 4, 5, 6, 7 and 8 : 40, 80, 120, 160 and 200 ppm CdCl<sub>2</sub>, respectively, of resistant strain. Arrowheads point at some of the new bands induced during stress of CdCl<sub>2</sub>. Star (\*) point at some bands which disappeared at 120, 160 and 200 ppm CdCl<sub>2</sub> but appeared in control

### Protein profiles of *P. aeruginosa* S22 under Cd<sup>2+</sup> stress

The obtained data in Figure 9 and Table 6 reveals an alteration of protein profile when *P. aeruginosa* S22 was treated with different concentrations of CdCl<sub>2</sub>. Three proteins of high molecular weights (208, 78 and 33.5 kDa) were lost and one protein with low molecular weight (2.5 kDa) was induced as a response of Cd<sup>2+</sup> stress. These findings are closely related to the data obtained by Mallick and Rai (1998), who detected production of a 3.3 kDa protein, rich in cadmium and -SH contents in the cyanobacterium *Anabaena doliolum* following 20 mM Cd exposure. They also observed that Cd is a strong inducer of the metal-binding proteins. On the other hand, Hamer (1986) reported that the synthesis of metal-binding proteins with low molecular weight (6 - 7 kDa) and rich with cysteine is well-known in bacteria. Fahmy (2001) also reported the induction of new sets of proteins (156, 75, 119 and 105 kDa) in *Rhodobacter capsulatus* strain B10 and 37b4, while a 67 kDa protein disappeared in both strains under cadmium stress. Similarly, Valerie and Gordon (1994) reported that the production of a copper-induced protein has been implicated in copper detoxifi-


cation in a wild-type *Vibrio alginoliticus*. Laplace et al. (1996) reported induction of new proteins by *Enterococcus faecalis* treated with high concentrations of Cd<sup>2+</sup>. Aiu et al. (1999) found that two Cobalt-resistant strains of *Pseudomonas* [*P. sp.* BS501 and *P. putida* BS394 (pBS501)] induced protective surface proteins of the cell wall with the molecular weights of 49, 40, and 32 kDa under the stress of cobalt. Moreover, Nies (1992) observed that 116-kDa CzcA, 55-kDa CzcB and 37-kDa Czc proteins were induced by the gram-negative bacterium *Alcaligenes eutrophus* under cadmium stress.

### Lipopolysaccharide (LPS) profiles of *P. aeruginosa* S22 under CdCl<sub>2</sub> stress

The data (Figure 10) revealed that the LPS profiles were modified under stress of CdCl<sub>2</sub>. Absence of bands in region I which represents the O-specific antigen and region II (core-oligosaccharide linked to lipid A) occurred at high concentrations of CdCl<sub>2</sub>. In this respect, Lins and Straatsma (2001) stated that lipopolysaccharides (LPS) form the major constituent of the outer membrane of Gram-negative bacteria, and are believed to play a key

**Table 6.** Molecular weights (Mr.wt.) and percentage amounts of protein bands of untreated S-strain of *Pseudomonas aeruginosa* S22 and R, strain grown in tris minimal broth containing 0, 40,80, 120, 160 and 200 ppmCdCl<sub>2</sub>, both strains were incubated at 37°Cwith agitation at 150 rpm for 12 h.

Band no.	Marker Mr. wt. (Kda)	Mr. wt. (Kda)	Amount of protein (%)						
			S-strain Untreated (Lane 2)	R-strain					
				Cont. (Lane 3)	40 ppm (Lane 4)	80 ppm (Lane 5)	120 ppm (Lane 6)	160 ppm (Lane 7)	200 ppm (Lane 8)
1		306	----	2.6	3.0	2.4	2.2	4.5	4.3
2		252	----	2.3	3.3	4.6	3.2	2.5	3.8
3	230	230	1.8	3.0	2.9	4.5	4.4	4.3	5.3
4		208	2.8	----	----	----	----	----	----
5	135	135	----	2.9	3.6	3.5	3.2	3.0	3.5
6		115	3.8	2.0	3.1	2.9	3.0	3.8	4.3
7	97	97	5.2	3.3	4.6	6.0	4.5	2.5	3.7
8		85	4.4	2.9	3.0	4.4	4.7	4.8	5.6
9	78	78	2.8	3.1	3.7	3.4	----	----	----
10	57.5	57.5	3.1	2.0	4.5	4.6	4.2	4.5	5.1
11	38.5	38.5	3.8	3.9	3.0	2.7	4.4	3.0	2.3
12	33.5	33.5	3.3	8.0	8.1	4.4	----	----	----
13		19	3.8	3.4	2.1	2.7	3.0	3.2	4.6
14		16	5.0	3.1	3.7	3.9	3.4	1.6	3.2
15		12	8.6	5.2	5.0	4.7	3.3	2.9	5.7
16		5	4.0	5.1	4.6	5.1	4.7	4.2	5.5
17		2.5	----	----	----	----	5.7	5.9	3.9



**Figure 10.** LPS profile in 12% SDS-PAGE of whole cell lysates extracted with lysing buffer and subjected to protinase K digestion (see materials and methods) of *Pseudomonas aeruginosa* S22 (untreated S-strain Lane 1) and R-strain (Lanes 2-7) CdCl<sub>2</sub>. Silver strained protinase K resistant bands (controls and treated are indicated by arrows (region I and II). T he absence of arrows indicated the removal of bands under stress of CdCl<sub>2</sub>.

role in processes that govern microbial metal binding, microbial adsorption to mineral surfaces, and microbe-mediated oxidation/reduction reactions at the bacterial exterior surface. Arredondo et al. (1994) concluded that raising the pH of the solution containing the cells of *Thiobacillus ferrooxidans* from 1.5 to 8.0 releases about

50% of LPS. On the other hand, Fahmy (2001) found that region I (O-specific antigen) and region II (core-oligosaccharide linked to lipid A) of *Rhodobacter capsulatus* strain B10 (capsulated strain) and strain 37b4 (non-capsulated strain) completely disappeared under cadmium stress, whereas, partial removal of region I was missing in non-capsulated strain under the same stress of heavy metal. Moreover, Poole (2002) reported that LPS mutations were associated with antibiotic treatment in *P. aeruginosa*.

## Conclusion

The results obtained in this study indicate that this isolated *P. aeruginosa* S22 could be a good candidate for heavy metal removal from polluted environments.

## REFERENCES

- Adam MS, Abdel-Baset R, Issa A (1997). Protective effect of Ca<sup>2+</sup> ions on wall polysaccharides of *Scenedesmus obliquus* (Turp.) Kutz. Cells treated with heavy metals. Acta Hydrobiol. 39: 61-65.
- Aiu I, Gavriushkin AV, Siunova TV, Khasanova LA, Khasanova ZM (1999). Resistance of certain strains of *Pseudomonas* bacteria to toxic effect of heavy metal ions. Mikrobiologiya. 68: 366-74.
- American Public Health Association (APHA) (1980). Standard methods for the examination of water and wastewater. 15th edition, Washington, D. C.
- Arredondo R, García A, Jerezv CA (1994). Partial removal of polysac-

- charide from *Thiobacillus ferrooxidans* affects its adhesion to solids. *Appl. Environ. Microbiol.* 60: 2846-2851.
- Banat K, Howari F, Al-Hamada A (2005). Heavy metals in urban soils of central Jordan: Should we worry about their environmental risks? *Environ. Res.* 97: 258-273.
- Bergey's manual of determinative bacteriology (1994). Eds Holt JG, Kreig NR, Sneath PHA, Staley JT, Williams ST, 9th edn Williams and Wilkins, Baltimore, Maryland 21202 (USA), ISBN 0-68300603-7.
- Bruins M, Kapil S, Oehme F (2000). Microbial Resistance to Metals in the Environment. *Ecotoxicol. Environ. Saf.* 45: 198-207.
- Fahmy SM (2001). The behavioral responses of *Rhodobacter capsulatus* B10 and 37b4 to cadmium and zinc toxicity. MSc. thesis. Faculty of science. Assiut University. Assiut-Egypt.
- Filali BK, Taoufik J, Zeroual Y, Dzairi FZ, Talbi M, Blaghen M (2000). Wastewater bacterial isolates resistant to heavy metals and antibiotics. *Curr. Microbiol.* 41: 151-156.
- Gadd GM (1992). Heavy metals pollutants: Environmental and biotechnological aspects. *Encycl. Microbiol.* Volume 2.
- Ghosh S, Mahapatra NR, Ramamurthy T, Banerjee PC (2000). Plasmid curing from an acidophilic bacterium of the genus *Acidocella*. *FEMS Microbiol. Lett.* 183: 271-274.
- Hamer DH (1986). Metallothionein. *Annu. Rev. Biochem. Sci.* 55: 913-951.
- Hassen A, Saidi N, Cherif M, Boudabous A (1998). Resistance of environmental bacteria to heavy metals. *Bioresour. Technol.* 64: 7-15.
- Havelaar AH, During M, Delfgou-Van Ash EHM (1985). Comparative study of membrane filtration and enrichment media for the isolation and enumeration of *Pseudomonas aeruginosa* from sewage, surface water, and swimming pools. *Can. J. Microbiol.* 31: 686-692.
- Hettiarachchi GM, Pierzynski GM, Ransom MD (2000). *In situ* stabilization of soil lead using phosphorus and manganese oxide. *Environ. Sci. Technol.* 34: 4614-4619.
- Hussein H, Farag S, Kandil K, Moawad H (2005). Tolerance and uptake of heavy metals by *Pseudomonads*. *Process Biochem.* 40: 955-961.
- Issa AA, Adam MS, Abdel-Baset R (1998). Effect of  $Ca^{2+}$  on the detoxification of  $Cd^{2+}$  by *Scenedesmus obliquus* cells at low or high temperature. *Folia Microbiol.* 43: 645-648.
- Kandeler E, Tschirko D, Bruce KD, Stemmer D, Hobbs PJ, Bardgett RD, Amelung W (2000). Structure and function of the soil microbial community in microhabitats of a heavy metal polluted soil. *Biol. Fert. Soils.* 32: 390-400.
- Kozdroj J, van Elsas JD (2001). Structural diversity of microorganisms in chemically perturbed soil assessed by molecular and cytochemical approaches. *J. Microbiol. Method.* 43: 197-212.
- Laemmli VK (1970). Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature.* 227: 680-685.
- Laplace J, Boutibonnes P, Auffray Y (1996). Unusual resistance and acquired tolerance to cadmium chloride in *Enterococcus faecalis*. *J. Basic Microbiol.* 5: 311-317.
- Lima Ana G, Etelvina M, Sofia PA (2005). Cadmium tolerance plasticity in *Rhizobium leguminosarum* bv. *Vicia*: glutathione as a detoxifying agent. *Can. J. Microbiol.* 51: 7-14.
- Lins RD, Straatsma TP (2001). Computer Simulation of the Rough Lipopolysaccharide Membrane of *Pseudomonas aeruginosa*. *Biophys. J.* 81: 1037-1046.
- Lowery OH, Rosebrough NH, Farr AL, Randall RJ (1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193: 291-297.
- Mahapatra NR, Ghosh S, Deb C, Banerjee PC (2002). Resistance to cadmium and zinc in *Acidiphilium symbioticum* KM2 is plasmid mediated. *Curr. Microbiol.* 45: 180-186.
- Mahapatra NR, Banerjee PC (1996). Extreme tolerance to cadmium and high resistance to copper, nickel and zinc in different *Acidiphilium* strains. *Lett. Appl. Microbiol.* 23: 393-397.
- Malik A, Jaiswal R (2000). Metal resistance in *Pseudomonas* strains isolated from soil treated with industrial wastewater. *World J. Microbiol. Biotechnol.* 16: 177-182.
- Malik A (2004). Metal bioremediation through growing cells. *Environ. Int.* 30: 261-278.
- Malik A, Sakamoto M, Kakii K (2002). Coaggregation of *Microbacterium esteraromaticum* S51 with other strains of non-flocculating sludge bacteria. *IWA's Water Environ. Manage. Ser.* 737-748.
- Mallick N, Rai LC (1998). Characterization of Cd-induced low molecular weight protein in a N2-fixing cyanobacterium *Anabaena doliolum* with special reference to co-/multiple tolerance. *BioMetals.* 11: 55-61.
- Mergey M (1995). Heavy metal resistance in microbial ecosystems. *Mole. Microbiol. Ecol. Mann.* 6: 7-17.
- Morita RY (1997). Bacteria in oligotrophic environment: starvation-survival lifestyle. pp. 286-287. Chapman and Hall. New York.
- Nepple BB, Flynn I, Bachofen R (1999). Morphological changes in phototrophic bacteria induced by metalloid oxyanions. *Microbiol. Res.* 154: 191-198.
- Nies DH (1992). CzcR and CzcD, gene products affecting regulation of resistance to cobalt, zinc and cadmium (czc system) in *Alcaligenes eutrophus*. *J. Bacteriol.* 174: 8102-8110.
- Nies DH (2000). Heavy-metal resistant bacteria as extremophiles: molecular physiology and biotechnological use of *Ralstonia* sp. CH34. *Extremophiles* 4: 77-82.
- Pardo R, Herguedas M, Barrado E, Vega M (2003). Biosorption of cadmium, copper, lead and zinc by inactive biomass of *Pseudomonas putida*. *Anal. Bioanal. Chem.* 376: 26-32.
- Pessoa-de-Franca F, Mora-Tavares A, Augusto-da-Costa A (2002). Calcium interference with continuous biosorption of zinc by *Sargassum* sp. (Phaeophyceae) in tubular laboratory reactors. *Biores. Technol.* 83: 159-163.
- Piotrowska-Seget Z, Cycon M, Kozdroj J (2005). Metal-tolerant bacteria occurring in heavily polluted soil and mine spoil. *Appl. Soil Ecol.* 28: 237-246.
- Plette A, Benedetti M, Van Rimsdijk (1996). Comparative binding of protons, calcium, cadmium and zinc to isolated cell walls of a Gram-positive soil bacteria. *Environ. Sci. Technol.* 30: 1902-1910.
- Poole K (2002). Outer membranes and efflux: The Path to Multidrug Resistance in Gram- Negative Bacteria. *Curr. Pharm. Biotechnol.* 3: 77-98.
- Poole RK, Gadd GM (1989). Metals: Microbe Interactions, pp. 137. IRL Press, Oxford. Programme on Chemical Safety (Concise International Chemical Assessment Document 12).
- Rosbach S, Kukuk ML, Wilson TL, Feng SF, Pearson MM, Fisher MA (2000). Cadmium-regulated gene fusions in *Pseudomonas fluorescens* *Environ. Microbiol.* 2: 373-382.
- Sabry SA, Ghozlan HA, Abou-zeid DM (1997). Metal tolerance and antibiotic resistance patterns of bacterial population isolated from sea water. *J. Appl. Microbiol.* 82: 245-252.
- Sandrin TR, Maier M (2003). Impact of Metals on the Biodegradation of Organic Pollutants. *Environ. Health. Perspect.* 111: 1093-1101.
- Shoreit A, Soltan E (1992). Fluorescent and non-fluorescent *Pseudomonas* species from Sohag Governorate (Upper Egypt). *Bull. Fac. Sci., Assiut Univ.* 21: 133-143.
- Soltan EM (2001). Isolation and Characterization of antibiotic and heavy metal resistant *Pseudomonas aeruginosa* from different polluted waters in Sohag District, Egypt. *Microbiol. Biotechnol.* 11: 50-55.
- Suh JH, Kim DS, Yun JW, Song SK (1998a). Process of  $Pb^{2+}$  accumulation in *S. cervisiae*. *Biotechnol. Lett.* 20: 153-156.
- Suh JH, Yun JW, Kim DS (1998b). Comparison of  $Pb$  accumulation characteristics between live and dead cells of *S. cervisiae* and *Aureobasidium pullulans*. *Biotechnol. Lett.* 20: 247-251.
- Suhadolc M, Schroll R, Gatterer A, Schlöter M, Munch JC, Lestan D (2004). Effects of modified  $Pb$ -,  $Zn$ -, and  $Cd$ - availability on the microbial communities and on the degradation of isoproturon in a heavy metal contaminated soil. *Soil Biol. Biochem.* 36: 1943-1954.
- Teitzel GM, Parsek MR (2003). Heavy Metal Resistance of Biofilm and Planktonic *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 69: 2313-2320.
- Ünalı MN, Korkmaz H, Arıkan B, Coral G (2003). Plasmid-Encoded heavy metal resistance in *Pseudomonas* sp. *Bull. Environ. Contam. Toxicol.* 71: 1145-1150.
- Valerie JH, Gordon AS (1994). Regulation of extracellular copper-binding proteins in copper-resistant and copper sensitive mutants of *Vibrio alginolyticus*. *Appl. Environ. Microbiol.* 60: 1749-1753.
- Wang CL, Michels, Schott PC, Sawson C, Kitisakul S, Baross JA, Keasling JD, Clark DS (1997). Cadmium removal by a new strain of *Pseudomonas aeruginosa* in aerobic culture. *Appl. Environ. Microbiol.* 63: 4075-4078.

- WHO (2005). Guidelines for drinking water Quality Vol I, Geneva, Switzerland.
- Yilmaz EI (2003). Metal tolerance and biosorption capacity of *Bacillus circulans* strain EB1. Res. Microbiol. 154: 409-415.