

Full Length Research Paper

Growth and physiological responses to water and nutrient stress in oil palm

Cheng-xu Sun¹, Hong-xing Cao^{1*}, Hong-bo Shao^{2,3*}, Xin-tao Lei¹ and Yong Xiao¹

¹Hainan Key Laboratory of Tropical Oil Crops Biology, Coconut Research Institute, Chinese Academy of Tropical Agricultural Sciences, Wenchang 571339, Hainan, China.

²The CAS/Shandong Provincial Key Laboratory of Coastal Environmental Process, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences(CAS), Yantai 264003, Shandong, China.

³Institute for Life Sciences, Qingdao University of Science and Technology(QUST), Qingdao266042, Shandong, China.

Accepted 11 July, 2011

The research was conducted to detect changes in growth, physiology and nutrient concentration in response to two watering regimes (well-watered and water-stress conditions) and to two nutrient regimes (with or without fertilization) of oil palm. Under stress conditions, changes in plant growth, dry matter allocation, relative water content, leaf relative conductivity, leaf N, P and K concentration are usually observed. These characteristics and related parameters were determined and the experiment results are listed as follows: (1) fertilization promoted the growth of oil palm under well-watered conditions, while under water stress conditions its effects on growth was negative. The ratio of root/shoot was increased under water stress condition; (2) relative water content and chlorophyll a/b content were gradually decreased while leaf relative conductivity was increased quickly under water and nutrient stress conditions during the experiment. It is obvious that water stress had a greater influence than nutrient stress on these parameters; (3) water and nutrient stress decreased leaf nitrogen and phosphorus concentration but increased potassium concentration; the combination of water and nutrient stress made significant effects on nitrogen and phosphorus concentration, but no significant effects on potassium concentration. Moreover, deficiency of both water and nutrients in combination had the greatest impact on changes in these traits of oil palm.

Key words: Plant growth, physiology response, nutrient concentration, water stress, nutrient stress.

INTRODUCTION

Water and nutrient deficiency are major factors limiting the productivity and geographical distribution of many species, including important agricultural crops (Conner et al., 1998; Zhang et al., 2007; Tseira and Irit, 2009; Andrews et al., 2010). Fertilization is most effective when trees are not water-stressed, and irrigation is most effective when nutrients are not scarce (Sands and Mulligan, 1990). Therefore, understanding the mechanisms of plant tolerance to water and nutrient stress is a crucial environmental research topic (Wang et al., 2009a). Generally, exposure to water or nutrient stresses triggers many common reactions in plants that lead to a decrease in the

growth rate and relative water content; change of the biomass partition and nutrient distribution. Another consequence of exposure to these stresses is the increase in root/shoot ratios and leaf relative conductivity. Numerous studies have shown that plants will respond to a large set of parallel changes in growth, and in morphological and physiological responses when the plants are exposed to water or nutrient stress environment (Chapin, 1991).

Until now, many researchers clarified numerous effects on plant growth (Ismail et al., 1994; Wang et al., 2009b), dry matter allocation (Yin et al., 2005; Mandal et al., 2006; Wu et al., 2008; Li et al., 2009; Sun et al., 2009; Gerardeaux et al., 2010), photosynthesis (Syros et al., 2004), nutrient concentration and status (Wang et al., 1998; Jose et al., 2003; Slamaa et al., 2008; Keutgen and Pawelzik, 2009). The oil palm (*Elaeis guineensis* Jacq.) is a perennial monocotyledonous plant which belongs to the

*Corresponding authors. E-mail: hongxing1976@163.com, shaohongbuchu@126.com Tel: +86-532-84023984.

family Arecaceae originating from West Africa. The fruit pulp and nut that provide palm and kernel oil, respectively, made oil palm a high yielding oil-producing crop (Henderson and Osborne, 2000; Corley and Tinker, 2003). At present, palm oil production is second only to that of soybean oil in terms of world vegetable oil production and the demand for palm oil is expected to increase in future (Ho et al., 2007; Yusof, 2007; Corley, 2009; Bateman et al., 2010). In order to meet the increasing demand for palm oil, an improvement in yield is required despite the large body of literature on water and nutrient stress, as the important stress of oil palm in most tropical or subtropical area.

Although some reports showed that the responses of oil palm are related to water stress (Smith, 1988; Caliman, 1992; Gawankar et al., 2003; Kallarackal et al., 2004; Henson et al., 2005; Mohd Roslan Noor, 2006; Legros et al., 2009; Omorefe and Bonaventure, 2010; Cha-um et al., 2010), or the effects of nutrient on plant growth, physiological variance and yield of oil palm (Ng, 1979; Van Kraalingena et al., 1989; Remison et al., 2000; Bah and Rahman, 2004; Taryo-Adiwigandaa et al., 2006; Anuar et al., 2008; Tan et al., 2010), to our knowledge little is known about the effects of water and nutrient stress on this plant. Therefore, there is a pressing need to know in more detail how oil palm responds and adapts to such conditions. In this work, we examined the variance of plant growth, dry matter allocation, relative water content, leaf relative conductivity, chlorophyll a/b, changes of leaf total N and P concentration were also determined.

MATERIALS AND METHODS

Plant material and experimental design

The experiment design combined two water levels [90 and 50% field water capacity (FC)] and two nutrient levels (no fertilizer supplied to the plants or irrigated with Hoagland solution to each plant every 15 days). In the well-watered treatment, the pots were rewatered to FC by replacing the amount of transpired water every second day. In the water-stressed treatments, the pots were watered to 50% of FC every second day to maintain the drought level in the soil. Therefore, four experimental treatments were conducted: +W+F and -W+F, were control treatment (90% field water capacity with applied fertilizer) and water stress treatment (50% field water capacity with applied fertilizer) respectively; while +W- F and -W- F, were nutrient stress treatment (90% field water capacity without applied fertilizer), and water and nutrient stress treatment (50% field water capacity without applied fertilizer), respectively. Water leakage from bottom of the pot and evaporation from the soil surface was prevented by enclosing all soil in plastic bags. Thereafter, pots were divided into four groups; three replicates each. The pots were randomly switched places often, so as to decrease the differences in microclimates. The seedlings were randomly sampled at 0, 30, 60, and 90 days for measuring the investigated traits.

Growth measurements

Growing plant height was measured as the distance from soil surface to upper end of the longest leaf. The total leaf area was

determined by a portable laser area meter (CI-203, CID Inc., USA). Whole plants from all treatments were harvested at 0, 30, 60, and 90 days of the experiment, and divided into leaves, stems and roots. Plant samples were first dried at 120°C for 2 h, and then oven-dried at 80°C until constant weight and weighted with analytical balance. Root/shoot ratio (Rs) were then calculated.

Measurements of relative water content (RWC), leaf relative conductivity, and chlorophyll content

RWC of leaves

RWC was estimated by recording the turgid weight of fresh leaf samples by keeping them in water for 24 h, followed by drying in a hot air oven until constant weight was achieved. $RWC = (FW - DW) \times 100 / (TW - DW)$, where FW is the fresh weight, TW is the turgid weight of dehydrating samples which were enclosed in black envelope at about 25°C for 24 h and DW is the dry weight of leaves after oven-drying samples at 85°C for 24 h (Whetherley, 1950).

Leaf relative conductivity

After treatment, the fully expanded leaves were randomly selected from three seedlings per replication, and a DDS-11A Conductivity Meter was used to measure the conductivity of the solutions. The percentage of membrane damage was calculated as a relative conductance using the formula: $(C_1 - C_w) / (C_2 - C_w) \times 100$, where C_1 is the electrical conductivity value of samples at the first measurement, C_2 is the conductivity after boiling and C_w is the conductivity of deionized water. The relative conductivity was carried out according to the method described by Premachandra et al. (1991) described with little modification.

Chlorophyll content

The fresh leaves were ground in 80% acetone as quickly as possible at room temperature, and the chlorophyll (Chl) contents (Chla, Chlb and Chla/b) were determined using a spectrophotometer (SCHOTT UVLINER 9400, German) at 470, 646 and 663 nm and by the calculations described by Wellburn (1994).

Determination of nitrogen, phosphorus, and potassium content in oil palm leaves

At the end of the experiment, the leaves for the nitrogen, phosphorus, and potassium content analysis were dried at 80°C until constant weight and then ground. The dried samples were milled and subsequently digested with concentrated H_2SO_4 and H_2O_2 . The total N and its proportional concentration (N%) were analyzed with the micro-Kjeldahl method (Liang and MacKenzie, 1994). The total P was determined by using molybdate-blue colorimetric method. The calculation of phosphorus content was as described by Kitson and Melon (1944) and Hocking and Meyer (1991). Potassium from samples was determined by analyzing with a flame photometer. Potassium content was calculated as mentioned by Siddiqi and Glass (1981) and Yang and Zhou (2007).

Statistical analyses

The results presented are the mean values \pm standard errors obtained from at least three replicates. Significant differences between the treated and control plants were determined using ANOVA test ($P < 0.05$ and $P < 0.01$). Statistical analyses were conducted using the statistical soft ware package SPSS11.5 for Windows.

Table 1. The effect of watering and fertilization on plant growth and biomass partitioning.

Variable	Day	Treatment			
		CK (+W+F)	-W+F	+W-F	-W-F
Plant height	0	68.94±1.22 ^a	69.07±1.30 ^a	68.81±1.21 ^a	69.09±1.21 ^a
	30	73.50±1.34 ^a	72.40±1.33 ^a	70.00±1.25 ^{ab}	69.92±1.22 ^b
	60	77.00±2.00 ^a	72.67±1.60 ^a	72.08±1.31 ^a	71.25±1.37 ^a
	90	79.00±2.10 ^a	73.60±1.71 ^{ab}	73.13±1.35 ^{ab}	72.50±1.31 ^b
Leaf area	0	260.08±1.53 ^a	259.43±0.78 ^a	258.93±2.01 ^a	259.29±1.90 ^a
	30	303.57±2.15 ^a	269.49±1.90 ^b	295.35±0.47 ^a	265.69±1.42 ^b
	60	316.19±1.32 ^a	272.98±2.67 ^b	308.48±2.01 ^a	267.95±1.09 ^b
	90	331.18±1.91 ^a	279.53±0.69 ^b	321.12±2.36 ^a	272.05±1.37 ^c
Root/shoot ratios	0	0.19±0.41 ^a	0.19±0.47 ^a	0.19±0.43 ^a	0.19±0.56 ^a
	30	0.23±1.07 ^b	0.26±0.54 ^a	0.25±1.11 ^b	0.26±1.23 ^a
	60	0.26±1.81 ^b	0.28±3.10 ^a	0.25±1.54 ^c	0.27±1.13 ^a
	90	0.27±0.57 ^{ab}	0.34±1.43 ^a	0.26±2.21 ^b	0.29±1.03 ^a

Different letters in the columns indicate significant differences between the watering and fertilization treatments at $P < 0.05$ level.

RESULTS

Plant growth and biomass partitioning

The plant growth that was sustainably increased during the experimental period was observed. After about 3 months of continuous growth, the dry matter accumulation and allocation were significantly affected by water and nutrient stress (Table 1). With the duration of stress time, the growth rate was slower than the initial stage. At the end of the experiment, water and nutrient stress decreased both plant height and leaf area, resulting in a significant water \times nutrient interaction. Seedling grew taller in the control treatment compared to other treatments. Similarly, a significant increase in leaf area was observed for seedlings in the control treatment over the water and nutrient stress treatment.

The root : shoot ratio was mainly influenced by water treatments. More biomass was partitioned to above-ground tissue in both nutrient stress treatment and control treatment than water stress treatment, respectively. Conversely, below ground allocation was greater in the water treatment than other treatments. These distribution patterns resulted in increased root : shoot ratio for seedling grown under water stress. It is obvious that water stress had a far greater influence on root : shoot ratio than nutrient stress.

RWC, leaf relative conductivity, chlorophyll a/b

RWC, leaf relative conductivity, and chlorophyll a/b were affected by the experimental factors (Table 2). RWC, and

chlorophylla/b gradually decreased and leaf relative conductivity greatly increased under water and nutrient stress during the experimental period. The RWC was highest in the control (+W+F) treatment, but lowest in the -W-F treatments. Conversely, leaf relative conductivity was significantly higher in the water stress treatments than in the well-watered treatments. At the end of the experiment, Chlorophyll a/b was significantly influenced by water and nutrient stress. However, Chlorophyll a/b remained stable under control treatment. These results also showed that water stress had a greater influence on these traits than nutrient stress.

Nitrogen, phosphorus and potassium concentration

As shown in Table 3, at the end of the experiment, the main effects, the interaction between water and nutrient stress, was significant for leaf nitrogen and phosphorus concentration ($P < 0.05$), but had no significant effect on potassium concentration. Nitrogen, and phosphorus concentration was lower in the nutrient and nutrient-water stressed plants compared to the control plant. The highest decrease in nitrogen, phosphorus, potassium concentrations were 11.4 and 27.4%, respectively. But potassium concentration was increased under stress condition.

Relationship between physiological and biochemical parameters of oil palm seedlings

It can be seen from Table 4, that under +W+P (control)

Table 2. Effect of watering and fertilization on relative water contents, leaf relative conductivity, Chlorophyll a/b.

Variable	Day	Treatment			
		CK (+W+F)	-W+F	+W-F	-W-F
Relative water contents	0	97.74±1.33 ^a	93.72±1.30 ^a	95.70±1.31 ^a	96.84±1.34 ^a
	30	96.4±1.25 ^a	92.28±0.89 ^a	93.85±0.84 ^a	90.04±0.68 ^b
	60	96.37±1.22 ^a	84.94±0.47 ^b	85.24±0.87 ^b	75.6±0.31 ^c
	90	96.34±1.31 ^a	70.43±0.21 ^{bc}	78.78±0.46 ^b	56.55±0.09 ^c
Leaf relative conductivity	0	104.22±1.88 ^a	102.21±1.78 ^a	103.22±1.68 ^a	101.12±1.58 ^a
	30	149.87±1.88 ^c	173.2±1.91 ^a	169.29±1.85 ^{bc}	170.83±2.01 ^a
	60	151.96±1.92 ^c	593.92±2.048 ^{ab}	209.03±2.18 ^{bc}	870.62±2.58 ^a
	90	162.99±1.98 ^c	1053.25±3.68 ^a	368.25±2.44 ^{bc}	1153.99±3.88 ^a
Chlorophyll a/b	0	2.01±1.08 ^a	2.11±1.31 ^a	2.13±1.64 ^a	2.14±1.41 ^a
	30	2.00±2.12 ^a	1.81±1.41 ^a	1.83±1.54 ^a	1.78±1.03 ^b
	60	1.99±1.88 ^a	1.63±1.12 ^b	1.79±1.24 ^{ab}	1.48±0.87 ^c
	90	1.93±1.67 ^a	1.42±0.78 ^b	1.62±1.02 ^{ab}	1.21±0.47 ^c

Different letters in the columns indicate significant differences between the watering and fertilization treatments at $P < 0.05$ level.

Table 3. Effect of watering and fertilization on leaf nitrogen, phosphorus, and potassium concentration at final harvest.

Variable	Treatment			
	CK (+W+F)	-W+F	+W-F	-W-F
Nitrogen concentration (%)	1.15±0.12 ^a	1.14±0.14 ^a	0.97±0.05 ^b	0.93±0.05 ^b
Phosphorus concentration (%)	0.12±0.01 ^a	0.11±0.01 ^a	0.10±0.00 ^{ab}	0.09±0.00 ^b
Potassium concentration (%)	0.97±0.09 ^b	1.26±0.12 ^a	1.02±0.11 ^{ab}	0.99±0.11 ^b

Different letters in the columns indicate significant differences between the watering and fertilization treatments at $P < 0.05$ level.

treatment, leaf area showed a significant positive and negative correlation with relative water content and potassium concentration, respectively ($P < 0.05$). Under -W+F treatment, Chlorophyll a/b showed a positive correlation with plant height ($P < 0.05$) and leaf relative conductivity ($P < 0.01$). In addition, the correlation between plant height and leaf relative conductivity, phosphorus concentration and root/shoot ratios, relative water content and nitrogen concentration also showed significant positive correlation ($P < 0.05$). Under +W-F treatment, leaf area and phosphorus concentration, nitrogen and potassium concentrations were significantly positive and negatively correlated at the significance level, ($P < 0.01$) respectively. The positive correlation between relative water content and root/shoot ratios was also highly significant ($P < 0.05$). Under -W-F treatment, leaf area and Chlorophyll a/b, root/shoot ratios and potassium concentration were significantly positive and negatively correlated at the significance level, ($P < 0.01$) respectively, and phosphorus concentration showed a significant positive correlation with leaf area and

Chlorophyll a/b ($P < 0.05$).

DISCUSSION

Under environmental stresses, plants would change their biomass allocation between organs, and the growth rate would gradually become slow with the duration of stress time. At the end of experiment, the effects of the interaction between water and fertilization on plant height, leaf area, and dry mass accumulation (including total biomass, root, stem and leaf mass) were significant. Water and nutrient stresses suppressed growth rate, as also previously observed (Hsiao, 1973; Munns, 2002; Karageorgou et al., 2002; Stefens et al., 2005; Song et al., 2010; Weih, 2010). Judging by the variance of plant growth in Table 1, there was no significant difference between water and nutrient stress, when applied separately on plant growth. Some reports argue that water is more limiting than nutrient because the increase in growth between well-watered and water stressed with

Table 4. Relationship between parameters, plant height (pH), leaf area (LAR), R/S (root/shoot ratios), RWC (relative water content), RC (leaf relative conductivity), Chl a/b (Chlorophyll a/b), N con (nitrogen concentration), phosphorus concentration (P con), potassium concentration (K con) of oil palm seedlings grown under water and nutrient stress at final harvest.

Treatment	Correlation Coefficient									
	Traits	PH	LAR	R/S	RWC	RC	Chl a/b	N con	P con	k con
+W+F	LAR									
	R/S									
	RWC		*							
	RC									
	Chl a/b									
	N con									
	P con									
	k con			-*						
-W+F	LAR									
	R/S									
	RWC									
	RC	*								
	Chl a/b	*					**			
	N con					*				
	P con				*					
	k con									
+W-F	LAR									
	R/S									
	RWC				*					
	RC									
	Chl a/b									
	N con									
	P con			**						
	k con								-**	
-W-F	LAR									
	R/S									
	RWC									
	RC									
	Chl a/b		**							
	N con									
	P con		*				*			
	k con				-**					

** and * indicate that positive correlation was significant at 0.01, and 0.05 probability levels, respectively; -** and -* indicate that negative correlation was significant at 0.01 and 0.05 probability levels, respectively.

fertilization (nutrients not limiting) was greater than between fertilized and non-fertilized in the well-watered treatment (Yin et al., 2009). But Stroup et al. (2003) have suggested that nutrient determines the plant growth even more than water availability does.

The nutrient availability has been found to alter allocation patterns: the values of the parameter root: shoot, decreased in the conditions with fertilization when compared to conditions without fertilization, despite the same watering regimes. These results indicate that supply fertilization treatments could reduce the growth of roots as well as promote the growth of shoots, especially the growth of leaves. Water or nutrient stress alone did not alter seedling root/shoot ratio, nutrient stress alone had less effect than water stress on this trait, but the com-

ination of water and nutrient stress significantly increased the root/shoot ratio. It was consistent with the report by Stark (1992) and Zainudin et al. (2003), which suggested that simultaneous water and nutrient had the greatest impact on root/shoot ratio. Plant modification of physiological traits such as RWC, leaf relative conductivity, and Chlorophyll a/b are the earlier response to stress environment than plant growth. As shown in Table 2, the results show that RWC and Chlorophyll a/b decreased with the water stress in both fertilizer supply and without fertilizer supply conditions, while the parameter, leaf relative conductivity decreased under the fertilizer supply only under well-watered conditions but greatly increased under water-stressed conditions. Similar results were also reported by previous studies (Legros et

al., 2009; Song et al., 2010; Sanchez et al., 1983; Correia et al., 1989; Marulanda et al., 2010; Li et al., 2010; Yousfi et al., 2010). When the treatment was observed up to 60 or 90 days, our result showed that RWC and Chlorophyll a/b was significantly lower and leaf relative conductivity was significantly higher in water-stressed treatment than in the well-watered treatment, but it increased with nutrient supply. This indicated that water stress applied separately had more effect than nutrient stress on these three traits.

It is shown in Table 3, that at the end of the experiment, water and nutrient stress decreased leaf nitrogen and phosphorus concentration but increased potassium concentration compared to the control treatment ($P < 0.05$); the combination of water and nutrient stress made significant effects on nitrogen and phosphorus concentrations, but there was no significant difference between the control and the potassium concentration. Many studies have been carried out to indicate that water deficit reduced N and P concentrations in leaves; N and P were affected more than K and plant nutrient content were lower under drought stress and soil nutrient shortage and overdose; their findings are in accordance with the present results (Eck and Musick, 1979; Shangguan et al., 2000; Bruyn et al., 2002). It can be seen from Table 4 that the more significant correlations between leaf area, root/shoot ratios, Chlorophyll a/b and phosphorus concentration with other index under different treatments than other investigated traits, may be good indicators for the physiological and growth characters, nutrient concentration responses to water and nutrient stress of oil palm seedlings.

In conclusion, the growth, biomass partitioning, nutrient concentration and the morphological and physiological properties of oil palm were affected by water and nutrient availability, but it appeared more affected by water stress. Fertilization under well-watered conditions alleviated the nutrient stress, while fertilization under water-stressed conditions aggravated the drought stress, by dissolving the fertilizer. The result also suggests that water \times nutrient interaction had the greatest influence on changes in these traits of oil palm.

ACKNOWLEDGEMENT

The work is jointly supported by the Central Public-interest Scientific Institution Basal Research Fund (ITBKF081), Feasibility Analysis of Development of Oil Palm and Tea-oil Tree Industry (2010-YBJ-03), One Hundred-Talent Plan of Chinese Academy of Sciences (CAS), the CAS/SAFEA International Partnership Program for Creative Research Teams, the National Natural Science Foundation of China(41171216), the Strategic Priority Research Program of the Chinese Academy of Sciences(CAS)(XDA01020304),Yantai Double-hundred Talent Plan(XY-003-02) and the Science and Technology Development Plan of Yantai City (2011016).

REFERENCES

- Andrews M, Hodge S, Raven JA (2010). Positive plant microbial interactions. *Ann. Appl. Biol.* 157: 317-320.
- Anuar AR, Goh KJ, Heoh TB, Ahmed OH (2008). Spatial-temporal yield trend of oil palm as influenced by nitrogen fertilizer management. *Am. J. Appl. Sci.* 5: 1376-1383.
- Bah AR, Rahman ZA (2004). Evaluating urea fertilizer formulations for oil palm seedlings using the ^{15}N isotope dilution technique. *J. Oil Palm Res.* 16: 72-77.
- Bateman IJ, Fisher B, Fitzherbert E, Glew D, Naidoo R (2010). Tigers, markets and palm oil: market potential for conservation. *Oryx* 44: 230-234.
- Bruyn LD, Scheirs J, Verhagen R (2002). Nutrient stress, host plant quality and herbivore performance of a leaf-mining fly on grass. *Plant Anim. Interactions* 130: 594-599.
- Caliman JP (1992). Oil palm and water deficit: production, adapted cropping techniques. *Oleagineux* 47: 205-216.
- Chapin FS (1991). Integrated responses of plants to stress. *BioScience* 41: 29-36.
- Cha-um S, Takabe T, Kirdmanee C (2010). Osmotic potential, photosynthetic abilities and growth characters of oil palm (*Elaeis guineensis* Jacq.) seedlings in responses to polyethylene glycol-induced water deficit. *Afr. J. Biotechnol.* 39: 6509-6516.
- Conner J K, Zangori LA (1998). Combined effects of water, nutrients and UV-B stress on female fitness in *Brassica* (Brassicaceae). *Am. J. Bot.* 85: 925 - 931.
- Corley RHV (2009). How much palm oil do we need ? *Environ. Sci. Polic.* 12: 134-139.
- Corley RHV, Tinker PB (2003). *The Oil Palm* 4th edition. UK: Blackwell pub.
- Correia MJ, Torres F, Pereira JS (1989): Water and nutrient supply regimes and the water relations of juvenile leaves of *Eucalyptus globulus*. *Tree Physiol.* 5: 459-471.
- Eck HV, Musick JT (1979). Plant water stress effects on irrigated grain sorghum. Effects on nutrients in plant tissues. *Crop Sci.* 19: 592-598.
- Gawankar MS, Devmore JP, Jamadagni BM, Sagvekar VV, Khan HH (2003). Effect of water stress on growth and yield of Tenera oil palm. *J. Appl. Hort.* 5: 39-40.
- Gerardeaux E, Jordan-Meille L, Constantin J, Pellerin S, Dingkuhn M, (2010). Changes in plant morphology and dry matter partitioning caused by potassium deficiency in *Gossypium hirsutum* (L.). *Environ. Exp. Bot.* 67: 451-459.
- Henderson J, Osborne DJ (2000). The oil palm in our lives: how this came about. *Endeavour* 24-63.
- Henson IE, Noor MR, Harun MH, Yahya Z, Mustakim SNA (2005). Stress development and its detection in young oil palms in north Kedah, Malaysia. *J. Oil. Palm Res.* 17: 11-26.
- Ho CL, Kwan YY, Choi MC, Tee SS, Ng WH, Lim KA, Lee YP, Ooi SE, Lee WW, Tee JM, Tan SH, Kulaveerasingam H, Alwee SSRS, Abdullah MO, (2007): Analysis and functional annotation of expressed sequence tags (ESTs) from multiple tissues of oil palm (*Elaeis guineensis* Jacq.). *BMC Genomics* 381: 1-12.
- Hocking PJ, Meyer CP (1991). Effects of CO₂ enrichment and nitrogen stress on growth, and partitioning of dry matter and nitrogen in wheat and maize. *Aust. J. Plant Physiol.* 18: 339-356.
- Hsiao TC (1973). Plant responses to water stress. *Ann rev. plant physiol.* 24: 519-570.
- Ismail MR, Burrage SW, Tarmizi H, Aziz MA, (1994). Growth, plant water relations, photosynthesis rate and accumulation of proline in young carambola plants in relation to water stress. *Sci. Hort.* 60: 101-114.
- Jose S, Merritt S, Ramsey CL (2003). Growth, nutrition, photosynthesis and transpiration responses of longleaf pine seedlings to light, water and nitrogen. *Forest Ecol Manag.* 180: 335-344.
- Kallarackal J, Suman P, George J (2004). Water use of irrigated oil palm at three different arid locations in Peninsular India. *J. Oil. Palm Res.* 16: 45-53.
- Karageorgou P, Levizou E, Manetas Y (2002). The influence of drought, shade and availability of mineral nutrients on exudate phenolics of *Dittrichia viscosa*. *Morphol., Distribution, Functional Ecol. Plants.* 197:

- 285-289.
- Keutgen AJ, Pawelzik E (2009). Impacts of NaCl stress on plant growth and mineral nutrient assimilation in two cultivars of strawberry. *Environ. Exp. Bot.* 65: 170-176.
- Kitson RE, Melon MG (1944) Colorimetric determination of phosphorus as molybdovanado phosphoric acid. *Ind.Eng.Chem.Res.* 16: 379-383.
- Legros S, Mialet-Serra I, Caliman J-P, Siregar FA, Clement-Vidal A, Fabre D, Dingkuhn M (2009). Phenology, growth and physiological adjustments of oil palm (*Elaeis guineensis* Jacq.) to sink limitation induced by fruit pruning. *Ann Bot.* 104: 1183-1194.
- Li F L, Bao WK, Wu N (2009). Effects of water stress on growth, dry matter allocation and water-use efficiency of a leguminous species, *Sophora davidii*. *Agroforestry System* 77:193-201
- Li XW, Shen XF, Li JM, Eneji AE, Li ZH, Tian XL, Duan LS (2010). Coronatine alleviates water deficiency stress on winter wheat seedlings. *J. Integrative Plant Biol.* 52: 616-625.
- Liang BC, MacKenzie AF (1994). Corn yield, nitrogen uptake and nitrogen use efficiency as influenced by nitrogen fertilization. *Can. J. Soil Sci.* 74: 235-240.
- Mandal KG, Hati KM, Misra AK, Bandyopadhyay KK (2006). Assessment of irrigation and nutrient effects on growth, yield and water use efficiency of Indian mustard (*Brassica juncea*) in central India. *Agr. Water Manage.* 85: 279-286.
- Marulanda A, Azcón R, Chaumont F, Ruiz-Lozano JM, Aroca R (2010). Regulation of plasma membrane aquaporins by inoculation with a *Bacillus megaterium* strain in maize (*Zea mays* L.) plants under unstressed and salt-stressed conditions. *Planta* 232: 533-543.
- Mohd Roslan Noor MD (2006). Effects of water stress on the physiological processes and water use efficiency in oil palm. Masters thesis, Univ. Putra Malaysia
- Munns R (2002) Comparative physiology of salt and water stress. *Plant, Cell and Environ.* 25: 239-250.
- Ng SK (1979). Greater productivity of the oil palm (*Elaeis guineensis* Jacq.) with efficient fertilizer practices. In: Proceedings 9th Congress Inter. Potash & Phosphate Institute. .
- Omorefe A, Bonaventure C (2010). Identification of moisture stress tolerant oil palm genotypes African. *J. Agric. Res.* 5: 3116-3121.
- Premachandra GS, Saneoka H, Kanaya M, Ogata S (1991). Cell membrane stability and leaf surface wax content as affected by increasing water deficits in maize. *J. Exp. Bot.* 42: 167-171.
- Remison SU, Onwubuya I, Mgbeze GC (2000). The effect of fertilizer application and tapping position on sap flow in the oil palm (*Elaeis guineensis* Jacq.) . *Nigerian J. Agric. Forestry* 1:39-48.
- Sanchez R A, Hall AJ, Trapani N, Cohen Hunau R (1983): Effects of water stress on the chlorophyll content, nitrogen level and photosynthesis of leaves of two maize genotypes. *Photosynth. Res.* 4: 35-47.
- Sands R, Mulligan DR (1990). Water and nutrient dynamics and tree growth. *For. Ecol. Manage.* 30: 91-111.
- Shangguan ZP, Shao MA, Dyckmans J (2000): Nitrogen nutrition and water stress effects on leaf photosynthetic gas exchange and water use efficiency in winter wheat. *Environ. Exp. Bot.* 44:141-149.
- Siddiqi MY, Glass ADM (1981). Utilization index: A modified approach to the estimation and comparison of nutrient utilization efficiency in plants. *J. Plant Nutr.* 4: 249-302.
- Slamaa I, Ghnaya T, Savouré A, Abdelly C (2008). Combined effects of long-term salinity and soil drying on growth, water relations, nutrient status and proline accumulation of *Sesuvium portulacastrum*. *C. R. Biol.* 331: 442-451.
- Smith BG (1988). The effects of soil water and atmospheric vapour pressure deficit on stomatal behaviour and photosynthesis in the oil palm. *J. Exp. Bot.* 40: 647-651.
- Song CJ, Ma KM, Qu LY, Liu Y, Xu XL, Fu BJ, Zhong JF (2010). Interactive effects of water, nitrogen and phosphorus on the growth, biomass partitioning and water-use efficiency of *Bauhinia faberi* seedlings. *J. Arid. Environ.* 74: 1003-1012.
- Stark N (1992). The effects of water and multi-nutrient stress on xylem sap chemistry, photosynthesis and transpiration of seedlings of two Eucalypts. *Trees.* 6, 7-12.
- Stefens D, Hütsch BW, Eschholz T, Losak T, Schubert S (2005). Water logging may inhibit plant growth primarily by nutrient deficiency rather than nutrient toxicity. *Plant Soil Environ.* 51: 545-552.
- Stroup JA, Sanderson MA, Muir JP, McFarland MJ, Reed RL (2003). Comparison of growth and performance in upland and low land switch grass types to water and nitrogen stress. *Bioresour Technol.* 86, 65-72.
- Sun ZX, Zheng JM, Sun WT (2009). Coupled effects of soil water and nutrients on growth and yields of maize plants in a semi-arid region. *Pedosphere.* 19: 673-680.
- Syros T, Yupsanis T, Omiroua M, Economou A (2004). Photosynthetic response and peroxidases in relation to water and nutrient deficiency in gerbera. *Environ. Exp. Bot.* 52: 23-31.
- Tan NP, Zaharah AR, Siti Nor Akma A, Jamaluddin N (2010). Evaluating the variability of Gafsa phosphate rock uptake by oil palm genotypes at nursery stage. *Pertanika J. Trop. Agric. Sci.* 33: 223-231.
- Taryo-Adiwigandaa Y, Balaman T, Berani P (2006). Effect of Bio fertilizer on mature oil palm in North Sumatra and Riau. *Indonesian J. Agric. Sci.* 7: 20-26.
- Tseira M, Irit AC (2009). The effectiveness of the protection of riparian landscapes in Israel. *Land Use Policy* 26: 911-918.
- Van Kraalingena DWG, Breureb CJ, Spitters CJT (1989). Simulation of oil palm growth and yield . *Agric. For. Meteorol.* 46: 227-244.
- Wang JR, Hawkins CDB, Letchford T (1998): Photosynthesis, water and nitrogen use efficiencies of four paper birch (*Betula papyrifera*) populations grown under different soil moisture and nutrient regimes. *Forest Ecol. Manage.* 112: 233-244.
- Wang WB, Kim YH, Lee HS, Kim KY, Deng XP, Kwak SS (2009b). Analysis of antioxidant enzyme activity during germination of alfalfa under salt and drought stresses. *Plant Physiol. Biochem.* 47: 570-577.
- Wang ZQ, Wu LH, Animesha S (2009a). Nutrient shift in *Parthenocissus tricuspidata* seedling in response to soil water stress. *Scientia Horticulturæ* 123, 253-257.
- Weih M (2010). Evidence for increased sensitivity to nutrient and water stress in a fast-growing hybrid willow compared with a natural willow clone .*Tree Physiol.* 21:1141-1148.
- Wellburn AR (1994). The spectra determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *J. Plant Physiol.* 144: 307-313.
- Whetherley PE (1950). Studies in the water relations of cotton plants .I.The field measurement of water deficit in leaves. *New Phytol.* 49, 81-87.
- Wu FZ, Bao WK, Li FL, Wu N (2008). Effects of drought stress and N supply on the growth, biomass partitioning and water-use efficiency of *Sophora davidii* seedlings. *Environ. Exp. Bot.* 63: 248-255.
- Yang QQ, Zhou CH (2007). The test of potassium oxide on mixed and compound fertilizer by flame atomicabsorption spectrometry. *Contemporary Chem. Industry* 36: 209-211.
- Yin CY, Wang X, Duan BL, Luo JX, Li CY (2005). Early growth, dry matter allocation and water use efficiency of two sympatric *Populus* species as affected by water stress. *Environ. Exp. Bot.* 53: 315-322.
- Yin CY, Pang XP, Chen K (2009). The effects of water, nutrient availability and their interaction on the growth, morphology and physiology of two poplar species. *Environ. Exp. Bot.* 67: 196-203.
- Yousfi N, Slama I, Ghnaya T, Savouré A, Abdelly C (2010). Effects of water deficit stress on growth, water relations and osmolyte accumulation in *Medicago truncatula* and *M. laciniata* populations. *C. R. Biol.* 333: 205-213.
- Yusof B (2007). Palm oil production through sustainable plantations. *Eur.J. Lipid Sci.Technol.* 109: 289-295.
- Zainudin SR, Awang K, Hanif AHM (2003). Effects of combined nutrient and water stress on the growth of *Hopea Odorata* Roxb and *Mimusops Elengi* Linn. Seedlings. *J. Arboricul.* 29: 79-83.
- Zhang ZB, Shao HB, Xu P, Chu LY, Lu ZH, Tian JY (2007). On evolution and perspectives of bio-watersaving. *Biointer.*, 55: 1-9.