

Full Length Research Paper

Biological activity of sugarcane pyroligneous acid against *Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera: Noctuidae) larvae

Douglas Antunes Freitas Ferreira¹, Marcos Barbosa Ferreira², Silvio Favero² and Carlos Alexandre Carollo^{1*}

¹University of Mato Grosso do Sul (UFMS), Laboratory of Pharmacognosy, Av. Costa e Silva s/nº., 79070-900, Campo Grande-MS, Brazil;

²Universidade Anhanguera-Uniderp, Av. Alexandre Herculano, 1.400, 79037-280 - Campo Grande

Accepted 20 September, 2013

***Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera: Noctuidae) annually cause enormous loss to the producers and their combat has become a worldwide challenge mainly due to several reports of pesticides resistance. Today, one of the best alternatives used in this combat is the application of natural insecticides such as neem oil and pyroligneous acid. This study demonstrates a method to obtain a hexane fraction from sugarcane pyroligneous acid, which can be easily applied, as well as its effectiveness against *S. frugiperda*. The hexane fraction exhibited LC₅₀ of 2206,41 ppm after 24 h of exposure with a linear dose-response, indicating that the fraction can be used as a bio-insecticide against *S. frugiperda*.**

Key words: Wood vinegar, insecticide, *Saccharum officinarum*, fall armyworm, pyroligneous extract.

INTRODUCTION

The fall armyworm, *Spodoptera frugiperda* (Smith, 1797) (Lepidoptera: Noctuidae), is a polyphagous species of tropical and semitropical zones (Sparks, 1979) that causes severe damage to crops (Kamaraj et al., 2008). The control of this pest is usually made with the use of synthetic insecticides (Tavares et al., 2010a; Wititsiri, 2011) which, although effective, can cause various problems, such as the presence of residues in food, poisoning of applicators, and outbreaks of secondary pests (Roel et al., 2000; Berlitz et al., 2012). Furthermore, the continued use of these pesticides may aid in development of a resistant pest population (Munoz et al., 2013).

An alternative for combating crop pests is the use of natural products such as neem (Correia et al., 2009;

Tavares et al., 2010b), plant extracts (Roel et al., 2000; Tavares et al., 2009, 2011, 2013), and pyroligneous acid (Azevedo et al., 2007; Tavares et al., 2010a). The latter one is produced from the condensation of smoke from wood carbonization (Mendonça et al., 2006). This product is mainly characterized by low environmental impact and is a good alternative in organic agriculture. The pyroligneous acid, also known as wood vinegar, has been demonstrated to have low mammalian toxicity, lack neurotoxicity, have low persistence in the environment, and have high biodegradability (Céspedes et al., 2000).

The pyroligneous acid is normally composed of sugars, carboxylic acids, phenols (Fengel and Wegener, 1984; Kim et al., 2008), aldehydes, ketones, esters, furans and pyrans derivatives, nitrogen compounds, and other com-

*Corresponding author Email: carlos.carollo@ufms.br. Tel: +55 67 33457366.

pounds (Ninomiya et al., 2004), and the major part is constituted by water (85%) and acetic acid (5.1%) (Zanetti et al., 2003). Studies on the pyroligneous acid have reported bactericidal, fungicidal, and termiticidal activity (Mu et al., 2003).

The extract is usually obtained from species used to make charcoal (Cai et al., 2012). In Brazil, the species most used for charcoal production is the *Pinus elliottii* var. *elliottii* Engelm. (Pinaceae) (Porto et al., 2007). One of the viable alternatives for obtaining this extract is from sugarcane bagasse (*Saccharum officinarum* L., Poaceae); this residue has been routinely burned as an energy source (Zandersons et al., 1999), and hence, by a simple condensation of the smoke released, the extract can be obtained, which can be processed and used as a bio-insecticide. This process could avoid the annual emission of millions of tons of gases into the atmosphere (Alves et al., 2010), and produce a product with low environmental impact. The aim of this study was to obtain and process sugarcane pyroligneous acid and evaluate it against *S. frugiperda* larvae.

MATERIALS AND METHODS

Plant material and extract fractionation

The bagasse of *S. officinarum* was collected from Santa Olinda S/A, located in the municipal of Sidrolândia - Mato Grosso do Sul State, Brazil. The pyroligneous acid was obtained from the Laboratory of Pharmacognosy - (Federal University of Mato Grosso do Sul - UFMS) in Campo Grande, Mato Grosso do Sul State, Brazil. For that purpose, the bagasse was placed in an Erlenmeyer flask and completely burned in a muffle at 300°C, and then the smoke was condensed and the extract was collected. The crude extract had its pH neutralized and then was partitioned with hexane; this hexane fraction of the pyroligneous acid (HF-PA) was used in the biological assay.

Bioassay procedures

The eggs of *S. frugiperda* were provided by the Laboratory of Entomology of the Anhanguera University (Uniderp) in Campo Grande, Mato Grosso do Sul State, Brazil. After the eggs hatched, the larvae were placed on artificial diet (Greene et al., 1976) with 12-12-h photoperiod, at 27 ± 3°C, and under 70 ± 5% of relative humidity. The tests were divided into three groups of 10 neonatal larvae for each concentration: Negative group, Artificial diet and acetone; positive group, artificial diet plus neem oil at concentration of 12000 ppm (Neem oil 95%, Serôdia®, Campo Grande, Mato Grosso do Sul State, Brazil) plus acetone; and test group, artificial diet plus the HF-PA (60, 120, 360, 600, 1200, 3600, 6000, 12000, and 36000 ppm) and acetone to dilute the extract.

Statistical analysis

The experiment was monitored at 24 h period, observing mortality in relation to concentrations of HF-PA, and the statistical program used was SPSS Statistics 20.0.0 to apply simple linear regression model ($P < 0.05$). To calculate the LC_{50} , Probit analysis from the logarithms of the concentrations was used.

RESULTS AND DISCUSSION

The yield of pyroligneous extract was approximately 40%. This extract was neutralized once acetic acid constituted approximately 5% of the extract (Zanetti et al., 2003) because this compound could interfere in the bioassay and degrade other compounds present. Another step was the fractionation of the extract; this step was aimed mainly to obtain an extract without water. The HF-PA showed a neutral pH and can be easily concentrated; these characteristics allowed easy development of new products and application of this extract.

The bioassay using HF-PA showed significant change, when compared with the negative control group. All the different concentrations of the extract caused significant mortality in 24 h, which was also observed in the positive control group (neem oil). The highest concentration evaluated (36000 ppm) resulted in 100% death, and the lowest concentration (60 ppm) caused only 10% of deaths (Figure 1). The LC_{50} of HF-PA was 2206,41 ppm. Our results suggested a linear dose-response (Figure 2), which was also confirmed by r^2 (0.83). Neonatal nymphs of *Bemisia tabaci* (Gennadius, 1889) (Hemiptera: Aleyrodidae) showed 66.59% mortality caused by neem oil (Natuneem®) and 67.45% of deaths caused by pyroligneous acid (Pironat®) (Azevedo et al., 2005). Neem oil (Natuneem®) and crude pyroligneous extract (Biopiról7M®) tested on eggs of *S. frugiperda* and *Diatraea saccharalis* (Fabricius, 1794) (Lepidoptera: Crambidae), confirmed mortality of eggs of different ages with different concentrations of the extract, and the extracts' satisfactory effect in controlling these pests (Tavares et al., 2010a, b). However, neem oil (Natuneem®) and commercial crude pyroligneous acid (Biopiról7M® and Pironat®), no significant mortality was observed when the extract was applied to adults of *Anastrepha fraterculus* (Wiedemann, 1830) (Diptera: Tephritidae) (Efrom et al., 2011). These contradictions among different studies are related to the stages of the insects used in the experiments. Another possibility is related to the tolerance of the insect to insecticides in the course of their development (Yu, 1983; Schmutterer, 1990).

One of the apparent causes of the activity against the larvae is the synergistic effect among the compounds (Richards et al., 2010); this synergism occurs between different classes of compounds or structurally similar compounds, and the effects may range from antifeeding effect to toxicity activity (Dyer et al., 2003). The wide range of compound prevents resistance development (Rice, 1993). These secondary metabolites have different sites of action and molecular targets. They interact with important metabolic and enzymatic inhibitors (Céspedes et al., 2000; Torres et al., 2003) against microbial pathogens and invertebrates (Wink and Schimmer, 1999). Thus, for example, some secondary metabolites such as the nitrogenous compounds may act by blocking




Figure 1. Distribution of percentages of mortality of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae caused by HF-PA.




Figure 2. Graph regression analysis ($P < 0.05$) presenting the log of the dose against the number of dead *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae during 24 h.

the nervous system activity (Kagabu, 2008), or by inhibiting synthesis of deoxyribonucleic acid (DNA), and modifying the permeability of the membrane by changing the composition with respect to structural proteins (Schmeller et al., 1997). Another mechanism of action of PA-HF could be related to the presence of phenolic compounds, one of the main groups being investigated for pest control (Henn, 1997).

The pyroligneous extract used was toxic to *S. frugiperda* larvae and displayed a linear dose-response with an LC_{50} of 2206,41 ppm. Furthermore, the possibility of preparation of a low-cost fraction of the sugarcane pyroligneous acid was also demonstrated. These results open new possibilities for the fraction to be tested against different pests of crops and agriculture, providing useful application for the smoke produced by the burning of

sugarcane bagasse.

REFERENCES

- Alves OJ, Zhuo C, Levendis YA, Tenorio JAS (2010). Analysis of light hydrocarbon gases in the pyrolysis process of sugarcane bagasse. Congr. Anu. ABM 65th, 766-773.
- Azevedo FR, Guimarães JA, Sobrinho RB, Lima MAA (2005). Eficiência de produtos naturais para o controle de *Bemisia tabaci* biótipo B (Hemiptera: Aleyrodidae) em meloeiro. Arq. Inst. Biol. 72:73-79.
- Azevedo FR, Leitão ACL, Lima MAA, Guimarães JA (2007). Eficiência de produtos naturais no controle de *Callosobruchus maculatus* (Fab.) em feijão caupi (*Vigna unguiculata* (L.) Walp) armazenado. Rev. Ciênc. Agron. 38:182-187.
- Berlitz DL, Azambuja AO, Sebben A, Oliveira JV, Fiuza LM (2012). Mortality of *Oryzophagus oryzae* (Costa Lima, 1936) (Coleoptera: Curculionidae) and *Spodoptera frugiperda* (JE Smith, 1797) (Lepidoptera: Noctuidae) larvae exposed to *Bacillus thuringiensis* and extracts of *Melia azedarach*. Braz. Arch. Biol. Technol. 55:725-731.
- Cai K, Jiang S, Ren C, He Y (2012). Significant damage-rescuing effects of wood vinegar extract in living *Caenorhabditis elegans* under oxidative stress. J. Agric. Food Chem. 92:29-36.
- Céspedes CL, Calderón JS, Lina L, Aranda E (2000). Growth inhibitory effects on fall armyworm *Spodoptera frugiperda* of some limonoids isolated from *Cedrela* spp. (Meliaceae). J. Agric. Food Chem. 48:1903-1908.
- Correia AA, Teixeira VW, Teixeira AAC, Oliveira JV, Torres JB (2009). Morfologia do canal alimentar de lagartas de *Spodoptera frugiperda* (J E Smith) (Lepidoptera: Noctuidae) alimentadas com folhas tratadas com nim. Neotrop. Entomol. 38:83-91.
- Dyer LA, Dodson CD, Stireman III, JO, Tobler, MA, Smilanich AM, Fincher RM, Letourneau DK (2003). Synergistic effects of three *Piper* amides on generalist and specialist herbivores. J. Chem. Ecol. 29:2499-2514.
- Efrom CFS, Redaelli LR, Meirelles RN, Ourique CB (2011). Laboratory evaluation of phytosanitary products used for control of the South American fruit fly, *Anastrepha fraterculus*, in organic farming. Crop Prot. 30:1162-1167.
- Fengel D, Wegener G (1984). Wood: Chemistry, ultrastructure, reactions. Walter de Gruyter, Berlin.
- Greene GL, Leppla NC, Dickerson WA (1976). Velvetbean caterpillar: a rearing procedure and artificial medium. J. Econ. Entomol. 69:487-488.
- Henn MW (1997). Adsorption von Tanninen aus Eichenblättern an Kohlen e hydrathaltiger Verbindungen. Mitt. Dtsch. Entomol. Ges. 11:495-499.
- Kagabu S (2008). Pharmacophore of neonicotinoid insecticides. J. Pestic. Sci. 33:9-13.
- Kamaraj C, Rahuman AA, Bagavan A (2008). Antifeedant and larvicidal effects of plant extracts against *Spodoptera litura* (F.), *Aedes aegypti* L. and *Culex quinquefasciatus* Say. Parasitol. Res. 103:325-331.
- Kim DH, Seo HE, Lee SC, Lee KY (2008). Effects of wood vinegar mixed with insecticides on the mortalities of *Nilaparvata lugens* and *Laodelphax striatellus* (Homoptera: Delphacidae). Anim. Cells Syst. 12:47-52.
- Mendonça JMA, Carvalho GA, Guimarães RJ, Reis PR, Rocha LCD (2006). Produtos naturais e sintéticos no controle de *Leucoptera coffeella* (Guérin-Mèneville & Perrottet, 1842) (Lepidoptera: Lyonetiidae) e seus efeitos sobre a predação por vespas. Ciênc. agrotec. 30:892-899.
- Mu J, Uehara T, Furuno T (2003). Effect of bamboo vinegar on regulation of germination and radicle growth of seed plants. J. Wood Sci. 49:262-270.
- Muñoz E, Lamilla C, Marin JC, Alarcon J, Cespedes CL (2013). Antifeedant, insect growth regulatory and insecticidal effects of *Calceolaria talcana* (Calceolariaceae) on *Drosophila melanogaster* and *Spodoptera frugiperda*. Ind. Crop Prod. 42:137-144.
- Ninomiya Y, Zhang L, Nagashima T, Koketsu J, Sato A (2004). Combustion and De-SO_x behavior of high-sulfur coals added with calcium acetate produced from biomass pyrolytic acid. Fuel 83: 2123-2131.
- Porto PR, Sakita AEN, Sakita MN (2007). Efeito da aplicação do extrato pirolenhoso na germinação e no desenvolvimento de mudas de *Pinus Elliottii* var. *elliottii*. Instituto Florestal de São Paulo 31:15-19.
- Rice M (1993). Built-in Resistance Prevention (BIRP): a Valuable Property of Azadirachtin. In: World Neem Conference, India, Bangalore.
- Richards LA, Dyer LA, Smilanich AM, Dodson CD (2010). Synergistic effects of amides from two *Piper* species on generalist and specialist herbivores. J. Chem. Ecol. 36:1105-1113.
- Roel AR, Vendramim JD, Toyoko SFR, Frighetto N (2000). Efeito do extrato acetato de etila de *Trichillia pallida* Swartz (Meliaceae) no desenvolvimento e sobrevivência da lagarta-do-cartucho. Bragantia 59:53-58.
- Schmeller T, Latz-Bruning B, Wink M (1997). Biochemical activities of agberberine, palmatine and sanguinarine mediating chemical defence against microorganisms and herbivores. Phytochemistry 44:257-266.
- Schmutterer H (1990). Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. Annu. Rev. Entomol. 35:271-297.
- Sparks AN (1979). A review of the biology of the fall armyworm. Fla. Entomol. 62:82-87.
- Tavares WD, Cruz I, Petacci F, de Assis SL, Freitas SD, Zanuncio JC, Serrão JE (2009). Potential use of Asteraceae extracts to control *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and selectivity to their parasitoids *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) and *Telenomus remus* (Hymenoptera: Scelionidae). Ind. Crop Prod. 30:384-388.
- Tavares WD, Freitas SD, Graziotti GH, Parente LML, Lião LM, Zanuncio JC (2013). Ar-turmerone from *Curcuma longa* (Zingiberaceae) rhizomes and effects on *Sitophilus zeamais* (Coleoptera: Curculionidae) and *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Ind. Crop Prod. 46:158-164.
- Tavares WS, Costa, MA, Cruz I, Silveira RD, Serrão JE, Zanuncio JC (2010a). Selective effects of natural and synthetic insecticides on mortality of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and its predator *Eriopsis connexa* (Coleoptera: Coccinellidae). J. Environ. Sci. Health B 45:557-561.
- Tavares WS, Cruz I, Fonseca FG, Gouveia NL, Serrão JE, Zanuncio JC (2010b). Deleterious activity of natural products on postures of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Diatraea saccharalis* (Lepidoptera: Pyralidae). Z. Naturforsch. C. 65:412-418.
- Tavares WS, Cruz I, Petacci F, Freitas SS, Serrão JE, Zanuncio JC (2011). Insecticide activity of piperine: toxicity to eggs of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Diatraea saccharalis* (Lepidoptera: Pyralidae) and phytotoxicity on several vegetables. J. Med. Plant Res. 5:5301-5306.
- Torres P, Avila JG, Romo De Vivar A, García AM, Marín JC, Aranda E, Céspedes CL (2003). Antioxidant and insect growth regulatory activities of stilbenes and extracts from *Yucca periculosa*. Phytochemistry 64:463-473.
- Wink M, Schimmer O (1999). Modes of action of defensive secondary metabolites. In: Wink, M. (Ed.), Functions of Plant Secondary Metabolites and Their Exploitation in Biotechnology. Annual Plant Reviews No. 3. Sheffield Academic Press, Sheffield.
- Wititsiri S (2011). Production of wood vinegars from coconut shells and additional materials for control of termite workers, *Odontotermes* sp. and striped mealy bugs, *Ferrisia virgata*. Songklanakarin J. Sci. Technol. 33:349-354.
- Yu SJ (1983). Age variation in insecticide susceptibility and detoxification capacity of fall armyworm (Lepidoptera: Noctuidae) larva. J. Econ. Entomol. 76:219-222.
- Zandersons J, Gravitis J, Kokorevics A, Zhurinsh A, Bikovens O, Tardenaka A, Spince B (1999). Studies of the Brazilian sugarcane bagasse carbonisation process and products properties. Biomass Bioenergy 17:209-219.
- Zanetti M, Cazetta JO, Júnior Dirceu M, Carvalho SA (2003). Uso de subprodutos de carvão vegetal na formação do porta-enxerto limoeiro 'Cravo' em ambiente protegido. Rev. Bras. Frutic. 25:508-512.