

Full Length Research Paper

Optimisation of phenolic compound biosynthesis in grape (Bogazkere Cv.) callus culture

Mehmet KARAASLAN^{1*}, Mustafa OZDEN², Hasan VARDİN¹ and Fatih Mehmet YILMAZ¹

¹Harran University, Faculty of Agriculture, Department of Food Engineering, Sanliurfa TR-63040 Turkey.

²Harran University, Faculty of Agriculture, Department of Horticulture, Sanliurfa TR-63040 Turkey.

Accepted 17 June, 2013

The objective of this study was to improve metabolite accumulation of grape cells by manipulating culture and environmental conditions. Response surface methodology (RSM) was used to design and conduct the study with the independent variables being: percent depletion of KH_2PO_4 or NH_4NO_3 and light intensity. Wounded grape leaves (Bogazkere cv.) were used for the initiation of callus formation. Grape calli were placed on the modified Murashige and Skoog (MS) media (from 0 to 75% KH_2PO_4 or NH_4NO_3 depleted) under various light regimes (from 30 to $120 \mu\text{mol m}^{-2} \text{s}^{-1}$), and incubated in a controlled environment for three weeks to stimulate secondary metabolite biosynthesis. The anthocyanin - total phenolic (TP) contents of the calli were determined using chemical analyses and the antioxidant capacity of the cultures were determined by using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay. Higher KH_2PO_4 depletion and $75 \mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity favored the biosynthesis of anthocyanins and the other phenolic compounds and resulted in elevated antioxidant capacity in the cultures. However, an increase in KH_2PO_4 or NH_4NO_3 depletion resulted in a significant decrease in cell weight. Similar to that, increase of light intensity was correlated with decrease of cell weight. Further elevation of light intensity and KH_2PO_4 - NH_4NO_3 depletion was fatal to grape cells and resulted in formation of non-growing watery-brownish cultures. Individual phenolic compounds found in the grape cell cultures were identified using the gas chromatography-flame ionization detector (GC-FID) system. Catechin and epicatechin were identified as the most abundant simple phenolic compounds present in the cultures.

Key words: Antioxidant capacity, callus, grape, phenolic compounds, response surface methodology.

INTRODUCTION

Phenolic compounds have been the subject of numerous scientific reports for their ability to scavenge intracellularly produced free radicals. Free radical activity causes uncontrolled cell death, DNA damage, tumor formation and cardiovascular disorders (Fresco et al., 2010; Loke et al., 2010; von Sonntag, 2006). Demand for natural products expanded following epidemiological studies documenting the health promoting properties of phenolic compounds for prevention of disorders related to

free radical accumulation in the organisms (Szajdek and Borowska, 2008). Solvent extracts of the plants, fruits, leaves, and seeds are being used to obtain natural food colorants and functional compounds (Wallace and Giusti, 2008) valuable for industrial production of food additives, dietary supplements, functional foods and phyto-tablets. However, economical considerations, geographical limitations, seasonal restrictions and environmental problems (harsh chemical extraction) pushed researchers to seek

*Corresponding author. E-mail: mk385@cornell.edu. Tel: 90 554 819 6160. Fax: 90 414 318 368.

Abbreviations: RSM, Response surface methodology; HPLC, high performance liquid chromatography; NAA, 1-naphthaleneacetic acid; BA, 6-benzylaminopurine; 2,4-D, 2, 4- dichlorophenoxy acetic acid; DPPH, 1-diphenyl-2-picrylhydrazyl; TP, total phenolic; GC, gas chromatography; MS, Murashige and Skoog; Chs, Chalcone synthase.

alternative renewable resources for biosynthesis of flavonoids, anthocyanins, and other phenolic compounds (Blando et al., 2004; Nicoletti et al., 2008). *In vitro* cell cultures may pose an alternative and valuable tool for the production of bioactive compounds. Plant calli and cell cultures exhibit many advantages over plant tissues in obtaining of aforementioned compounds such as ease of extraction, manipulation of direction of biosynthesis, and scalable production. Secondary metabolite production using callus cultures has been reported in *Prunus cerasus*, *Daucus carota*, *Cleoma rosea*, *Rosmarinus officinalis*, *Vaccinium macrocarpon* (Blando et al., 2004; De Bellis et al., 2005; Glagben et al., 1992; Simoes et al., 2009; Kuhlmann and Rohl 2006; Mathavi et al., 1995).

Grape is one of the most potent fruits suitable for initiation of cell culture system for the production of phenolics and it displays high antioxidant activity with a wide range of secondary metabolites. Bogazkere cv. is a predominant grape variety growing in Southeastern and Eastern regions of Turkey with stringent taste and high phenolic content which makes it a good candidate to initiate cell cultures for production of secondary metabolites. Manipulation of medium composition, light intensity, temperature, growth regulators, and presence - concentration of intermediaetic compounds may affect the productivity and viability of *in vitro* cultures (Georgiev et al., 2008). Optimization of the callus growth conditions is one of the prerequisites for the stimulation of secondary metabolite biosynthesis and development of protocols allowing establishment of viable cultures. Response surface methodology (RSM) is being used to design experiments to meet the best operation conditions to achieve acceptable product yields, improve efficiency - profitability, while consuming limited time and amount of energy. The aim of this investigation was to determine the best medium composition regarding KH_2PO_4 or NH_4NO_3 concentrations and environmental conditions by using RSM skills to induce secondary metabolite biosynthesis in grape callus. The anthocyanin accumulation, phenolic content and antioxidant activity were investigated in the grape callus grown in various media and under different light regimes. The medium composition and light intensity were modified to improve the biosynthesis of the secondary metabolites by using a three factorial experimental design.

MATERIALS AND METHODS

Materials

High performance liquid chromatography (HPLC)-grade ethanol and methanol, HCl, ammonium nitrate, potassium chloride, sodium chloride, sodium carbonate, sodium acetate, Folin-Ciocalteu reagent, were purchased from MERCK Chemical Company, Germany. Gamborg-B5 medium, 1-naphthaleneacetic acid (NAA), 6-benzylaminopurine (BA), 2,4-dichlorophenoxy acetic acid (2,4-D), reagent-grade gallic acid, were purchased from Sigma Chemical Co. 1,1-diphenyl-2-picrylhydrazyl (DPPH) was obtained from Fluka

(Chemie AG, Bucks).

Grape callus induction and secondary metabolite biosynthesis

The grapevine leaves of *Vitis vinifera* L. cv. Bogazkere used for callus formation were taken from long-term *in vitro* cultures established and maintained according to the method described by Ozden et al. (2009). Leaf discs from grapevine (*V. vinifera*) cv. Bogazkere were excised and placed abaxial side up on solid Gamborg B-5 (Gamborg et al., 1968) medium with vitamins in petri dishes. Plant growth regulators, 0.2 mg/L BA and 0.02 mg/L 2, 4-D were added to the medium to reinforce callus formation (Keller et al., 2000). The pH value of the medium was adjusted to 5.8 prior to autoclaving (121°C , 1 kg cm^{-2} for 20 min). Calli were initiated from young leaf explants and incubated at $24 \pm 1^\circ\text{C}$ in darkness in controlled growth room. After four weeks, calli were separated from the explants and subcultured to fresh media for four weeks more for callus proliferation under 16 h photoperiod at $24 \pm 1^\circ\text{C}$. Afterwards, the calli were submerged and cultivated in Petri dishes with modified Murashige and Skoog (MS) media supplemented with 1 mg/L NAA and 0.1 mg/L BA for the induction of anthocyanin and the other phenolic compounds' biosynthesis. After three weeks of culture, calli were harvested, pooled, and used to obtain acidified methanol extracts of callus. Modifying the medium composition and light intensity on the biosynthesis of the secondary metabolites in grape callus was evaluated according to the response surface, 3-level factorial design (Table 1) with two variables; percent depletion (0 to 75%) of KH_2PO_4 and NH_4NO_3 in the MS and light intensity (30 to $120 \mu\text{mol m}^{-2} \text{ s}^{-1}$) based on the results obtained in the preliminary studies.

Extraction of phenolics from callus

For sample extraction, calli were homogenized in an ice cold blender and 10 g of the homogenate was macerated in 20 mL of methanol containing 20 μl conc. HCl and was left incubated in darkness overnight at 4°C . Then the extract was filtered over Whatman No. 1 paper under vacuum, and fresh acidified methanol was used to re-extract the residue until it became colorless. The alcohol in the extract was evaporated using rotary evaporator.

Total phenolic (TP) analysis

TP content of the callus were estimated according to the method of Slinkard and Singleton (1977) with some modifications. Samples of 0.03 mL were introduced into test tubes, 2.370 mL distilled water and 0.15 mL Folin-Ciocalteu's reagent was added and the tubes were vortexed vigorously. After 8 min, a 0.45 mL of saturated Na_2CO_3 was added to each tube, and then each mixture was vortexed again and allowed to stand for 30 min at room temperature. Absorbance was measured at 750 nm. Gallic acid was used to create a standard curve. The results were expressed as mg gallic acid equivalent.

Total anthocyanin analysis

Anthocyanin content of the callus was determined by the pH differential method described by Giusti and Wrolstad (1996) using a ultraviolet (UV)-vis spectrophotometer (Model U-2000, Hitachi). Absorbance of the samples in 0.025 M potassium chloride buffer (pH 1.0) and 0.4 M sodium acetate buffer (pH 4.5) were measured at 520 and 700 nm. Anthocyanin content was determined using the equation: $A = (A_{520} - A_{700})_{\text{pH } 1.0} - (A_{520} - A_{700})_{\text{pH } 4.5}$ with a molar extinction coefficient of 28,000. Results were expressed as mg of

Table 1. Experimental conditions of callus growth treatments for phenolic compounds.

Run Number	Independent variable				MS Media Component Categorical factor
	Depletion		Light Intensity		
	Coded level	Actual level	Coded level	Actual level	
1	-1	0.0	0	75	NH ₄ NO ₃
2	0	37.5	0	75	KH ₂ PO ₄
3	1	75.0	0	75	KH ₂ PO ₄
4	-1	0.0	1	120	NH ₄ NO ₃
5	0	37.5	0	75	NH ₄ NO ₃
6	0	37.5	0	75	KH ₂ PO ₄
7	0	37.5	0	75	NH ₄ NO ₃
8	-1	0.0	-1	30	NH ₄ NO ₃
9	0	37.5	1	120	NH ₄ NO ₃
10	0	37.5	0	75	KH ₂ PO ₄
11	-1	0.0	0	75	KH ₂ PO ₄
12	0	37.5	1	120	KH ₂ PO ₄
13	-1	0.0	1	120	KH ₂ PO ₄
14	1	75.0	1	120	KH ₂ PO ₄
15	0	37.5	0	75	NH ₄ NO ₃
16	0	37.5	0	75	NH ₄ NO ₃
17	1	75.0	1	120	NH ₄ NO ₃
18	1	75.0	0	75	NH ₄ NO ₃
19	-1	0.0	-1	30	KH ₂ PO ₄
20	0	37.5	-1	30	NH ₄ NO ₃
21	1	75.0	-1	30	NH ₄ NO ₃
22	1	75.0	-1	30	KH ₂ PO ₄
23	0	37.5	0	75	KH ₂ PO ₄
24	0	37.5	0	75	KH ₂ PO ₄
25	0	37.5	-1	30	KH ₂ PO ₄
26	0	37.5	0	75	NH ₄ NO ₃

malvidin-3-O-glucoside equivalent in per kg fresh fruit (Wrolstad, 1976).

Antioxidant capacity determination

Free radical scavenging activity (antioxidant capacity) of the samples were determined by the DPPH assay with minor modifications as described by Blois (1958). A 0.1 mM solution of DPPH in methanol was prepared. A 0.1 mL of various concentrations of the samples diluted in methanol was added to 2.9 mL of DPPH solution. The decrease in absorbance at 517 nm was measured after 30 min of incubation at room temperature. Radical scavenging capacity of each sample was expressed as percent DPPH radical scavenging effect using the following equation.

$$\text{DPPH scavenging effect (\%)} = [(A_c - A_s) / A_c \times 100]$$

Where, A_c is the absorbance of the control in ethanol, and A_s is the absorbance of the sample. Assays were carried out in triplicates.

Gas chromatography (GC) analysis

The phenolic compounds found in grape calli were identified and

quantified using a Thermo Quest, CE Instrument trace GC equipped with a flame ionization detector (GC-FID). Phenolic compounds were separated by using a ZB-5 (Zebron) capillary GC column (30 m x 0.25 mm x 0.25 μ m film thickness). The method described by Soleas et al. (1997) was essentially used for solid phase extraction and derivatization. One μ l of solution injected to GC for analysis. Similar GC temperature program, carrier gas flow rate, and split ratio were used as mentioned in an earlier study (Soleas, Diamandis, Karumanchiri and Goldberg, 1997). Authentic standards were used to identify phenolic compounds according to their retention times. Quantification was accomplished by using fisetin (10mg/L) as internal standard. The standards used in this study were kindly provided either by Dr. Geza Hrazdina (Cornell University) or obtained from the Sigma company.

Statistical analysis

The response surface, 3-level factorial design was used to determine the influence of modifying the medium composition and light intensity on the biosynthesis of the secondary metabolites in grape callus. The responses were analyzed using the software trial version Design-Expert[®] V.7.1.6, StatEase, Inc., (MN, USA). The effects of the independent variables on the properties of the callus

Table 2. Experimental responses from the callus extracts treatments.

Run	Response			
	Anth.	TP	AC	CW
1	34.5±0.2	1690±12	35±1	6.2±0.2
2	49.4±0.3	3045±24	66±2	1.5±0.1
3	68.9±0.2	8021±72	91±3	1.6±0.1
4	31.5±0.2	1455±20	31±2	6.4±0.2
5	40.1±0.3	2618±26	54±1	4.0±0.2
6	49.1±0.2	3020±20	66±2	1.4±0.1
7	40.5±0.2	2590±29	55±3	4.0±0.2
8	24.8±0.1	1315±13	31±2	6.4±0.3
9	39.7±0.2	1920±14	49±1	4.0±0.1
10	49.6±0.2	3082±23	66±1	1.5±0.1
11	34.5±0.1	1690±15	34±2	6.2±0.2
12	43.5±0.2	2151±21	52±1	1.9±0.1
13	31.5±0.1	1455±11	31±1	6.4±0.3
14	54.6±0.2	7650±31	83±3	1.8±0.1
15	40.6±0.1	2580±32	55±2	4.0±0.2
16	40.8±0.1	2628±19	55±2	4.0±0.2
17	43.2±0.1	2650±32	70±1	4.1±0.1
18	51.0±0.1	3610±23	73±2	4.1±0.1
19	24.8±0.1	1315±14	31±2	6.5±0.2
20	36.5±0.1	1725±16	48±1	4.4±0.1
21	41.4±0.1	2511±23	68±2	4.2±0.1
22	50.7±0.2	6410±42	75±2	4.1±0.2
23	49.6±0.3	3051±26	64±2	1.4±0.0
24	49.7±0.3	3057±32	65±3	1.4±0.0
25	41.1±0.1	2015±27	51±2	4.3±0.2
26	39.4±0.1	2604±22	55±2	4.0±0.1

Anth, anthocyanin content of callus (mg/kg); **TP**, (mg/kg) total phenolics of callus; **AC**, (% Inhibition), antioxidant capacity of callus extracts; **CW**, (g) cell weight of callus.

were assessed with a design consisted of 26 treatments. A quadratic model was fitted to the experimental data to predict the dependent variable Y (percent depletion of chemical (0-75 %) and light intensity (30-120)):

$$Y = a + b x_1 + c x_2 + d x_1 x_2 + e x_1^2 + f x_2^2$$

Where *a*, *b*, *c*, *d*, *e*, *f*, denote the regression coefficients. The three level factorial design model was validated statistically by analysis of variance (ANOVA), calculation and evaluation of the multiple-correlation coefficients, and estimation of the lack of fit.

RESULTS AND DISCUSSION

Secondary metabolite producing grape callus cell lines were placed on MS media containing different concentrations of phosphate and nitrate ions (KH₂PO₄ or NH₄NO₃ concentration was decreased from 0% to 75%), and exposed to various light intensities (30 to 120 μmol m⁻² s⁻¹). Grape calli were harvested after three weeks of incubation in a tissue culture room and examined for their anthocyanin - total phenolic content, and antioxidant capacity by using chemical experiments. The experimental responses of the light and medium treatments on

grape calli are shown in Table 2, while the regression coefficients for each dependent variable (response) as a function of *x*₁ (% Depletion), *x*₂ (light intensity), and categoric factor (MS Media Nutrient Composition) are shown in Table 3.

Effect of media composition on secondary metabolite production

The percentage depletion of KH₂PO₄ exhibited highly significant linear positive effect on grape callus anthocyanin content at *p*<0.01. Anthocyanin contents of the grape calli were significantly increased by increasing *x*₁ and *x*₂. TP concentrations of the grape calli were drastically increased by increasing all variables mainly *x*₁ and *x*₂ while there was a negative quadratic effect of *x*₂ which was almost negligible. Anthocyanin and TP contents of the plant cells remarkably contribute to their free radical scavenging activity and antioxidant capacity. The antioxidant capacity of the calli was also increased with all the increasing variables (*R*² = 0.9823) except for the slight negative quadratic effect of *x*₂ which displayed very similar pattern obtained for calli's TP contents. For

Table 3. Regression equations of the models for the anthocyanin total phenolic contents and antioxidant capacity of grape callus grown on different media and light regimes.

Response	Equation
Anth.	$Y(\text{KH}_2\text{PO}_4) = 10.85 + 0.41x_1 + 0.58x_2 - 5.70 \times 10^{-4} x_1x_2 + 7.97 \times 10^{-5} x_1^2 - 3.41 \times 10^{-3} x_2^2$
	$Y(\text{NH}_4\text{NO}_3) = 10.42 + 0.24x_1 + 0.57x_2 - 5.70 \times 10^{-4} x_1x_2 + 7.97 \times 10^{-5} x_1^2 - 3.41 \times 10^{-3} x_2^2$
TP	$Y(\text{KH}_2\text{PO}_4) = -822.14 + 17.27 x_1 + 59.59x_2 + 0.08 x_1x_2 + 0.73 x_1^2 - 0.38 x_2^2$
	$Y(\text{NH}_4\text{NO}_3) = -572.49 + 26.88 x_1 + 66.36x_2 - 0.15 x_1x_2 - 0.014 x_1^2 - 0.42 x_2^2$
AC	$Y(\text{KH}_2\text{PO}_4) = 15.05 + 0.62x_1 + 0.61x_2 + 7.41 \times 10^{-4} x_1x_2 + 1.04 \times 10^{-4} x_1^2 - 4.00 \times 10^{-3} x_2^2$
	$Y(\text{NH}_4\text{NO}_3) = 15.91 + 0.44x_1 + 0.58x_2 + 7.41 \times 10^{-4} x_1x_2 + 1.04 \times 10^{-4} x_1^2 - 4.00 \times 10^{-3} x_2^2$
CW	$Y(\text{KH}_2\text{PO}_4) = 8.33 - 0.12x_1 + 0.06x_2 - 1.67 \times 10^{-4} x_1x_2 + 1.12 \times 10^{-3} x_1^2 + 3.52 \times 10^{-4} x_2^2$
	$Y(\text{NH}_4\text{NO}_3) = 7.86 - 0.10x_1 + 0.05x_2 - 1.67 \times 10^{-4} x_1x_2 + 1.12 \times 10^{-3} x_1^2 + 3.52 \times 10^{-4} x_2^2$

Anth., Anthocyanin content (mg/kg); **TP**, total phenolic contents (mg/kg); **AC**, antioxidant capacity (% inhibition); **CW**, cell weight (g); x_1 , % Depletion (coded values, ranging from -1 to 1, which correspond to 0 to 75 respectively); x_2 , light intensity (coded values, ranging from -1 to 1, which correspond to 30 to 120, respectively).

the anthocyanin contents of the grape calli, the increasing order of magnitude of their effect was $x_2 > x_1$, for the TP contents $x_2 > x_1$, and for the antioxidant capacity $x_1 > x_2$.

The depletion of NH_4NO_3 also significantly contributed to the stimulation of secondary metabolite biosynthesis in grape callus, however at a smaller rate in comparison to what was observed in the calli grown on KH_2PO_4 limited media. The x_1 and x_2 variables affected the metabolite accumulation in grape calli grown on NH_4NO_3 limited media at the same pattern to those of grown on KH_2PO_4 depleted media, although with lower magnitudes as shown in Figures 1 and 2 and Tables 1 and 2, except for TP content. Both x_1 and x_2 exhibited positive influence on anthocyanin - total phenolic contents, and antioxidant capacity (Table 3). However, it should be noted that quadratic effect of x_2 was the sole negative effect on TP content of the cultures obtained from KH_2PO_4 depleted media while $x_1 - x_2$ interaction, and the quadratic effects of both $x_1 - x_2$ negatively affected the TP content in calli from the NH_4NO_3 limited media. The F values and R^2 coefficients of the models were above 1 and 0.90, respectively, showing the adequacy of the model (Table 4).

Response surface graphs (Figures 1 and 2) show the influence of light intensity and KH_2PO_4 or NH_4NO_3 depletion on metabolite content and antioxidant activity of grape calli. Optimization of nutrient media composition (concentration and type of nitrate ions, phosphate ions, microelements, carbon source, growth regulators) is one of the most effective ways of the induction of secondary metabolite biosynthesis in various tissue and cell culture systems. Red beet hairy root cultures grown on phosphate and/or nitrate limited media displayed elevated metabolite production potential (Taya et al., 1994; Mukundan et al., 1999; Akita et al., 2002). Our data shows that both depletion of KH_2PO_4 and NH_4NO_3 positively affected the metabolite concentration of calli and antioxidant capacity of their extracts (Figures 1 and 2). The highest anthocyanin - TP contents, and antioxi-

dant activity was detected in the grape calli grown on 75% KH_2PO_4 depleted medium under $75 \mu\text{mol s}^{-2} \text{m}^{-1}$. Calli grown on 75% KH_2PO_4 depleted media finally displayed 1.35 fold higher anthocyanin content, 2.22 fold higher TP content, and 1.25 fold higher antioxidant capacity in comparison to those of grown on 75% NH_4NO_3 depleted media under the same light regime.

Light dependent secondary metabolite production

Initially, we have cultured wounded grape leaf explants in the dark for callus induction and monitoring possible pigmentation and phenolics accumulation. Supplementation of the MS media with 1 mg L^{-1} NAA and 0.1 mg L^{-1} BA was necessary to induce callus formation and growth. Callus induction from grape leaves was accomplished in the dark although there was no visible anthocyanin accumulation in grape calli (Figure 3a). We have applied different light regimes (from 30 to $120 \mu\text{mol s}^{-2} \text{m}^{-1}$) to the grape calli obtained from the leaf explants cultured in the dark for the stimulation of biosynthesis of anthocyanins and the other phenolics. On transferring callus cultures to light in production medium significant secondary metabolite production was observed (Figure 3b). Highest metabolite accumulation was detected on the 21st day (data not shown) and thus a three week time period was selected for the evaluation of the light effect on the natural compound accumulation in grape calli. Increasing light intensity initially favored the anthocyanin and TP accumulation that brought about a significant increase in antioxidant capacity (Figures 1 and 2). However, further elevation of light intensity resulted in a reduction of those metabolites' concentration in the grape cells. Light-induced secondary metabolite production in plant cells is species-specific and significantly affected by light source, intensity and duration. Light-induced secondary metabolite production was reported in callus cultures of many species such as *Fragaria ananassa* (Nakamura et al.,




Figure 1 Three dimensional plot depicting 1a) anthocyanin accumulation, 1b) total phenolic compound concentration, 1c) antioxidant capacity, 1d) cell weight as a function of light intensity and KH₂PO₄ depletion.

1999), *Prunus cerasus* L. (Blando et al., 2004), *Vaccinium macrocarpon* Ait (Madhavi et al., 1995), *Centaurea cyanus* (Kakegawa et al., 1987), and *Eleutheretococcus senticosus* (Shohael, 2006). According to the results obtained in this study, exposure of calli to light strongly induced anthocyanin and phenolics biosynthesis that is linked to increased antioxidant capacity and the highest metabolite production was detected in grape calli grown under 75 $\mu\text{mol s}^{-2} \text{m}^{-1}$ (Figures 1 and 2). It should be noted that elevation of light intensity to 120 $\mu\text{mol s}^{-2} \text{m}^{-1}$ causes a significant reduction in anthocyanin - phenolics accumulation and a decrease in antioxidant activity in grape cells. It was observed that light intensities above 75 $\mu\text{mol s}^{-2} \text{m}^{-1}$ impair cell formation and development which could be

linked to callus vitrification and a decrease in metabolite accumulation by affecting the viability of calli. Similar results were obtained by Chan et al. (2010). They found that *Melastoma malabathricum* cell cultures display highest anthocyanin accumulation when exposed to moderate light intensities (301 to 600 lux). In contrast, a significant decrease in measurable anthocyanin concentration was observed in the calli grown under higher light intensity (900 to 1200 lux). Chalcone synthase (Chs) gene is one of the key genes controlling the phenylpropanoid pathway that is responsible for anthocyanin biosynthesis and accumulation of Chs mRNA transcripts is closely related to elevation of anthocyanin concentration within the cell. Parsley Chs gene's promoter region contains several *cis*-acting elements responsible




Figure 2. Three dimensional plot depicting 2a) anthocyanin accumulation, 2b) total phenolic compound concentration, 2c) antioxidant capacity, 2d) cell weight as a function of light intensity and NH₄NO₃ depletion.

for light-mediated activation of Chs mRNA transcription (Schulze-Lefert et al., 1989). However, Chs transcription is strongly and exclusively stimulated by only low fluence and short wavelength light (Chappel and Hahlbrock, 1984). Arabidopsis cells respond to low level of UV-light irradiance by stimulating the Chs transcription without apparent oxidative damage (Christie and Jenkins, 1996; Frohmeyer et al., 1997) light irradiation may cause an increase in the cellular anthocyanin concentration whereas cell growth might be suppressed (Zhang et al., 2002). Over exposure of plant cells to light might cause suppression of Chs transcription and an oxidative stress leading to oxidative damage (Green and Fluhr, 1995) those might be the major factors causing a reduction in anthocyanin accumulation observed in our study. Therefore, under the light and nutrient conditions applied

in this study, the best growth conditions to obtain highest secondary metabolite by using grape cell culture cv. Bogazkere were as follows, 75 $\mu\text{mol m}^{-2}\text{s}^{-1}$ light intensity with 75% KH₂PO₄ depleted MS media.

Effects of media composition and light intensity on the cell weight

Grape cell proliferation was determined for all the calli placed on various KH₂PO₄ or NH₄NO₃ depleted media used in this study under different light regimes and expressed on the gram basis. Equal amount of grape callus (100 mg) was located on the media for culture initiation and further analysis of the effect of KH₂PO₄ or NH₄NO₃ concentrations on the cell growth under certain

Table 4. Statistical evaluation and validation tests applied to regression equations.

Parameter	Regression	Residual	Total	R ²	Lack of Fit	Pure error
Anth.						
Sum of squares	2197.06 ^a	104.62	2301.68	0.9545	102.80 ^a	1.82
Freedom degrees	8	17	25	-	9	8
Mean squares	274.63	6.15	-	-	11.42	0.23
F value	44.63	-	-	-	50.32	-
TP						
Sum of squares	7.21x10 ⁷ ^a	6.06x10 ⁶	7.82x10 ⁷	0.9225	6.06x10 ⁶ ^a	3538
Freedom degrees	8	17	25	-	9	8
Mean squares	9.01x10 ⁶	3.56x10 ⁵	-	-	6.73x10 ⁵	442.25
F value	25.30	-	-	-	1521.21	-
AC						
Sum of squares	6860.71 ^a	123.79	6984.50	0.9823	117.59 ^a	6.20
Freedom degrees	8	17	25	-	9	8
Mean squares	857.59	7.28	-	-	13.07	0.78
F value	117.77	-	-	-	16.86	-
CW						
Sum of squares	73.44 ^a	6.49	79.93	0.9189	6.48 ^a	1.82x10 ⁻³
Freedom degrees	8	17	25	-	9	8
Mean squares	9.18	0.38	-	-	0.72	2.28x10 ⁻⁴
F value	24.06	-	-	-	3166.96	-

Anth., Anthocyanin (mg/kg); TP, total phenolic (mg/kg); AC, antioxidant capacity (% inhibition); CW, cell weight (g). ^aSignificant for $\alpha = 0.01$.

light intensities. Cell weight was determined after three weeks of time period. KH₂PO₄ or NH₄NO₃ depletion deteriorated cell proliferation and negatively correlated with the culture growth. 75% depletion of KH₂PO₄ resulted in almost four times less cell growth weight (from 6.8 to 1.8 g) compared to that of control samples (0% KH₂PO₄ depletion) under highest light regime (120 $\mu\text{mol m}^{-2}\text{s}^{-1}$) (Figures 1 and 2). Similarly, grape cells grew approximately two times less on 75% NH₄NO₃ depleted media in comparison to the control samples growing under the same light regimes. According to the results obtained in this study; even though depletion of KH₂PO₄ or NH₄NO₃ positively contributes to accumulation of phenolic compounds, optimisation studies indicated a critical balance zone shown in contour plots (Figure 4a, b) for each of KH₂PO₄ or NH₄NO₃ depletion for the most optimum production efficiency for certain phenolic compounds while keeping in mind callus viability and cell weight which are also two important factors determining the best conditions of cellular production of industrially important natural compounds.

Light intensity was also another significant determinant of total concentrations of the analyzed compounds and final cell weight. Higher light intensities (75 $\mu\text{mol m}^{-2}\text{s}^{-1}$) remarkably stimulated anthocyanin - TP biosynthesis in comparison to that of grown under lower light regimes (30 $\mu\text{mol m}^{-2}\text{s}^{-1}$). Results indicated detrimental effect of light

intensities above 100 $\mu\text{mol m}^{-2}\text{s}^{-1}$ on the cell growth and brought about development of smaller cultures that was considered inefficient regarding metabolite accumulation on the basis of measured cell weight.

In conclusion, the grape callus cultures obtained with the optimal conditions displayed a higher TP and anthocyanin content together with greater antioxidant capacity. Cell viability and weight was limiting factor for our study interfering the potential increment of light intensity or further depletion of media components, namely KH₂PO₄ or NH₄NO₃. Healthy and viable callus formation and growth was the primary concern for establishing cell cultures and concluding most optimal conditions those will be recommended for bioproduction of natural compounds under mentioned conditions. Optimisation plot containing variables and responses are shown in Figure 4a, b. An increase in light intensity and depletion of both media components induced secondary metabolite production and intersection of those responses with cell weight determined the optimal contour plot indicating optimal zone showing most optimum variable factors for the best culture growth and secondary metabolite biosynthesis. Significant variables were utilised to generate response surface plots and their contours against each response. A multi-response optimization technique was employed to determine the optimum combination of KH₂PO₄ or NH₄NO₃ concentra-




Figure 3. a, Grape calli grown on MS media in the dark; b, pigmentation in grape calli when exposed to light ($75 \mu\text{mol m}^{-2} \text{s}^{-1}$)

tions and light intensity for the most optimum conditions of callus growth accumulating anthocyanin and phenolic compounds. The main criteria constraint optimization were growth and viability to be as high as possible with high antioxidant capacity, anthocyanin and phenolic compound content combined with improved concentrations of individual natural compounds as determined by GC-FID system.

The optimum culture growth conditions can then be determined by superimposing the contour plots of relevant responses. The contour plots indicated that 30% depletion of KH_2PO_4 or 50% depletion of NH_4NO_3 from

media exposed to 65 or $75 \mu\text{mol m}^{-2} \text{s}^{-1}$, respectively, light intensities represented the best growing conditions for the grape cultures.

Identification of individual phenolic compounds found in callus cultures

One of the main objectives of this study was to improve the concentrations of industrially important phenolics found in grape cultures as they are beneficial to human health and might be used as functional food ingredients.




Figure 4. Contour plots showing the effects of light intensity - KH₂PO₄ depletion (4a) and light intensity-NH₄NO₃ (4b) on the variables.

Amount of phenolic compounds found in grape cell cultures were analyzed and individually identified by using a GC coupled to a FID detector system. We analysed the cell cultures grown on 37.5% KH₂PO₄ depleted media exposed to increasing light intensities from 0 to 75 $\mu\text{mol m}^{-2}\text{s}^{-1}$

¹. Six simple phenolic compounds were identified by using GC-FID system, namely gentisic acid, gallic acid, epicatechin, vanillin, hesperidin and catechin as shown in GC chromatogram (Figure 5). The amount of identified phenolic compounds increased parallel to increasing light intensities.

The most prevalent phenolic compound was catechin and its amount increased from 29.40 to 137.80 (mg/kg) with the elevation of the light intensity up to 75 $\mu\text{mol m}^{-2}\text{s}^{-1}$. The amounts of the identified natural compounds were shown in **Table 5**.




Figure 5. Simple phenolic compounds determined using GC-FID system. 1, Gentisic acid; 2, gallic acid; 3, epicatechin; 4, vanillin; 5, hesperidin; 6, catechin.

Table 5. Simple phenolic compounds detected by GC-FID system and concentration of phenolic compounds at different light intensities.

Simple phenolic compound (mg/kg)	Light Intensity ($\mu\text{mol m}^{-2}\text{s}^{-1}$)		
	0	37.5	75.0
Gentisic acid	0.6	3.21	5.2
Gallic acid	7.2	12.09	16.13
Epicatechin	13.17	18.26	21.54
Vanillin	0.46	6.17	18.5
Hesperidin	0.24	0.43	0.59
Catechin	29.4	92.4	137.8

REFERENCES

- Akita T, Hina Y, Nishi T (2002). New medium composition for high betacyanin production by a cell suspension culture of table beet (*Beta vulgaris* L.). *Biosci. Biotech. Biochem.* 66:902-905.
- Blando F, Gerardi C, Nicoletti I (2004). Sour Cherry (*Prunus cerasus* L.) Anthocyanins as ingredients for functional foods. *J. Biomed. Biotechnol.* 5:253-258.
- Blois MS (1958). Antioxidant determinations by the use of a stable free radical. *Nature*, 181:1199-1200.
- Chan LK, Koay SS, Boey PL, Bhatt A (2010). Effects of abiotic stress on biomass and anthocyanin production in cell cultures of *Melastoma malabathricum*. *Biol. Res.* 43:127-135.
- Chappel J and Hahlbrock K (1984). Transcription of plant defense genes in response to UV light or fungal elicitor. *Nature*. 311:76-78.
- Christie J and Jenkins G (1996). Distinct UV-B and UV-A/blue light signal transduction pathways induce chalcone synthase gene expression in Arabidopsis cells. *Plant Cell* 8:1555-1567.
- De Bellis L, Nicoletti I, Giovinazzo F, Blando F, Scardino AP (2005). Characterization of in vitro anthocyanin-producing sour cherry (*Prunus cerasus* L.) callus cultures. *Food Res. Int.* 38:937-942.
- Fresco P, Borges F, Marques MPM, Diniz C (2010). The anticancer properties of dietary polyphenols and its relation with apoptosis. *Curr. Pharm. Des.* 21:114-134.
- Frohnmeier H, Bowler C, Schaefer E (1997). Evidence for some signal transduction elements involved in UV-light dependent responses in parsley protoplasts. *J. Exp. Bot.* 48:739-750.
- Gamborg O, Miller R, Ojima K (1968). Nutrient requirement suspensions cultures of soybean root cells. *Exp. Cell Res.* 50:151-158.
- Georgiev V, Ilieva M, Bley T, Pavlov A (2008). Betalain production in vitro systems. *Acta Physiol. Plant.* 30:581-593.
- Giusti MM and Wrolstad RE (1996). Radish anthocyanin extract as a natural red colorant for maraschino cherries. *J. Food Sci.* 61: 688-694.
- Glaben WE, Wray V, Strack D, Metzger JW, Seitz HU (1992). Anthocyanins from cell suspension cultures of *Daucus carota*. *Phytochem.* 31:1593-1601.
- Green R and Fluhr R (1995). UV-B- induced PR-1 accumulation is mediated by active oxygen species. *Plant Cell* 7:203-212.
- Kakegawa K, Kaneko Y, Hattori E, Koike K, Takeda K (1987). Cell cultures of *Centaurea cyanus* produce malonated anthocyanin in UV light. *Phytochem.* 26:2261-2263.
- Keller M, Steel CC, Creasy GL (2000). Stilben accumulation in grapevine tissues: developmental and environmental effects. XXV. International Horticultural Congress, Part 4: Culture Techniques with Special Emphasis on Environmental Implications, ISHS Acta Horticulturae. 514:275-286.
- Kuhlmann I and Rohl C (2006). Phenolic antioxidant compounds produced by in vitro cultures of rosemary and their anti-inflammatory effect lipopolysaccharide activated microglia. *Pharm. Biol.* 6:401-410
- Loke WM, Proudfoot JM, Hodgson JM, McKinley AJ, Hime N, Magat M, et al. (2010). Specific dietary polyphenols attenuate atherosclerosis in apolipoprotein E-knockout mice by alleviating inflammation and endothelial dysfunction. *Arterioscler. Thromb. Vasc. Biol.* 30: 749-57.
- Madhavi DL, Smith MAL, Berber-Jimenez MD (1995). Expression of anthocyanins in callus cultures of cranberry (*Vaccinium macrocarpon*) *J. Food Sci.* 60:351-355.
- Mukundan U, Bhide V, Dawda H (1999). Production of betalains by hairy root cultures of *Beta vulgaris* L. In: Fu TJ, Singh G, Curtis WR (eds) *Plant cell and tissue culture for the production of food ingredients*. Kluwer Academic/Plenum Publishers, New York. pp. 121-127.
- Nakamura M, Takeuchi Y, Miyanaga K, Seki M, Furusaki S (1999). High anthocyanin accumulation in the dark by strawberry (*Fragaria ananassa*) callus. *Biotechnol. Lett.* 21:695-699.
- Nicoletti I, Bello C, De Rossi A, Corradini D (2008). Identification and quantification of phenolic compounds in grapes by HPLC-PDA-ESI-MS on a semimicro separation scale. *J. Agric. Food Chem.* 56:8801-8808.
- Ozden M, Demirel U, Kahraman A (2009). Effects of proline on antioxidant system in leaves of grapevine (*Vitis vinifera* L.). *Sci. Hort.* 119:163-168.
- Schulze-Lefert P, Becker-Andre M, Schulz W, Hahlbrock K, Dangi JL (1989). Functional architecture of the light-responsive chalcone synthase promoter from parsley. *Plant Cell.* 1:707-714.
- Shohael AM, Ali MB, Yu KW, Hahn EJ, Islam R, Paek KY (2006). Effect of light on oxidative stress, secondary metabolites and induction of antioxidant enzymes in *Eleutheretococcus senticosus* somatic embryos in bioreactor. *Process Biochem.* 41:1179-1185.
- Simoës C, Bizarri CHB, Cordeiro LS, Castro TC, Coutada LCM, Silva AJB, Albarello N, Mansur E (2009). Anthocyanin production in callus cultures of *Cleome rosea*: Modulation by culture conditions and characterization of pigments by means of HPLC-DAD/ESI-MS. *Plant Physiol. Biochem.* 47:895-903.
- Slinkard K and Singleton VL (1977). Total phenol analysis: Automation and comparison with manual methods. *Am. J. Enol. Viticult.* 28: 49-55.
- Szajdek A and Borowska EJ (2008). Bioactive Compounds and Health-Promoting Properties of Berry Fruits: A Review. *Plant Foods Hum. Nutr.* 63:147-156.
- Taya M, Yakura K, Kino-oka M, Tone S (1994). Influence of medium constituents on enhancement of pigment production by batch culture of red beet hairy root. *J. Ferment. Bioeng.* 77:215-217.
- von Sonntag C (2006) Free radical-induced DNA damage and its repair. A Chemical Perspective, Springer-Verlag, Berlin, Heidelberg.
- Wallace TC and Giusti MM (2008). Determination of color, pigment, and phenolic stability in yogurt system colored with nonacylated anthocyanins from *Berberis boliviana* L. as compared to other natural/synthetic colorants. *J. Food Sci.* 73:241-248.
- Wrolstad RE (1976). Color and pigment analyses in fruit products. In: *Agricultural Station Bulletin 624*, Oregon State University.
- Zhang W, Curtin C, Kikuchi M, Franco C (2002). Integration of jasmonic acid and light irradiation for enhancement of anthocyanin biosynthesis in *Vitis vinifera* suspension cultures. *Plant Sci.* 162:459-468.