

Full Length Research Paper

Sequence analysis of cereal sucrose synthase genes and isolation of sorghum sucrose synthase gene fragment

T. Sivasudha^{1*} and P. A. Kumar²

¹Department of Environmental Biotechnology, Bharathidasan University, Tiruchy-620 024, India.

²NRC on Plant Biotechnology, IARI, New Delhi, India-110 012, India.

Accepted 28 August, 2007

Sorghum (*Sorghum bicolor* (L.) Moench) is an important staple food for about 500 million people in semi arid regions of the world. Recently, sorghum has been identified as a main plant species for the comparative analysis of grass genomes and as a source of beneficial genes for agriculture. Recent studies have shown that there is conservation of gene order at the chromosome level in rice, sorghum and maize. Therefore, a high-resolution alignment between these genomes will be needed to utilize them constructively for sorghum gene discovery. Sorghum sucrose synthase gene fragment was amplified by primers designed at conserved exon position of cereal sucrose synthases. Sorghum sucrose synthase gene fragment I shared homology with other cereal sucrose synthase at the exon positions 6, 7, 8, 9 and 10. Sorghum sucrose synthase fragment II shared homology from exon 2 to 6.

Key words: Sorghum, sucrose synthase, multiple sequence alignment.

INTRODUCTION

Sorghum, a C₄ grass that diverged from maize just 15 million years ago, is the fifth most important cereal grown worldwide (Doggett, 1988). Recently, sorghum has been identified as a key plant species for the comparative analysis of grass genomes and as a source of beneficial genes for agriculture. Sorghum is relatively small genome (750 Mbp) (Arumuganathan and Earle, 1991) and incremental divergence from maize and rice (Doebley et al., 1990), make it ideally suited to aid the discovery and analysis of grass genes through comparative genomics. Sucrose synthase genes have been isolated from many of plants, mostly from starch storing plants such as maize (Werr et al., 1985; Hung et al., 1994), rice (Wang et al., 1992; Yu et al., 1992), wheat (Marana et al., 1990), potato (Salanoubat and Belliard, 1987), pea (Barratt et al., 2001) as well as *Arabidopsis thaliana* (Chopra et al., 1992; Martin et al., 1993).

Sucrose synthase [UDP-D-glucose (UDPG): D-fructose 2-glycosyl transferase, EC 2.4.1.13, *SuS*] catalyzes a

reaction of sucrose and UDP to form fructose and UDPG, the latter being a precursor of complex saccharide biosynthesis (Chourey and Nelson, 1976; Chourey et al., 1991; Carlson and Chourey, 1996). This study reports on multiple sequence alignment of sucrose synthase gene sequences from different cereals and isolation and sequencing of sucrose synthase gene fragment from sorghum using primers designed at their conserved exons.

MATERIALS AND METHODS

Multiple sequence alignment

Sucrose synthase gene sequences of various cereals like rice, maize, and barley were accessed from NCBI Genbank database and multiple sequence alignment was done using MegAlign programme of DNA star software.

Isolation of plant DNA

Two weeks old sorghum seedlings grown in ½ MS medium (Murashige and Skoog, 1962) were taken for total genomic DNA isolation. Sorghum (var. CSV 15) seeds were kept in 0.1% HgCl₂ for 10 min with intermittent shaking. After decanting 0.1% HgCl₂ solution, the sorghum seeds were thoroughly washed in sterile dis-

*Corresponding author. E-mail: sudacoli@yahoo.com. Tel: +91-431-2407088. Fax: +91-431-2407032.

Table 1. Nucleotide sequences of primers.

Primers	Sequence
GP1 F	5' GC GTCGAC CCAAGAGCTTGGTTTGGAGAAGG 3'
GP1 R	5' GC TCTAGA CTGTGAACTGGCATGAGAAGTGG 3'
GP2F	5' CTCCTCTCATCCCAATG 3'
GP2 R	5' CCCCTTCTCCAAACCAAG 3'

GP- Gene specific primer; R- Reverse primer; F- Forward primer.

tilled water two times and then placed in tissue culture bottle containing ½ MS medium. Sorghum seeds were kept at 28°C (BOD incubator) for germination. Sorghum genomic DNA was isolated by CTAB method (Dellaporta et al., 1983). The isolated genomic DNA was treated with RNase to make it RNA free. DNA quantitation was carried out by agarose gel analysis. 2 - 5 µl of sorghum genomic DNA was loaded in 0.7% agarose gel. Known amount of 1 Kb ladder was loaded as control in adjacent well. The quantity of DNA in the sample was estimated by comparison with the intensity of 3 kb band by eye judgment.

PCR

Primer sequences were designed at the conserved coding region to amplify a sucrose synthase gene fragment from sorghum (Table 1). Purification of amplified DNA from PCR reactions was performed using Qiaquick PCR purification kit (QIAGEN, Germany). pBlue-script and UA cloning vector (QIAGEN PCR Cloning Kit) was used to clone the PCR amplified product

Transformation of E. coli cells

One vial of competent cells was taken out from -70°C and thawed by keeping on the ice. 2 ml of plasmid DNA or 15 µl of ligation mixture was added and kept on ice for 30 min. The cells were subjected to heat shock at 42°C for 2 min, immediately plunged on ice for 5 min. To this 900 µl of Luria broth was added and kept at 37°C with shaking speed of 220 rpm for 1 h. The Cells were pelleted at 10,000 rpm for 1 min and resuspended in 100 µl of Luria broth, plated and incubated at 37°C for growth of transformed cells.

Cloning and sequencing

The recombinant clone was sequenced using T₇ and SP₆ reverse primers. Sequencing was carried out by Sanger dideoxy DNA sequencing method.

Results

Multiple sequence alignment

Sucrose synthase is found in all plant tissues and is found at high levels particularly in sink tissues. In monocotyledonous plants sucrose synthase is encoded by a small gene family. Sucrose synthase gene sequences of cereals such as rice, maize, and barley were taken from

NCBI Genbank data base and multiple sequence alignment was carried out using MegAlign programmer of DNA star software. Genes Suc S₂, RSS₁ and SUS₂ are highly conserved with respect to their exon and intron position and number. All the three genes have 16 exons. All the gene sequences compared have conserved gene sequences at their coding part and not in the non-coding region. 13th exon of rice RSS₂ is split in two. 14th exon of RS₂ is sharing homology with 16th exon of maize SuS₂ and rice RSS₁. In case of rice RSS₃ and Ss₂ of *Hordeum vulgare* the 3rd exon position is replaced with a new exon. This new exon of RSS₃ of rice and Ss₂ of *H. vulgare* does not share homology with 3rd exons of RSS₁, RSS₂ of rice, SucS₂, SuS₁ of maize, SS₂ of *Saccharum officinarum*. 5th exon of RSS₃ of rice and Ss₂ of *H. Vulgare* is missing in both. Various cereal sucrose synthase gene sequences taken for MSA are summarized in the table with their gene name, accession number, exon numbers, exon position and region of homology (Table 2)

Primer design and PCR amplification

As the sucrose synthase gene sequence is highly conserved among cereals especially at the coding region, from multiple sequence alignment, highly conserved exon positions were chosen to design primers and using those to amplify sucrose synthase gene fragment from sorghum. Schematic diagram showing the exon position where the primers were designed and sorghum sucrose synthase gene fragments were cloned is shown in Figure 1. The forward primer GP1F 5' GC GTC GAC CCA AGA GCT TGG TTT GGA GAA GG 3' was designed based on the region of homology at the exon position 6 and the reverse primer GP1R 5' GC TCT AGA CTG TGA ACT GGC ATG AGA AGT GG3' was designed based on the region of homology at the exon position 10. The purified sorghum genomic DNA was used for amplification. Various annealing temperatures ranging from 53 – 63°C were tested for amplification of the specific sequence. PCR reaction was run for 25 cycles and the reaction was electrophoresed on the 0.8% agarose gel. Among various annealing temperature tested, amplification was obtained at 56°C, which showed a single band of expected size of about 1 kb. No nonspecific amplification was observed (Figure 2). The amplified product was eluted from gel cloned in *Sal I* and *Xba I* pBluescript. To confirm the size of the insert, plasmid isolated from the transformed colony was digested with same two restriction enzymes. The expected fragment of ~1kb was obtained.

Cloning of sorghum sucrose synthase gene fragment

Sorghum sucrose synthase gene fragment cloned in pBluescript was sequenced and the sequence was aligned with other cereal sucrose synthase gene sequences, which were already taken for multiple sequence

Table 2. Gene name, accession number, exon numbers, exon position and region of homology different cereal sucrose synthases

S/N	Plant	Gene name	Access-ion No.	Exon No.	Alignment position	Exon position	Region of homology
1.	<i>Oryza sativa</i>	RSs ₁	X 64770	2	20-130	2882-2992	2899-2996
	<i>Oryza sativa</i>	RSs ₂	X 59046	2	20-120	2632-2770	2689-2765
	<i>Oryza sativa</i>	RSs ₃	L 03366	2	40-100	14-123	52-133
	<i>Zea mays</i>	SucS ₂	X 02382	2	10-110	2208-2321	2227-2325
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	2	10-110	1333-1443	1333-1443
	<i>Oryza sativa</i>	Sus ₂	L 39940	2	10-110	4191-4285	4205-4290
2.	<i>Oryza sativa</i>	RSs ₁	X 64770	3	870-1060	3404-3524	3398-3581
	<i>Oryza sativa</i>	RSs ₂	X 59046	3	880-990	3497-3627	3511-3630
	<i>Oryza sativa</i>	RSs ₃	L 03366	3 ^Δ	-	658-787	Missing
	<i>Zea mays</i>	SucS ₂	X 02382	4*	880-990	884-1035	857-872
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	3 ^Δ	700-820	1856-1985	1880-2015
	<i>Oryza sativa</i>	Sus ₂	L 39940	4*	950-990	2064-2215	2110-2150
3.	<i>Oryza sativa</i>	RSs ₁	X 64770	4	1090-1240	3614-3765	3607-4831
	<i>Oryza sativa</i>	RSs ₂	X 59046	4	1090-1240	3722-3823	3725-3826
	<i>Oryza sativa</i>	RSs ₃	L 03366	5 [∇]	119-1311	1142-1204	1142-1202
	<i>Zea mays</i>	SucS ₂	X 02382	4	1090-1240	3041-3192	3031-3189
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	5 [∇]	1150-1230	2304-2496	2297-2385
	<i>Oryza sativa</i>	Sus ₂	L 39940	4	1090-1230	4907-5058	4910-5059
4.	<i>Oryza sativa</i>	RSs ₁	X 64770	5	1360-1560	3849-4041	3846-4049
	<i>Oryza sativa</i>	RSs ₂	X 59046	5	1360-1460 1480-1540	3994-4186	3994-4074 4104-4173
	<i>Oryza sativa</i>	RSs ₃	L 03366	-	-	Missing ^α	-
	<i>Zea mays</i>	SucS ₂	X 02382	5	1366-1470	3271-3464	3271-3477
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	-	-	Missing	-
	<i>Oryza sativa</i>	Sus ₂	L 39940	5	1360-1560	5142-5334	5151-5352
5.	<i>Oryza sativa</i>	RSs ₁	X 64770	6 ^l	1650-1680 1690-1830	4042-4190	4136-4166 4178-4315 ⁰
	<i>Oryza sativa</i>	RSs ₂	X 59046	6 ^l	1720-1730	4312-4430	4316-4433
	<i>Oryza sativa</i>	RSs ₃	L 03366	6 ^l	1720-1800	1489-1607	1496-1586
	<i>Zea mays</i>	SucS ₂	X 02382	6 ^l	1710-1840	3615-3733	3604-3739
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	6 ^l	1720-1830	2834-2952	2839-2959
	<i>Oryza sativa</i>	Sus ₂	L 39940	6 ^l	1720-1830	5474-5602	5498-568
6.	<i>Oryza sativa</i>	RSs ₁	X 64770	7	1940-2160	4405-4621	4400-4630
	<i>Oryza sativa</i>	RSs ₂	X 59046	7	1840-2160	4550-4766	4549-4766
	<i>Oryza sativa</i>	RSs ₃	L 03366	7	1940-2040	1695-1911	1697-1914
	<i>Zea mays</i>	SucS ₂	X 02382	7	1950-2170	3832-4048	3829-4053
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	7	1950-2150	3039-3255	3046-3251
	<i>Oryza sativa</i>	Sus ₂	L 39940	7	1960-2160	5698-5914	5714-5924
7.	<i>Oryza sativa</i>	RSs ₁	X 64770	8	2310-2450	4704-4799	4698-4826
	<i>Oryza sativa</i>	RSs ₂	X 59046	8	2320-2420	4851-4946	4849-4949
	<i>Oryza sativa</i>	RSs ₃	L 03366	8	2320-2410	2022-2117	2024-2119
	<i>Zea mays</i>	SucS ₂	X 02382	8	2320-2430	4125-4219	4125-4223
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	8	2320-2400	3411-3768	3603-3769
	<i>Oryza sativa</i>	Sus ₂	L 39940	8	2320-2400	5997-6092	6001-6091

Table 2. Contd.

8.	<i>Oryza sativa</i>	RSs ₁	X 64770	9	2520-2700	4896-5069	4891-5071
	<i>Oryza sativa</i>	RSs ₂	X 59046	9	2530-2710	5028-5201	5029-5204
	<i>Oryza sativa</i>	RSs ₃	L 03366	9	2530-2670	2233-2406	2240-2404
	<i>Zea mays</i>	SucS ₂	X 02382	9	2550-2710	4315-4488	4315-4491
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	9	2530-2700	3595-3768	3603-3769
	<i>Oryza sativa</i>	Sus ₂	L 39940	9	2560-2700	9189-6362	6222-6372
9.	<i>Oryza sativa</i>	RSs ₁	X 64770	10	2810-2940	5154-5270	5155-5277
	<i>Oryza sativa</i>	RSs ₂	X 59046	10	2800-2930	5289-5905	5281-5408
	<i>Oryza sativa</i>	RSs ₃	L 03366	10	2810-2930	2514-2630	2496-2634
	<i>Zea mays</i>	SucS ₂	X 02382	10	2840-2920	4572-4688	4598-4680
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	10	2840-2910	3852-3968	3882-3952
	<i>Oryza sativa</i>	Sus ₂	L 39940	10	2830-2930	6447-6563	6466-6576
10.	<i>Oryza sativa</i>	RSs ₁	X 64770	11	3040-3210	5353-5519	5354-5526
	<i>Oryza sativa</i>	RSs ₂	X 59046	11	3040-3190	5489-5655	5486-5656
	<i>Oryza sativa</i>	RSs ₃	L 03366	11	3040-3200	2726-2892	2725-2892
	<i>Zea mays</i>	SucS ₂	X 02382	11	3040-3210	4798-4964	4800-4963
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	11	3070-3210	4054-4220	4078-4228
	<i>Oryza sativa</i>	Sus ₂	L 39940	11	3040-3200	6696-6812	6649-6819
11.	<i>Oryza sativa</i>	RSs ₁	X 64770	12	3240-3540	5066-5832	5591-5838
	<i>Oryza sativa</i>	RSs ₂	X 59046	12	3310-3450	5760-5984	5752-5819
	<i>Oryza sativa</i>	RSs ₃	L 03366	12	3310-3540	2971-3195	2974-3206
	<i>Zea mays</i>	SucS ₂	X 02382	12	3310-3480	5045-5269	5039-5210
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	12	3310-3540	4304-4528	4308-4533
	<i>Oryza sativa</i>	Sus ₂	L 39940	12	3310-3535	6901-7125	6903-7113
12.	<i>Oryza sativa</i>	RSs ₁	X 64770	13	4110-4130 4200-4230	6172-6490	6212-6242 6286-6316
	<i>Oryza sativa</i>	RSs ₂	X 59046	13 [□]	4210-4240 3640-3700	6071-6389	6295-6325 6077-6137
	<i>Oryza sativa</i>	RSs ₃	L 03366	13 [□]	3640-3700 3790-3840	3283-3601	3292-3352 3442-3475
	<i>Zea mays</i>	SucS ₂	X 02382	13	4090-4140 4240-4320	5404-5722	5417-54740 5547-5637
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	13 [□]	3640-3930	4607-4927	4616-4916
	<i>Oryza sativa</i>	Sus ₂	L 39940	13	3700-3720	7465-7783	7221-7251
13.	<i>Oryza sativa</i>	RSs ₁	X 64770	14 [■]	4700-4810	6621-6865	6649-6748
	<i>Oryza sativa</i>	RSs ₂	X 59046	14 [■]	4030-4050	4688-6732	6467-6497
	<i>Oryza sativa</i>	RSs ₃	L 03366	14	4120-4780 4860-4920	3688-3932	3734-3794 3859-3919
	<i>Zea mays</i>	SucS ₂	X 02382	14	4720-4920	5851-6095	5891-6078
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	14	4720-4920	5021-5265	5062-5249
	<i>Oryza sativa</i>	Sus ₂	L 39940	14 [■]	4720-4920	7914-8158	7964-8151
14.	<i>Oryza sativa</i>	RSs ₁	X 64770	15	5110-5120 5190-5260	6977-7115	6986-6993 7030-7080
	<i>Oryza sativa</i>	RSs ₂	X 59046	15 [■]	5380-5400	6827-9225	7814-7844
	<i>Oryza sativa</i>	RSs ₃	L 03366	15	5190-5230	4027-4165	4083-4105
	<i>Zea mays</i>	SucS ₂	X 02382	15	5280-5290	6258-6396	6250-6260
	<i>Hordeum vulgare</i>	Ss ₂	Y 15802	15	5190-5210	5374-5580	5428-5463
	<i>Oryza sativa</i>	Sus ₂	L 39940	15 [■]	5170-5220	5270-8408	8322-8353
15.	<i>Oryza sativa</i>	RSs ₁	X 64770	16	5370-5400 5490-5495	7213-7456	7143-7173 7224-7239

Table 2. Contd.

<i>Oryza sativa</i>	RSs ₂	X 59046	-	-	-	-
<i>Oryza sativa</i>	RSs ₃	L 03366	16	-	4262-4538	-
<i>Zea mays</i>	SucS ₂	X 02382	16	-	6516-6811	-
<i>Hordeum vulgare</i>	Ss ₂	Y 15802	-	-	-	-
<i>Oryza sativa</i>	Sus ₂	L 39940	16	-	8506-8553	-

Δ= 3rd exon of RSs₃ (*Oryza sativa*) and SS₂ (*Hordeum vulgare*) share homology with each other but does not share a region of homology with 3rd exon of other sucrose synthase gene sequences compared; * = 4th exon of RSs₃ and SS₂ share region of homology with 3rd exon of other sucrose synthase gene sequences compared; ∇= 5th exon of RSs₃ and SS₂ share region of homology with 5th exon of other sucrose synthase gene sequences compared.; α= RSs₃ and SS₂ does not have an exon to share homology with 5th exon of other sucrose synthase gene sequences compared. ! = 6th exon is highly conserved and shares highest region of homology in all the sucrose synthase gene sequences. □ = 13th exon of RSs₂ shares similarity with 13th exon of RSs₃ and Ss₂ and not with the RSs₁, SucS₂ and SVS₂. ■ = 14th exon of RSs₂ shares similarity with 14th exon of RSs₁ and Sus₂ and not with the RSs₃, Ss₂ and SucS₂. 0 = 15th exon of RSs₂ shares region of homology only with Sus₂ and not with other sucrose synthase gene sequences compared.




Figure 1. Schematic diagram showing the exon position where primers were designed and sorghum sucrose synthase gene fragments were cloned. GP- Gene specific primer; F - forward primer; R - reverse primer.




Figure 2. PCR Amplification of Sorghum sucrose synthase gene fragment using gene specific primers. Lane M 1 kb ladder. Lane 1 PCR Product of Sorghum sucrose synthase gene fragment I.

alignment. The cloned fragment shared homology in the region of exon 6, exon 7, exon 9 and exon 10 with Rss¹, Rss², and Rss³ of rice, SUC², SUC1 of maize, Ss₂ of *H.*

vulgare and *S. officinarum*, so the PCR amplified and cloned fragment in pBluescript vector was confirmed as sucrose synthase gene fragment of sorghum (Figure 3).

To clone an upstream sequence of sucrose synthase gene, one more forward primer GP₂F was designed at the exon position 2 and reverse primer GP2F was designed at exon position 6. PCR amplification was done at various annealing temperatures ranging from 50 to 60°C using sorghum genomic as template. The electrophoresis of PCR products was done on 0.8% agarose gel. The sharp single band of expected size of about 1.5 kb was obtained at 55°C annealing temperature.

Thermostable polymerase enzyme will add extra adenine nucleotide at the end of the amplifying strand at each cycle. This property was utilized to clone a Taq amplified PCR product in TA cloning vector. The 1.5 kb PCR amplified product was cloned in pDrive TA cloning vector. Sequencing was done by Sanger dideoxy DNA sequencing method and the nucleotide sequence was given in Figure 4. The sequence of 1.5 kb gene fragment was compared with other cereal sucrose synthase. It shared homology at exon No. 2, 3, 4, 5, and 6 of RSS₁, RSS₂, RSS3 of rice, Suc 2, Suc1C of maize, SS2 of *H.*

1 CAAGAGCTTG GTTTGGAGAA GGTTGGGGTG AACTGCAAA GCGCCGTACT TGACACAC
 61 CACTTGCTTC TTGACCTTCT TGAGGCCCT GATCCTGCCA ACTTGGAGAA GTTCCTTG
 121 ACTATACCAA TGATGTTNAA TGTTGTTATC CTGTCTCCTC ATGGCTACTT TGCCCAATC
 181 AATGTGCTTG GATACCTTGA CACTGGTGGC CAGGTACAGA AGCTTAGTGA TTTTTTTT
 241 AGACACTGAT TGTTTTCTT TTAGCTATTA TAGCTTTTAG GTTTCATT TGCAATCATT
 301 TTGCAGGTTG TGTACATTTT GGATCAAGTC CGTGCTTTGG AGAATGAGAT GCTTCTTA
 361 ATTAAGCAGC AAGGCCTTGA CATCACCCCG AAGATCCTCA TTGTATGTTT CATGTTTG/
 421 ACCATGTTG CTTTCTGAAC CTTTTCGTA TGCTGGNTGA AGTCATGCAT TCTGTGCT/
 481 AGGATGTTGC CAGTGAATA ATGTTAGAAA TGCAGGCCAA GCCTGACCTT ATCGTTGC/
 541 ACTACAGTGA TGGCAACCTA GTCGCCACTC TGCTCGCACA CAAGTTGGGA GTTACTC/
 601 TCTGTTTTGC TGTACATGAA TAATTGAGTT TTTTTTATA TAAAATTATT AAGTTCTCCA
 661 AATGCCTAAT AGTTTTGTAC ATACTTGCAG TGTACCATTG CCCATGCCTT GGAGAAAA/
 721 AAATACCCCA ACTCGGACAT ATACTTGGAC AAATTTGACA GCCAATACCA CTTCTCAT
 781 CAGTTCACAG TCTAGAGC

Figure 3. Nucleotide sequence of Sorghum sucrose synthase gene fragment I Gp1F-Gp1R. Highlighted portions are exon sequences sharing homology with cereal sucrose synthases.

1 CTCCTCTCAT CCCAATGAGC TGATTGCACT CTTCTCCAG GTGGGCATAC CAAAATATG1
 61 AACTTGCATT TCATTTCTG TACTGGAATT TGTTAATTTG GTATTCTCTT CATCCCAAAT
 121 GTAAACACGA GCATATGCAA CTTCTTTCTT GGTTTCTTTT GTTAAACACCA TCATGCATG
 181 TAATTGCTAT TCATCATCGA CTCATTGATC ATATATAATG ATTTTATGAT CAGGAGATTA
 241 TTGATTGTAA AGCATAGTGT TGCTGCTCTT CAGTTTTTGA AGCCTTTTGG TTTGATTAG
 301 ACAATTAGTT GATAAGACAG TATACTTTGT GTTACATCAT TTGGCAGATT GTTTGACTT
 361 AGTTGGTACA GTGCCATTTA ATATTTACAT CTTTCAGATC TAAATAGGAT ATAAAATGT
 421 CATCACAGCA GGGGAAAAGG TACATGATAT GAGATGTAAC ATCCATTTTA TTTGTGAA
 481 TCACTTTTAC AGGTATGTTA ACCAGGGCAA GGGAATGCTT CAGCGCCATC AACTGCTT
 541 TGAGTTTGAT GCCCTGTTTG ATAGTGACAA GGAGAAGTAT GCGCCC? (Gap of 250-300
 TTCG AAGACTTTCT
 601 TCGTGCTGCT CAGTAACACT GCTGAGATGC CTGCTTGAGT GATTGCGCCA GTCAACA/
 661 ACTTGTGCTG AGCTGATTTG AGCCTTCAAT GCTCGTTCCC TCGTCCTTCC ATGTCAAA/
 721 CCATGGGAAT GGAGTGCAAT TCTTAACCGA CACCTGTCTT CCAAGTTGTT ACAGGAC/
 781 GAGAGCCTGT ACCCATTGCT GAATTTCTC AAAGCCATA ACTACAAGGG CACGGTG.
 841 TTACAATTCA GAATCTTCCA AGCACATGCT TCACAATGGA TGATGACAAT ATTTATTTA/
 901 GAACTTTACA TAATCTGAAA ATGGATTAAA TGATGCCACC CAACTCCCTC ATTTGTAAC
 961 CTTTTTTTTT TCTGTTACAG ACGATGATGT TGAATGACAG AATTCAGAGC CTCCGTGGC
 1021 TCCAGTCATC CTTTAGAAAG GCAGAAGAGT ATCTACTGAG TGTCCCTCAA GACACTC
 1081 ACTCAGAGTT CAACCATAGG TGATTCATCA ATAAATTGTC CTTGCCATTT AACTTTGG
 1141 GAACTAGCAA ATGTATTAAC TGCTTGATG CCACCATGAT CTGCATTAGG TTCCAAG/
 1201 TTCCAAGAGC TTGGTTTGGG GAAGGGG

Figure 4. Nucleotide sequence of Sorghum sucrose synthase gene fragment II Gp2F-Gp2R.

vulgare and *S. officinarum*. So the 1.5 kb fragment was confirmed as sorghum synthase gene fragment. Highlighted portions are exon sequences sharing homology with cereal sucrose synthases

DISCUSSION

Sucrose synthase is found in all plant tissues and is found at high levels particularly in sink tissues. In monocotyledonous plants sucrose synthase is encoded by a small gene family. Sucrose synthase gene sequences of cereals such as rice, maize, sugarcane and barley were taken from NCBI Genbank database and multiple sequence alignment was carried out using MegAlign programme of DNA star software. Genes Suc S₂, RS₁ and SUS₂ are highly conserved with respect to their exon

and intron position and number. All the three genes have 16 exons. All the gene sequences compared have conserved gene sequences at their coding part and not in the non-coding region. 13th exon of rice RSS₂ is split in two. 14th exon of RS₂ is sharing homology with 16th exon of maize SuS₂ and rice RSS₁. In case of rice RS₃ and Ss₂ of *H. vulgare* the 3rd exon position is replaced with a new exon. This new exon of RSS₃ of rice and Ss₂ of *H. vulgare* does not share homology with 3rd exons of RSS₁, RSS₂ of rice, SucS₂, SuS₁ of maize, SS₂ of *S. officinarum*. 5th exon of RSS₃ of rice and Ss₂ of *H. vulgare* is missing in both. A long leader intron is characteristic of sucrose synthase genes (Werr et al., 1985, Wang et al., 1992; Yu et al., 1992). It is present in all the sucrose synthase genes isolated so far, except for ASuS I from Arabidopsis (Martin et al., 1993). The Cis elements present in the leader intron may be involved in gene re-

gulation.

Sorghum sucrose synthase gene fragment cloned in pBluescript was sequenced and the sequence was aligned with other cereal sucrose synthase gene sequences, which were already taken for multiple sequence alignment. The cloned fragment shared homology in the region of exon 6, exon 7, exon 9 and exon 10 with Rss¹, Rss² and Rss³ of rice, SUC², SUC1 of maize, Ss₂ of *H. vulgare* and *S. officinarum*. Therefore the PCR amplified and cloned fragment in pBluescript vector was confirmed as sucrose synthase gene fragment of sorghum.

It was decided to amplify the gene portion between exon 2 and exon 6. Primers were designed at exon 2 and exon 6 and 1.5 kb sucrose synthase gene fragment II was PCR amplified and cloned in pDrive cloning vector. The nucleotide sequence, which shared sequence similarity at their exon positions 2 to 6 with other cereal sucrose synthases. The position and the length of sequence homology were highlighted in the Figure 4. Exonic sequences not encoding protein (exon 1 and 16) of all the sucrose synthases taken for comparison do not exhibit conservation as that of coding region of the gene.

High level sequence divergence of the introns relative to that of exons was observed. This indicates that selection pressure against mutation abolishing or reducing the function of the gene. It was observed that sequences at exon/intron boundaries are highly conserved. These findings suggest that mutations in such a manner do not alter their ability to be recognized as introns. A very small percentage of sequence variations in exonic sequences encoding protein observed is an indication of the significance of evolution.

REFERENCES

- Arumuganathan E, Earle ED (1991). Nuclear DNA content of some important plant species. *Plant Mol. Bio. Repl* 9: 208-218.
- Barratt DHP, Barber L, Kruger NJ, Smith AM, Wang TL, Martin C (2001). Multiple distinct isoforms of sucrose synthase in pea. *Plant Physiol.* 127: 655-664.
- Carlson SJ, Chourey PS (1996). Evidence for plasma membrane associated forms of sucrose synthase in maize. *Mol. Gen. Genet.* 252: 303-310
- Chopra S, Delfavero J, Dolferus R, Jacobs M (1992). Sucrose synthase of *Arabidopsis*: genomic cloning and sequence characterization. *Plant Mol. Biol.* 18: 131-134
- Chourey PS, Chen YC, Miller ME (1991). Early cell degeneration in developing endosperms is unique to the *shrunken* mutation in maize. *Maydica*, 36: 141-146.
- Chourey PS, Nelson OE (1976). The enzymatic deficiency conditioned by the *shrunken-1* mutations in maize. *Biochem. Genet.* 14: 1041-1055
- Dellaporta SL, Wood J, Hicks JB (1983). A plant DNA miniprep: Version II. *Plant Mol. Biol. Rep.*, 1: 19-21.
- Doebley J, Durbin M, Golenberg EM, Clegg MT, Ma DP (1990). Evolution analysis of the large subunit of carboxylase (*rbcl*) nucleotide sequence among the grasses (Gramineae). *Evolution* 44: 1097-1108.
- Doggett H (1988). Sorghum. 2nd Edn. Longman Scientific and Technical Publishers, London, pp. 428-453.
- Hung XF, Nguge-Quoc B, Chourey PS, Yelle S (1994). Complete nucleotide sequence of the maize (*Zea mays* L.) sucrose synthase 2 cDNA. *Plant Physiol.* 104: 293-294
- Marana C, Garcia-Olmedo F, Carbonero P (1990). Different expression of two types of sucrose synthase-encoding genes in wheat in response to anaerobiosis, cold shock and light. *Gene*, 88: 167-172.
- Martin T, Frommer WB, Salanoubat M, Willmottzer R (1993). Expression of a *Arabidopsis* sucrose synthase gene indicates a role in metabolism of sucrose both during phloem loading and sink organs. *Plant J.* 4: 367-377
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant* 15: 473-497
- Salanoubat M, Belliard G (1987). Molecular cloning and sequencing of sucrose synthase cDNA from potato (*Solanum tuberosum* L.): preliminary characterization of sucrose synthase mRNA distribution. *Gene*, 60: 47-56
- Wang MB, Boulter D, Gatehouse P (1992). A complete sequence of the rice sucrose synthase-1 (RSs1) gene. *Plant Mol. Biol.* 19: 881-885.
- Werr W, Frommer WB, Mass C, Starlinger P (1985). Structure of the sucrose synthase gene on chromosome 9 of *Zea mays* L. *EMBO J.* 4: 1373-1380.
- Yu WP, Wang AY, Juang RH, Sung HY, Su JC (1992). Isolation and sequences of rice sucrose synthase cDNA and genomic NDA. *Plant Mol. Bio.* 18: 139-142.