

Full Length Research Paper

Protective roles of nitric oxide on antioxidant systems in tall fescue leaves under high-light stress

Yue-Fei Xu¹, Xiao-Ling Sun¹, Jing-Wei Jin² and He Zhou^{1*}

¹Department of Grassland Science, College of Animal Science and Technology, China Agricultural University, Beijing 100193, China.

²Department of Soil and Water Science, College of Resources and Environment, China Agricultural University, Beijing 100193, China.

Accepted 20 November, 2009

Nitric oxide (NO) is an important molecule involved in many physiological processes. In this study, the effect of NO on oxidative damage caused by high levels of light was investigated in tall fescue leaves. Tall fescue was developed in relative low light intensity ($100 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 21 days and then transferred to high light ($500 \mu\text{mol m}^{-2} \text{s}^{-1}$). Tall fescue leaves was supplied with NO donor, sodium nitroprusside (SNP), before high-light treatment to determine the physiological mechanisms of NO on tall fescue tolerance to high-light stress. Treatment of tall fescue leaves with $100 \mu\text{M}$ SNP before high-light stress resulted in alleviated light-induced electrolyte leakage, malondialdehyde and carbonyl contents in tall fescue leaves. The levels of H_2O_2 and superoxide radical ($\text{O}_2^{\cdot-}$) were reduced as well. Moreover, the activities of superoxide dismutase (SOD), catalase (CAT) and ascorbate peroxidase (APX) increased in tall fescue in presence of SNP under high-light stress. This pattern was reversed by application of NO scavenger, 2-(4-carboxy-2-phenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (PTIO) before light treatment. Pronounced increases of NO production were found in tall fescue leaves after exposure to high-light stress. The results suggested that high-light stress elevated NO level and that NO might act as a signalling molecule to enhance antioxidant enzyme activities, further protecting against injuries caused by high-light stress.

Key words: Antioxidant, high-light stress, nitric oxide, tall fescue.

INTRODUCTION

Light is essential for plant growth and development, but when plants are subjected to excessive light, active oxygen generation is increased (Asada, 2006), often resulting in photo-oxidative damages; thus light can also be one of the most deleterious environmental factor. Singlet oxygen (O_2^1), superoxide radicals ($\text{O}_2^{\cdot-}$) hydrogen peroxide (H_2O_2) and hydroxyl radicals (HO^{\cdot}) are collectively called reactive oxygen species (ROS). The effects of these ROS can cause the oxidation of lipids, proteins and enzymes necessary for the proper functioning of the chloroplast and the cell as a whole. To avoid ROS induced cellular injury, plants employ various antioxidative enzymes such as superoxide dismutase (SOD: EC 1.15.1.1), catalase (CAT: EC 1.11.1.6) and

ascorbate peroxidase (APX: EC 1.11.1.6) and low molecular weight antioxidants such as ascorbate (ASC), glutathione (GSH) and α -tocopherol (Mittler, 2002; Apel and Hirt, 2004) in the scavenging of these radicals.

Nitric oxide (NO) is an important signaling molecule involved in many plant physiological processes (Lamotte et al., 2005; Crawford, 2006; Corpas et al., 2008; Zhang et al., 2009). NO generates dual effects on plant, which is determined by its concentration, action site and physiological conditions of cells (Beligni and Lamattina, 1999). On one hand, NO participates in plant growth and signal transduction of acclimatization through the direct effect of effective molecular reaction or indirect effect of changing potential difference of cellular redox. On the other hand, higher concentration of NO can react with $\text{O}_2^{\cdot-}$ to produce many exogenous peroxy-nitrite, which could form peroxy-nitrous acid after protonation and damage structure and function of bio-macromolecule (Beligni and Lamattina, 2000; Martinez et al., 2000; Wendehenne et al., 2001; Neill

*Corresponding author. E-mail: zhouhe@cau.edu.cn. Tel: +86-10-62734122. Fax: +86-10-62734122.

et al., 2003). Several studies have shown that NO is involved in the regulation of plant responses to various environmental stresses. Increasing exogenous NO content under abiotic stresses may alleviate injury and enable plant cells to survive better (Song et al., 2006; Shi et al., 2007; Sun et al., 2007; Vital et al., 2008; Zhao et al., 2008). However, excessive NO released in plants and sustained high NO concentration might be cytotoxic (Arasimowicz and Floryszak-Wieczorek, 2007).

Tall fescue (*Festuca arundinacea*) is a widely used cold-season turfgrass species in China. Tall fescue is used as understorey turfgrass species and has leaves that are adapted to low light (Burner, 2003). As a result, it is often propagated under low light levels and can suffer damage if transferred to full sunlight. For this reason, it is important to obtain new light-tolerant cultivars. In a preliminary experiment, different varieties of tall fescue were found to exhibit distinct photoacclimation. Arid3 was not photobleached under high-light stress and this result suggested that Arid3 was light-tolerant turfgrass. High-light stress leads to enhanced ROS production. ROS-scavenging enzymes activities were reported to increase under high-light stress to mitigate oxidative damage (Burrill and Mackenzie, 2003; Ali et al., 2005a; Jiang et al., 2005). There have been no reports in which the influence of light levels on the production of ROS and antioxidant metabolism in tall fescue had been considered. On the other hand, NO was able to increase antioxidant enzymes activities under abiotic stresses to alleviate oxidative damage (Arasimowicz and Floryszak-Wieczorek, 2007), but there is little information on how NO is involved in the regulation of plant responses to light stress. Therefore, the objectives of this study were to determine whether antioxidant metabolism play in the acclimation of tall fescue following transfer of plants grown in the low light, to high light; and to elucidate the role of NO (applied exogenous NO or depleted endogenous NO) in alleviating light-induced oxidative damage in leaves of tall fescue.

MATERIALS AND METHODS

Plant materials and treatments

Seeds of tall fescue (*F. arundinacea* cv. Arid3) were obtained from Beijing Clover Seed and Turf CO., Ltd., China. Seeds were surface sterilized in 0.1% (w/v) sodium hypochlorite, rinsed several times in distilled water and germinated on moist filter paper at room temperature for 7 d. Seedlings were selected and placed into 5 L black plastic containers containing 4 L of solution. Each plastic container contained six plants. Seedlings cultured hydroponically in a continuously aerated nutrient solution containing 4 mM $\text{Ca}(\text{NO}_3)_2$, 4 mM KNO_3 , 1 mM KH_2PO_4 , 2 mM MgSO_4 , 46 μM H_3BO_3 , 10 μM MnSO_4 , 50 μM Fe-EDTA, 1.0 μM ZnSO_4 , 0.05 μM H_2MoO_4 , 0.95 μM CuSO_4 . The nutrient solution pH was adjusted close to 6.5 by adding H_2SO_4 or KOH. Nutrient solution was re-newed once a week. The plants were grown in a plant incubator at a day/night temperature 25/20°C, a relative humidity of 70%, a day/night regime of 14/10 h and a photosynthetic photo flux density (PPFD) at the height of the plants of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Light was provided by

a fluorescent lamp.

The leaves of plants that were maintained in relative low light (L, 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD) and high light (H, 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD) for the duration of the experiment were referred to as low-light-leaves and high-light-leaves, respectively, while those that developed fully in the low light and were then transferred to high light were referred to as transferred-leaves. Stress treatments were carried out after 21 d of pre-culture. Plants developed fully in the low light (100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD) and were transferred to high light (500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD). Sodium nitroprusside (SNP; Sigma, USA) was used as NO donor. The potassium salt of 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazole-1-oxyl-3-oxide (PTIO; Sigma, USA) was used as NO scavenger. 100 μM SNP and 200 μM PTIO were applied to tall fescue seedlings through the roots incubated (Laspina et al., 2005; Sun et al., 2007) in 4 L of nutrient solution (regenerated once a day) with high-light treatments. The 21-day-old seedlings were incubated in the solutions for 5 and 10 d at a day/night temperature 25/20°C, a relative humidity of 70%, day/night regime of 14/10 h. After 5 and 10 d of treatment, plants were harvested and frozen in liquid nitrogen and then stored at -80°C for further analysis.

Membrane permeability measurement

Membrane permeability (MP) was determined by the modified method of Song et al. (2006). The fresh leaves (0.5 g) were washed in deionized water and placed in petri dishes with 5 ml of deionized water at 25°C for 2 h. After the incubation, the conductivity was measured (C_1). Then, the samples were boiled for 20 min and conductivity was read again (C_2). Electrolyte leakage was expressed as a percentage of the total conductivity after boiling ($\text{MP} \% = C_1/C_2 \times 100$).

Analysis of lipid peroxidation

The level of lipid peroxidation was expressed as the amount of malondialdehyde (MDA) produced with a slight modification of the thiobarbituric acid method described by Buege and Aust (1978). Leaves (0.5 g) were homo-genized with a mortar and pestle in 10% trichloroacetic acid and then the homogenate was centrifuged at 4000 \times g for 30 min. A 2 ml aliquot of supernatant was mixed with 2 ml of 10% trichloroacetic acid containing 0.5% thiobarbituric acid. The mixture was heated at 100°C for 30 min. The absorbance of the supernatant was measured at 532 nm, with a reading at 600 nm subtracted from it to account for non-specific turbidity.

Analysis of oxidative damage to protein

Leaves (1.0) was homogenized in 50 mM potassium phosphate buffer (pH 7.0) containing 3 mM EDTA and 1 mM PMSF and leupeptin and centrifuged at 10000 \times g for 15 min at 4°C. Contaminating nucleic acids were removed by treatment with streptomycin sulfate and the oxidative damage to proteins was estimated as the protein carbonyl content, as determined by reaction with 2,4-dinitrophenylhydrazine (Reznick and Packer, 1994).

Determination of hydrogen peroxide and superoxide radical

Hydrogen peroxide content was measured according to Veljovic-Jovanovic et al. (2002). Leaves (0.5 g) were ground in liquid nitrogen and the powder was extracted in 2 ml 1 M HClO_4 in the presence of 5% PVP. The homogenate was centrifuged at 12000 \times g for 10 min and the supernatant was neutralized with 5 M K_2CO_3 .

(pH 5.6) in the presence of 0.1 ml of 0.3 M phosphate buffer (pH 5.6). The solution was centrifuged at $12000 \times g$ for 1 min and the sample was incubated for 10 min with 1 U ascorbate oxidase to oxidize ascorbate prior to assay. The reaction mixture contained 0.1 M phosphate buffer (pH 6.5), 3.3 mM 3-(dimethylamino) benzoic acid, 0.07 mM 3-methyl-2-benzothiazoline hydrazone, 0.3 U POX and 200 μ l supernatant. Changes in absorbance at 590 nm were monitored at 25°C. Superoxide radical production rate was determined by the modified method of Elstner and Heupel (1976). Leaves (1.0 g) were homogenized in 3 ml of 50 mM potassium phosphate buffer (pH 7.8) and centrifuged at $12000 \times g$ for 20 min. The incubation mixture contained 1 ml of supernatant, 1 ml of 50 mM potassium phosphate buffer (pH 7.8) and 1 ml of 1 mM hydroxyl aminonium chloride and the mixture was incubated in 25°C for 20 min. The mixture was subsequently incubated with 2 ml of 17 mM sulphaniic acid and 2 ml of 7 mM α -naphthyl amine at 25°C for 20 min. The final solution was mixed with an equal volume of ethyl ether and the absorbance of the pink phase was read at 530 nm. The production rate of superoxide radical was calculated based on a standard curve.

Antioxidant enzyme activity

Leaves (1.0 g) were homogenized with a mortar and pestle at 4°C in 5 ml 50 mM phosphate buffer (pH 7.0) containing 1mM EDTA, 1% PVP. The homogenate was centrifuged at $12000 \times g$ for 30 min at 4°C and the supernatant was collected for enzyme assays.

The activity of SOD was measured by nitroblue tetrazolium (NBT) method of Beauchamp and Fridovich (1971). One unit of SOD was defined as the amount of enzyme required to cause 50% inhibition of the reduction of NBT as monitored at 560 nm. The activity of CAT was determined by following the consumption of H_2O_2 at 240 nm ($E = 39.4 \text{ mM}^{-1} \text{ cm}^{-1}$) by the method of Aebi (1984). The activity of APX was measured according to Nakano and Asada (1981) by monitoring the rate of ascorbate oxidation at 290 nm ($E = 2.8 \text{ mM}^{-1} \text{ cm}^{-1}$). Enzyme activities were expressed on the basis of per unit protein weight. Protein content in the supernatant was determined using bovine serum albumin (BSA) as a standard (Bradford, 1976).

NO content determination

NO content determination was performed according to Murphy and Noack (1994) with some modifications. Fresh Leaves (0.5 g) were incubated with 100 units of catalase and 100 units of superoxide dismutase for 5 min to remove endogenous ROS before addition of 5 ml oxyhaemoglobin (5 mM). After 2 min incubation, NO concentrations were estimated by following the conversion of oxyhaemoglobin to methaemoglobin spectro-photometrically at 577 and 591 nm.

Statistical analysis

Each experiment was repeated at least three times. Values were expressed as means \pm SD. Statistical analyses were performed by analysis of variance (ANOVA). Means were separated using Duncan's multiple range test at 5% level of significance.

RESULTS

Effect of NO on membrane permeability (MP)

Membrane permeability (MP) in low-light-leaves was not significantly different from those found in high-light-leaves.

Transfer to high light caused a significant rapidly increased MP (at $P < 0.05$). Five days following transfer, MP remarkably increased by 35.5% and then showed little decrease after 10 days treatment (Figure 1A). Treatment of plant leaves with NO donor, SNP before high-light stress resulted in significant decrease of MP in transferred-leaves (at $P < 0.05$), especially 10 d after treatment. NO scavenger, PTIO were utilized to further clarify the role of SNP. The results showed that addition of PTIO enhanced MP to similar levels in transferred-leaves under high-light stress, indicating that PTIO scavenged endogenous NO.

Effect of NO on lipid peroxidation

As observed for membrane permeability, the levels of lipid hyperoxides in low-light-leaves was not significantly different from those found in high-light-leaves, but increased rapidly when plants were transferred to high light (at $P < 0.05$). High-light stress significantly increased MDA content in transferred-leaves both 5 and 10 d after treatment (Figure 1B). Supplementation with NO donor, SNP before high-light stress remarkably (at $P < 0.05$) reduced MDA content in transferred-leaves (at $P < 0.05$). However, when endogenous NO was removed, MDA content rose evidently, indicating that severe lipid peroxidation was caused.

NO prevents protein oxidative damage

Carbonyl content in low-light-leaves was not significantly different from that of high-light-leaves of tall fescue (Figure 1C). Carbonyl content of transferred-leaves increased following transfer to high light (at $P < 0.05$); with levels remaining significantly elevated for 5 d and then declined to normal levels after 10 d treatment. Application of exogenous NO dramatically prevented protein oxidative damage in transferred-leaves under high-light stress (at $P < 0.05$), especially 10 d after treatment.

Effect of NO on H_2O_2 and $O_2^{\cdot -}$ production

Compared to low-light-leaves, high-light stress caused significant accumulations (at $P < 0.05$) of H_2O_2 and $O_2^{\cdot -}$ production in transferred-leaves (Figure 2). Application of NO significantly reduced the accumulation of H_2O_2 and $O_2^{\cdot -}$ (at $P < 0.05$). Addition of PTIO remarkably increased H_2O_2 and $O_2^{\cdot -}$ levels in transferred-leaves of tall fescue under high-light stress (Figure 2).

Antioxidant enzyme activity

High-light stress had different effects on SOD, CAT and




Figure 1. Effect of NO on Membrane permeability (A), MDA content (B) Carbonyl content (C) in tall fescue leaves under high-light stress. Different treatments represents: H, L, LH, LH + SNP (100 μM) and LH+PTIO (200 μM). Values are means ± SD (n = 3). Bars with different letters are significantly different at the 5% level.

APX activity both on the 5 and 10 days after treatment in transferred-leaves (Figure 3A). On the 10th day after treatment, high-light stress greatly induced SOD, CAT




Figure 2. Effect of NO on levels of H₂O₂ (A) and superoxide radical (B) in tall fescue leaves under high-light stress. Values are means ± SD (n = 3). Bars with different letters are significantly different at the 5% level.

and APX activity (at P < 0.05). Under high-light stress, application of NO kept SOD and CAT with a relative high activity on the 5th day of treatment but slightly decreased APX activity on the 10th day treatment. Under normal condition, high-light-leaves had higher activities of antioxidant enzymes than low-light-leaves.

NO production

To further reveal the relationship between NO accumulation and high-light stress, NO production was measured. Transfer to high-light caused significantly increased endogenous NO production both 5 and 10 d after treatment. NO production increased by 199.6 and 179.2% in transferred-leaves, respectively. Application of NO scavenger PTIO merely reduced NO content (Figure 4).




Figure 3. Effect of NO on the activities of antioxidant enzymes of SOD (A), CAT (B) and APX (C) in tall fescue leaves under high-light stress. Values are means \pm SD ($n = 3$). Bars with different letters are significantly different at the 5% level.

DISCUSSION

Light is the major environmental constraint to growth and reproduction of understory turfgrass species. High light intensities might have contributed to a degradation of




Figure 4. NO production in tall fescue leaves under high-light stress. Values are means \pm SD ($n = 3$). Bars with different letters are significantly different at the 5% level.

chlorophyll contents, whereas low light intensities likely prevented breakdown of chlorophyll in leaves of tall fescue (Wherley et al., 2005). Thus, anatomical and physiological adaptations limited photosynthetic capacity and the ability to respond to increased irradiance and CO₂ of tall fescue grown continuously in low light intensities (Allard et al., 1991). Recent research indicated that the value of Fv/Fm has been used to understand photosynthesis affected by light intensities. The lower Fv/Fm in plants was due to photo-inhibition under high-light stress and turfgrasses grown under low light might suffer photo-inhibition when they were removed to high-light stress (jiang et al., 2005).

Photo-inhibition occurs when plants are exposed to a photosynthetic photon flux density (PPFD) higher than that required for the rate of CO₂ fixation, which further led to increased ROS generation (Asada, 2006). Many studies have reported that transfer from low light to high light caused enhanced H₂O₂ accumulation in plant leaves (Ali et al., 2005; Burritt and Mackenzie, 2003). The result in the present study showed that the levels of H₂O₂ and O₂⁻ increased in tall fescue transferred-leaves (Figure 2). The increased SOD activity may account for the increased accumulation of O₂⁻ in transferred-leaves (Figure 3A). Over production of ROS caused the oxidation of membrane lipids, proteins and enzymes necessary for the proper functioning of the chloroplasts and cells as a whole (Mittler, 2002). The increase in membrane permeability, MDA and carbonyl content under high-light stress indicated that high light induced oxidative damage on membrane lipid and proteins (Figure 1).

Previous studies have indicated that enhancements in the activities of activated oxygen scavenging enzymes generally accompany exposure to high-light stress (Foyer et al., 1997). For example, it has been reported that ROS-scavenging enzymes activities increased under highlight

stress in pumpkin, periwinkle (Logan et al., 1998a), begonias (Burritt and Mackenzie, 2003), seashore paspalum, bermudagrass (Jiang et al., 2005) and seven-son flower leaves (Liu et al., 2006). The result of this study indicated that tall fescue transferred-leaves exhibited increases in the activities of antioxidant enzymes (SOD, CAT and APX) both 5 and 10 d after highlight treatment (Figure 3). The difference indicated that the influence of high-light stress on the antioxidant enzymes are very complex and related to the plant treatment time, light intensity, plant tissues, plant species and genotype. SOD is a major scavenger of O_2^- , catalyzing the dismutation of superoxide to H_2O_2 and O_2 . CAT and APX are important H_2O_2 detoxifying enzymes. Increases in the activities of all antioxidant enzymes suggested that generation of ROS induced increased activities of antioxidant enzymes located in different cellular compartments (Logan et al., 1998b). Pronounced increase in antioxidant enzymes activities and the relatively low level of ROS in transferred-leaves indicated that tall fescue alleviated oxidative injuries through raising antioxidant enzymes activities to scavenge newly-produced ROS.

In many studies, it was found that NO could counteract oxidative damage and had protective effect against various stressful conditions (Song et al., 2006; Arasimowicz and Floryszak-Wieczorek, 2007; Shi et al., 2007; Sun et al., 2007; Tian et al., 2007; Vital et al., 2008; Zhao et al., 2008; Zhang et al., 2009). This study provided evidence that 100 μ M SNP treatments was able to protect transferred-leaves of tall fescue from MDA increase by delaying the accumulation of H_2O_2 and O_2^- . Addition of exogenous NO with SNP regulated the ROS levels in transferred-leaves of tall fescue relating to the activation of SOD, CAT and APX, prevented lipid peroxidation and proteins damage, whereas a reversed pattern was found with the supplementation of NO scavenger PTIO. Protective effect of NO might be related to two possibilities. On one hand, low concentration of NO induced the expression or activation of antioxidant enzymes, which was proven in plants (Laspina et al., 2005; Shi et al., 2007; Zhang et al., 2009). Recent research also proved that NO can induce APX and CAT antioxidant genes in *Arabidopsis* suspension cells (Huang et al., 2002). On the other hand, NO can detoxify ROS, such as reacting with O_2^- and generating peroxynitrite (ONOO \cdot). In the physiological pH range, ONOO \cdot is an unstable product. However, ONOO \cdot can be protonated and decomposed to a nitrate anion and a proton, or it can react with hydrogen peroxide to yield a nitrite anion and oxygen (Martinez et al., 2000; Wendehenne et al., 2001). Thus, NO might provide protection against oxidative stress. Transferred-leaves of tall fescue suffered severe peroxidative damage when endogenous NO was removed by PTIO, which further proved that NO plays an important role in photo-protection.

In conclusion, ROS metabolism is clearly important for tall fescue during acclimatization to high-light stress. The

acquisition of tolerance to high-light stress tall fescue transferred-leaves, are owed to the significantly increased antioxidant enzymes activities. As a bioactive antioxidant, NO protects tall fescue leaves against light-induced oxidative damage by reacting with ROS directly or inducing activities of ROS-scavenging enzymes.

ACKNOWLEDGEMENT

This work was supported by Key Projects in the National Science and Technology Pillar Program in the Eleventh Five-year Plan Period (No.2006BAD16B09-2).

REFERENCES

- Aebi H (1984). Catalase in vitro. *Methods Enzymol.* 105: 121-126.
- Ali MB, Hahn EJ, Paek KY (2005). Effects of light intensities on antioxidant enzymes and malondialdehyde content during short-term acclimatization on micropropagated *Phalaenopsis* plantlet. *Environ. Exp. Bot.* 54: 109-120.
- Allard G, Nelson CJ, Pallardy SG (1991). Shade effects on growth of tall fescue: II. Leaf gas exchange characteristics. *Crop Sci.* 31: 167-171.
- Apel K, Hirt H (2004). Reactive oxygen species: metabolism, oxidative stress and signal transduction. *Annu. Rev. Plant Biol.* 55: 373-399.
- Arasimowicz M, Floryszak-Wieczorek J (2007). Nitric oxide as a bioactive signalling molecule in plant stress responses. *Plant Sci.* 172: 876-887.
- Asada K (2006). Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141: 391-396.
- Beauchamp C, Fridovich I (1971). Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal. Biochem.* 44: 276-287.
- Beligni MV, Lamattina L (1999). Nitric oxide counteracts cytotoxic processes mediated by reactive oxygen species in plant tissues. *Planta*, 208: 337-344.
- Beligni MV, Lamattina L (2000). Nitric oxide stimulates seed germination and de-etiolation and inhibits hypocotyl elongation, three light-inducible responses in plants. *Planta*, 210: 215-221.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248-254.
- Buege JA, Aust SD (1978). Microsomal lipid peroxidation. *Methods Enzymol.* 52: 302-310.
- Burner DM (2003). Influence of alley crop environment on orchardgrass and tall fescue herbage. *Agron. J.* 95: 1163-1171.
- Burritt DJ, Mackenzie S (2003). Antioxidant metabolism during acclimation of *Begonia* \times *erythrophylla* to high light levels. *Ann. Bot.* 91: 783-794.
- Corpas FJ, Chaki M, Fernández-Ocaña A, Valderrama R, Palma JM, Carreras A, Begara-Morales JC, Airaki M, del Río LA, Barroso JB (2008). Metabolism of reactive nitrogen species in pea plants under abiotic stress conditions. *Plant Cell Physiol.* 49: 1711-1722.
- Crawford NM (2006). Mechanisms for nitric oxide synthesis in plants. *J. Exp. Bot.* 57: 471-478.
- Elstner EF, Heupel A (1976). Inhibition of nitrite formation from hydroxylammoniumchloride: a simple assay for superoxide dismutase. *Anal. Biochem.* 70: 616-620.
- Foyer CH, Lopez-delgado H, Dat JF, Scott IM (1997). Hydrogen peroxide- and glutathione-associated mechanisms of acclimatory stress tolerance and signalling. *Physiol. Plantarum*, 100: 241-254.
- Huang X, Rad U, Durner J (2002). Nitric oxide induces transcriptional activation of the nitric oxide-tolerant alternative oxidase in *Arabidopsis* suspension cells. *Planta*, 215: 914-923.
- Jiang YW, Carrow RN, Duncan RR (2005). Physiology acclimation of seashore paspalum and bermudagrass to low light. *Sci. Hortic.* 105: 101-115.

- Lamotte O, Courtois C, Barnavon L, Pugin A, Wendehenne D (2005). Nitric oxide in plants: the biosynthesis and cell signalling properties of a fascinating molecule. *Planta*, 221: 1-4.
- Laspina NV, Groppa MD, Tomaro ml, Benavides MP (2005). Nitric oxide protects sunflower leaves against Cd-induced oxidative stress. *Plant Sci*. 169: 323-330.
- Liu P, Yang YS, Xu GD, Hao CY (2006). Physiological response of rare and endangered seven-son-flower (*Heptacodium miconioides*) to light stress under habitat fragmentation. *Environ. Exp. Bot.* 57: 32-40.
- Logan BA, Demmig-Adams B, Adams WW III (1998a). Antioxidants and xanthophyll cycle-dependent energy dissipation in *Cucurbita pepo* L. and *Vinca major* L. upon a sudden increase in growth PPF in the field. *J. Exp. Bot.* 49: 1881-1888.
- Logan BA, Demmig-Adams B, Adams WW III, Grace SC (1998b). Antioxidants and xanthophyll cycle-dependent energy dissipation in *Cucurbita pepo* L. and *Vinca major* L. acclimated to four growth PPFs in the field. *J. Exp. Bot.* 49: 1869-1879.
- Martinez GR, Mascio PD, Bonini MG, Augusto O, Briviba K, Sies H, Maurer P, R othlisberger U, Herold S, Koppenol WH (2000). Peroxynitrite does not decompose to singlet oxygen ($^1\Delta O_2$) and nitroxyl (NO). *Proc. Natl. Acad. Sci. USA*, 97: 10307-10312.
- Mittler R (2002). Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7: 405-410.
- Murphy ME, Noack E (1994). Nitric oxide assay using haemoglobin method. *Methods Enzymol.* 233: 240-250.
- Nakano Y, Asada K (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22: 867-880.
- Neill SJ, Desikan R, Hancock JT (2003). Nitric oxide signalling in plants. *New Phytol.* 159: 11-35.
- Reznick AZ, Packer L (1994). Oxidative damage to proteins: spectrophotometric method for carbonyl assay. *Methods Enzymol.* 233: 357-363.
- Shi QH, Ding F, Wang XF, Wei M (2007). Exogenous nitric oxide protect cucumber roots against oxidative stress induced by salt stress, *Plant Physiol. Biochem.* 45: 542-550.
- Song LL, Ding W, Zhao MG, Sun BT, Zhang LX (2006). Nitric oxide protects against oxidative stress under heat stress in the calluses from two ecotypes of reed. *Plant Sci.* 171: 449-458.
- Sun BT, Jing Y, Chen KM, Song LL, Chen FJ, Zhang LX (2007). Protective effect of nitric oxide on iron deficiency-induced oxidative stress in maize (*Zea mays*). *J. Plant Physiol.* 164: 536-543.
- Tian QY, Sun DH, Zhao MG, Zhang WH (2007). Inhibition of nitric oxide synthase (NOS) underlies aluminum-induced inhibition of root elongation in *Hibiscus moscheutos*. *New Phytol.* 174: 322-331.
- Veljovic-Jovanovic S, Noctor G, Foyer CH (2002). Are leaf hydrogen peroxide concentrations commonly overestimated? The potential influence of artefactual interference by tissue phenolics and ascorbate. *Plant Physiol. Biochem.* 40: 501-507.
- Vital SA, Fowler RW, Virgen A, Gossett DR, Banks SW, Rodriguez J (2008). Opposing roles for superoxide and nitric oxide in the NaCl stress-induced upregulation of antioxidant enzyme activity in cotton callus tissue. *Environ. Exp. Bot.* 62: 60-68.
- Wendehenne D, Pugin A, Klessig DF, Durner J (2001). Nitric oxide: comparative synthesis and signaling in animal and plant cells. *Trends Plant Sci.* 6: 177-183.
- Wherley BG, Gardner DS, Metzger JD (2005). Tall fescue photomorphogenesis as influenced by changes in the spectral composition and light intensity. *Crop Sci.* 45: 562-568.
- Zhang LG, Zhou S, Xuan Y, Sun M, Zhao LQ (2009). Protective effect of nitric oxide against oxidative damage in *Arabidopsis* leaves under ultraviolet-B irradiation. *J. Plant Biol.* 52: 135-140.
- Zhao L, He JX, Wang XM, Zhang LX (2008). Nitric oxide protects against polyethylene glycol-induced oxidative damage in two ecotypes of reed suspension cultures. *J. Plant Physiol.* 165: 182-191.