

Full Length Research Paper

Effects of dietary L-arginine on orthodontic tooth movement in rats

Amir Mohammadi^{1*} and Ramin Azar²

¹Orthodontics Department, Dental Faculty, Tabriz Medical Science University, Golgasht Ave., Tabriz, Iran.

²Orthodontist, private practice.

Accepted 4 November, 2011

The purpose of this study was to investigate the effects of dietary L-arginine as nitric oxide (NO) precursor on orthodontic tooth movement in rats. 36 male ten-week old Wistar rats were randomly divided into experimental and control groups. The experimental group received 2% (w/w) dietary L-arginine in drinking water six days before the insertion of springs to elevate their blood level. On the seventh day, in both groups, maxillary incisors was moved by the insertion of springs and 12 days after insertion of springs, the rats were sacrificed, then the mesioincisal distance between maxillary incisors was measured. Afterwards, 12 and six rats from both groups were selected randomly for preparing histological section to count osteoclasts under a light microscope and for examining the surface area of root resorption lacunae under a scanning electron microscope, respectively. The data on the extent of orthodontic tooth movement and the number of osteoclasts were analyzed by independent sample t test and findings on root resorption were analyzed by using Mann-Whitney U test. The results showed that in L-arginine group, the orthodontic tooth movement ($p < 0.001$) and the number of osteoclasts ($p < 0.05$) were significantly higher when compared with the control group. However, there was no significant difference between the two groups in terms of the surface area of resorption lacunae.

Key words: L-Arginine, dietary, orthodontic tooth movement, nitric oxide, root resorption, osteoclast, nitric oxide synthase (NOS).

INTRODUCTION

An increase in the rate of orthodontic tooth movement (OTM) without observable damage to the tooth supporting structures can shorten the active treatment period. The OTM is mediated through a slow bone remodeling process. Osteoblasts and osteoclasts have a great role in this process. Shortening the duration of orthodontic treatment is a primary goal for the majority of the researchers involved in this field (Yamasaki et al., 1989).

Various factors and mediators are involved in the remodeling process, retarding or accelerating tooth movement, depending on the factor involved. An important signaling molecule which has recently been implicated in orthodontic tooth movement is nitric oxide

(NO), which is derived from the amino acid L-arginine by the nitric oxide synthase (NOS) enzyme in the human body (Marletta, 1993). Nilforoushan and Mansolon (2009) also showed that all NOS isoforms are involved in OTM with different expression patterns between tension and pressure sides, with nNOS being more involved in early OTM events. Both endothelial nitric oxide synthase (eNOS) and inducible nitric oxide synthase (iNOS) seem to be important regulators of bone remodeling during orthodontic force application (Tan et al., 2009). NO is a short-lived, soluble, free radical gas produced by a variety of cells and capable of mediating a number of various functions. The short half-life of NO indicates that its effects are regulated primarily by the rate of synthesis (Mitchell and Cotran, 2003).

The main activity of L-arginine in the human body is mediated through the production of NO molecule. L-arginine/NO pathway is involved in various biological activities in the cardiovascular, nervous, immune and

*Corresponding author. E-mail: amirortho@gmail.com. Tel: +989141161971. Fax: +984113356977.

other systems in the human body (Calver et al., 1993).

NO is an important role in inflammatory processes (Roberts et al., 2004), regulating the remodeling and turnover processes of bone (Ralston et al., 1994, 1995; Jung et al., 2003). Prescribing L-arginine for human and laboratory animals has different influences such as reducing stress (Gupta et al., 1995; Wu et al., 2010), lymphocyte activation (Ochoa et al., 2001; Tan et al., 2009), increases pregnancy chances in women and animal (Battaglia et al., 1999; Liu et al., 2011; Ren et al., 2011), enhances intestinal development and expression of vascular endothelial growth factor in weanling piglets (Yao et al., 2011), retardation of atherosclerosis progression in rabbits (Boger et al., 1997), decreases human blood pressure (Siani et al., 2000) and protects against nervous damage in rats (Sharma et al., 2005). Dietary L-arginine supplementation appears to affect the metabolism of lipoproteins and might alleviate some gastrointestinal functions, commonly seen in diabetes mellitus (Miguez et al., 2004).

Local injection of L-arginine and also nitro-L-arginine as NOS precursor adjacent to teeth in rats increases tooth movement and the number of osteoclasts, which is accomplished through NO synthesis and subsequent increase in cGMP level (Shirazi et al., 2002; Akin et al., 2004).

L-arginine is a non-essential amino acid and is under the influence of dietary intake. Under certain catabolic conditions such as trauma, infection and burns, it is an essential amino acid (Sy et al., 2006). After the dietary intake of this amino acid in laboratory animals, its blood level reaches the maximum after three days and after three weeks, its blood level is still higher than that in the control group (Jeremy et al., 1996). Dietary intake of L-arginine at a dose of 1.25 to 5% (w/w) in drinking water is not toxic for rats (Tsubuku et al., 2004).

Considering the fact that the local injection of L-arginine accelerates tooth movement and considering the importance of L-arginine/NO pathway in the orthodontic movement of teeth, this study was carried out to evaluate the effects of dietary L-arginine on the rate of orthodontic tooth movement, the number of osteoclasts in the area under pressure and root resorption in rats.

MATERIALS AND METHODS

36 approximately ten-week-old male Wistar rats with an average weight of 200 ± 22 g were selected and randomly divided into the control and L-arginine groups. To adapt the rats to the environmental conditions, they were kept in standard 12 h light and dark intervals at 24°C for a week.

The rats in both the experimental and control groups received drinking water and standard laboratory food *ad libitum* during the study period. The drinking water in the experimental group contained 2% (w/w) L-arginine as a solute (Jeremy et al., 1996; Boger et al., 1997; Miguez et al., 2004). To elevate the blood level of L-arginine, the experimental group had access to this kind of drinking water six days before the placement of the tooth-moving springs until the day when the rats were sacrificed (12 days after

spring placement). The rats were monitored during the study and were weighed at the beginning and at the end of the study.

On the 7th day, the tooth-moving spring were placed on the maxillary incisors in both the experimental and control groups. Holes were prepared on maxillary incisors, and 30 g of reciprocal force was applied to the teeth with a spring bent from 0.35-mm stainless steel wire, modified with respect to Akin et al. (2004). The springs were placed on a grid and activated on a single arm with a plier. The force was measured with a gauge, and the springs were not reactivated during the experiment. Prior to spring placement, the rats were anesthetized using 85 mg/kg ketamine (Alfasan, Woerden, Holland) in a peritoneal procedure. After being sure of anesthesia depth and fixing the animals, a special device that is specifically designed for this study was used to open the mouths of the rats and the springs were placed (Figure 1).

Considering the results of the previous studies, 30 g force was used to move the maxillary incisors of the rats (King et al., 1991; Akin et al., 2004). This amount of force is less than 90 g force necessary to open the maxillary suture in rats (Chang et al., 1997).

12 days after spring placement, the rats were sacrificed with an overdose of the anesthetic agent. Then, the distance between the mesioincisal line angles of maxillary incisors in both the control and the L-arginine group was measured using digital calipers, with a measuring accuracy of 0.01 mm (Figure 2). Subsequent to measuring the amount of orthodontic movement, 12 rats were randomly chosen from each group and their premaxillae were removed along with the incisor teeth and placed in 10% formalin for four days for fixation so that the number of osteoclasts could subsequently be counted. Then, the specimens were placed in 10% nitric acid for two days for decalcification. After decalcification, the specimens were rinsed for 2 h and again placed in 10% formalin for three days for fixation. Finally, the specimens were placed in special molds in a specific direction. The specimens were finally placed in paraffin molds and were ready to be sectioned. The specimens were sectioned serially by a microtome at 5 μ m thicknesses perpendicular to the long axis of the incisor teeth in the bone. Five serial sections were prepared at 5 μ m thicknesses below the alveolar crest.

The specimens were stained with Hematoxylin and Eosin. To standardize the area for osteoclastic count, a tangential line was drawn on the histological plate on the upper border of the teeth (toward the nasal cavity) and perpendicular to the middle septum. Another line was drawn parallel to the line 2 mm away from that on the lower side of the teeth. The third line was drawn tangential to the external border of the alveolus and perpendicular to the two earlier mentioned lines.

The fourth line was drawn 1 mm away from the third line and parallel to it on the nasal septum side. This way, we had a 2 mm² surface area on the external side of the alveolus for each incisor tooth, which added up to 4 mm² of the alveolar surface on both sides for osteoclastic count (Figure 3).

Therefore in each rat, five sections and in each section the distal surfaces of the right and left incisors were evaluated. On the whole, ten pressure areas with a total surface area of 20 mm² were measured and evaluated in each rat. Osteoclastic count was carried out only on the internal surface of the alveolus, which was under pressure, and resorption lacunae with multinuclear cells were considered as osteoclasts. Two histologists counted the osteoclasts twice at different times under a light microscope.

The incisors in the six remaining rats from each group, which had not been histological studied, were prepared to evaluate the resorption lacunae under a scanning electron microscope. The images of the scanned specimens were stored at 30x magnification and then the surface areas of resorption lacunae were calculated in mm² using AutoCAD software and the results were compared between the experimental and control groups (Figure 4). To evaluate the normal distribution of data, one sample Kolmogorov-Smirnov analysis was used. Levene's test was used to evaluate the




Figure 1. Device for fixation and opening of rat's mouth.

homogeneity of variances.

Finally, independent sample t test was used to compare the rate of tooth movement, the number of osteoclasts and the weight of the animals before and after the experiment between the two groups. Mann-Whitney U test was used to compare the surface area of the resorption lacunae on the root surface which was under pressure (the average of resorption lacunae at distal surface of the left and right roots in each specimen).

RESULTS

The average weight of the rats prior to the experiment was 200 ± 22 g. There was no significant difference between the weights of the rats before the experiment or after the experiment between two groups.

The finding of the OTM in the L-arginine and control groups is presented in Table 1. As shown in Table 1, the rate of the OTM in L-arginine group exceeds that in the control group and the difference is statistically significant ($P < 0.001$), and also, the number of osteoclasts in L-arginine group was more than that in the control group and the difference was statistically significant ($P < 0.05$).

Scanning electron microscope evaluation of the maxillary incisors in rats in both groups to estimate the extent of root resorption revealed that the mean surface area of resorption lacunae difference between two groups was not statistically significant (Table 2).

DISCUSSION

In animals, arginine improves protein synthesis, micro-vascular development, immune function, antioxidant activity and cell proliferation by regulating some intracellular signaling pathway (Yin and Tan, 2010).

Various factors and mediators including prostaglandins, cAMP, interleukin 1-beta and neurotransmitters are involved in the orthodontic tooth movement (Davidovitch et al., 1975; Shirazi et al., 2002). A mediator which has recently attracted attention as being involved in orthodontic tooth movement and bone remodeling is NO, which is derived from the amino acid L-arginine in the human body by the action of the enzyme NOS (Ralston et al., 1995). This molecule is a key messenger involved in the synthesis of the cyclic guanosine monophosphate (cGMP) (MacIntyre et al., 1991; Hsu et al., 2003), subsequently increasing cellular activity and osteoclasts recruitment. NO can have a direct role in the activation and synthesis of other messengers such as prostaglandin E2 (PGE₂). Under mechanical stress, human fibroblasts and osseous cells demonstrate an increase in NO synthesis (Pitsillides et al., 1995; Nakago-Matsuo et al., 2000).

The impact of various factors on the orthodontic movement of the teeth in rats five to 14 days after the




Figure 2. Measurement of orthodontic tooth movement by digital calipers.

application of force has been investigated in various studies and it has been demonstrated that the difference in tooth movement between experimental and control groups is more significant from day seven onward (Yamasaki et al., 1984; Engstrom et al., 1988; Kale et al., 2004; Akin et al., 2004).

It is shown in this study that 12 days after force application, the rate of orthodontic tooth movement in oral L-arginine group was 1.6 times more than that in the control group ($P < 0.001$). It was concluded that dietary L-arginine accelerates orthodontic tooth movement in rats which is similar to local L-arginine as Shirazi et al. (2002) demonstrated in a research study. Hayashi et al. (2002) demonstrated a decrease in the OTM in group receiving NOS inhibitor as compared to the control group. Akin et al. (2004) in their study showed an increase in the tooth movement by the use of nitro-L-arginine as NOS precursor.

Movement of the teeth during orthodontic treatment involves osteoclastic resorption of the alveolar bone around the tooth root in the direction of the movement. Recruitment of osteoclasts and an increase in their number is considered as an important factor in the evaluation of the rate of orthodontic movement (Kaku et

al., 2001; Shirazi et al., 2002; Akin et al., 2004; Mavragani et al., 2005). In addition, the maximum osteoclastic activity and concentration during orthodontic force application is usually observed five to seven days after force application (Tanne et al., 1998; Ren et al., 2005). In this study, the premaxillae of 12 rats, which contained the central incisors, were prepared for osteoclastic count under a light microscope. The results demonstrate that the number of osteoclasts in L-arginine group was more than that in the control group ($p < 0.05$). This finding is consistent with the results of a study carried out by Akin et al. (2004) and Shirazi et al. (2002).

Considering the results of this study, the difference in the number of osteoclasts and the rate of orthodontic tooth movement can only be attributed to the presence of L-arginine in the drinking water in the experimental group.

Resorption is a cell-mediated process; various mediators and messengers are involved in the recruitment and differentiation of the cells in the periodontal ligament (PDL) during orthodontic tooth movement. Therefore it is logical to believe that cell-mediated factors in tooth movement can also influence root resorption (Rygh et al., 1986; Leiker et al., 1995).

The result demonstrate that the difference in root




Figure 3. Area of osteoclasts count.




Figure 4. SEM view of root and resorption lacunae.

Table 1. Comparison of orthodontic tooth movement, osteoclasts number and roots resorption in control and L-arginine groups, using independent sample t test.

Parameter	Control			L-arginine			Diff.	t	df	P*
	Number	Mean	SD	Number	Mean	SD				
Initial weights (g)	18	198.3	21.9	18	203.0	23	4.7	-0.063	34	0.523
Final weights (g)	18	216.0	22.2	18	215.7	24	-0.3	0.036	34	0.971
OTM (mm)	18	1.54	0.31	18	2.45	0.36	0.89	-7.99	34	0.000
Osteoclasts count (n/mm ²)	12	3.86	1.84	12	5.57	1.44	1.71	-2.52	22	0.019

*p<0.05 is significant.

Table 2. Comparison of roots resorption in control and L-arginine groups, using Mann-Whitney U test

Parameter	Control			L-arginine			Diff.	U	P*
	Number	Mean	SD	Number	Mean	SD			
Resorption lacunae surface (mm ²)	6	2.80	1.63	6	4.73	1.29	1.93	14.50	0.575

* p<0.05 is significant.

resorption between two groups was not statistically significant. This finding is contrary to the finding of the study carried out by Shirazi et al. (2002), in which root resorption was less in L-arginine group as compared to the control group.

The relationship between NO and OTM was shown in previous studies (Shirazi et al., 2002; Hayashi et al., 2002; Akin et al., 2004). In light of these studies and our findings, it can be said that NO and L-arginine as NO precursor increases the number of osteoclasts in compression site and result to increase in the rate of orthodontic tooth movement.

Conclusion

Adding 2% (w/w) L-arginine to the drinking water of rats during orthodontic movement of the teeth leads to the following results:

- 1) Increase in orthodontic tooth movement.
- 2) Increase in the number of osteoclasts around the tooth in the movement direction in rats.
- 3) No significant increase in root resorption during orthodontic tooth movement in rats.

REFERENCES

- Akin E, Gurton AU, Olmez H (2004). Effects of nitric oxide in orthodontics tooth movement in rats. *Am. J. Orthod. Dentofac. Orthop.* 126: 608-614.
- Battaglia C, Salvatori M, Maxia N, Petraglia F, Facchinetti F, Volpe A (1999). Adjuvant L-arginine treatment for in vitro fertilization in poor responder patients. *Hum. Reprod.* 14: 1690-1697.
- Boger RH, Bode-Boger SM, Brandes RP, Phivthong-ngam L, Bohme M, Nafe R, Mugge A, Frolich JC (1997). Dietary L-arginine reduces the

- progression of atherosclerosis in cholesterol-fed rabbits: comparison with lovastatin. *Circulation*, 96: 1282-1290.
- Calver A, Collier J, Vallance P (1993). Nitric oxide and cardiovascular control. *Exp. Physiol.* 78: 303-326.
- Chang HN, Garetto LP, Potter RH, Katona TR, Lee CH, Roberts WE (1997). Angiogenesis and osteogenesis in an orthopedically expanded suture. *Am. J. Orthod. Dentofac. Orthop.* 111: 382-390.
- Davidovitch Z, Shanfeld JL (1975). Cyclic nucleotide level in alveolar bone of orthodontically treated cats. *Arch. Oral. Biol.* 20: 567-574.
- Engstrom C, Granstrom G, Thilander B (1988). Effect of orthodontic force on periodontal tissue metabolism. A histologic and biochemical study in normal and hypocalcemic young rats. *Am. J. Orthod. Dentofac. Orthop.* 93: 486-495.
- Gupta V, Gupta A, Saggi S, Divekar HM, Grover SK, Kumar R (1995). Anti-stress and adaptogenic activity of L-arginine supplementation. *Evid. Based. Complement. Alternat. Med.* 2: 93-97.
- Hayashi K, Igarashi K, Miyoshi K, Shinoda H, Mitani H (2002). Involvement of nitric oxide in orthodontic tooth movement in rats. *Am. J. Orthod. Dentofac. Orthop.* 122: 306-309.
- Jeremy RW, McCarron H, Sullivan D (1996). Effect of dietary L-arginine on atherosclerosis and endothelium-dependent vasodilatation in the hypercholesterolemic rabbit. Response according to treatment duration, anatomic site, and sex. *Circulation.* 94: 498-506.
- Jung JY, Lin AC, Ramos LM, Faddis BT, Chole RA (2003). Nitric oxide synthase I mediates osteoclast activity in vitro and in vivo. *J. Cell Biochem.* 89: 613-621.
- Kaku M, Kohno S, Kawata T, Fujita I, Tokimasa C, Tsutsui K, et al., (Provide Complete Name) (2001). Effects of vascular endothelial growth factor on osteoclast induction during tooth movement in mice. *J. Dent. Res.* 80: 1880-1883.
- Kale S, Kocaderli I, Atilla P, Asan E (2004). The effects of 1,25 dihydroxy cholecalciferol and Prostaglandin E2 on orthodontics tooth movement. *Am. J. Orthod. Dentofac. Orthop.* 25: 607-614.
- King GJ, Keeling SD, McCoy EA, Ward TH (1991). Measuring dental drift and orthodontic tooth movement in response to various initial forces in adult rats. *Am. J. Orthod. Dentofac. Orthop.* 99: 456-465.
- Leiker BJ, Nanda RS, Currier GF, Howes RI, Sinha PK (1995). The effects of exogenous prostaglandins on orthodontic tooth movement in rats. *Am. J. Orthod. Dentofac. Orthop.* 108: 380-388.
- Liu XD, Wu X, Yin YL, Liu YQ, Geng MM, Yang HS, Wu GY (2011). Effects of dietary L-arginine or N-carbamylglutamate supplementation during late gestation of sows on the miR-15b/16, miR-221/222, VEGFA and eNOS expression in umbilical vein. *Amino Acids.* DOI10.1007/s00726-011-0936-9.

- MacIntyre I, Zaidi M, Towhidul Alam ASM, Datta HK, Moonga BS, Lidbury PS, Hecker M, Vane JR (1991). Osteoclastic inhibition: an action of nitric oxide not mediated by cyclic GMP. *Proc. Natl. Acad. Sci. USA.* 88: 2936-2940.
- Marletta MA (1993). Nitric oxide synthase structure and mechanism. *J. Biol. Chem.* 268: 12231-12234.
- Mavragani M, Brudvik P, Selvig KA (2005). Orthodontically Induced root and alveolar bone resorption: inhibitory effect of systemic doxycycline administration in rats. *Eur. J. Orthod.* 27: 215-225.
- Miguez I, Marino G, Rodriguez B, Taboada C (2004). Effects of dietary L-arginine supplementation on serum lipids and intestinal enzyme activities in diabetic rats. *J. Physiol. Biochem.* 60: 31-37.
- Mitchell RN, Cotran RS (2003). Acute and chronic inflammation. In: Kumar V, Cotran RS, Robbins SJ (Eds) *Robbins basic pathology.* Elsevier, p. 50.
- Nakago-Matsuo C, Matsuo T, Nakago T (2000). Basal nitric oxide production is enhanced by hydraulic pressure in cultured human periodontal ligament fibroblasts. *Am. J. Orthod. Dentofac. Orthop.* 117: 474-478.
- Nilforoushan D, Manolson MF (2009). Expression of nitric oxide synthases in orthodontic tooth movement. *Angle Orthod.* 79: 502-508.
- Ochoa JB, Strange J, Kearney P, Gellin G, Edean E, Fitzpatrick E (2001). Effect of L-arginine on the proliferation of T-lymphocyte subpopulations. *J. Parenter. Enteral. Nutr.* 25: 23-29.
- Pitsillides AA, Rawlinson SC, Suswillo RF, Bourrin S, Zaman G, Lanyon LE (1995). Mechanical strain-induced NO production by bone cells: a possible role in adaptive bone remodeling? *FASEB J.* 9: 1614-1622.
- Ralston SH, Todd D, Helfrich M, Benjamin N, Grabowski PS (1994). Human osteoblast-like cells produce nitric oxide and express inducible nitric oxide synthase. *Endocrinology.* 135: 330-336.
- Ralston SH, Ho LP, Helfrich MH, Grabowski PS, Johnston PW, Benjamin N (1995). Nitric oxide: a cytokine-induced regulator of bone resorption. *J. Bone Miner. Res.* 10: 1040-1049.
- Ren Y, Kuijpers-jagtman AM, Maltha JC (2005). Immunohistochemical evaluation of osteoclast recruitment during experimental tooth movement in young and adult rats. *Arch. Oral Biol.* 50: 1032-1039.
- Roberts WE, Huja S, Roberts JA (2004). Bone modeling: biomechanics, molecular mechanisms, and clinical perspectives. *Semin. In Orthod.* 10: 123-161.
- Rygh P, Bowling K, Hovlandsdal L, Willias S (1986). Activation of the vascular system: a main mediator of periodontal remodeling in orthodontic tooth movement. *Am. J. Orthod.* 89: 453-458.
- Sharma HS, Badgaiyan RD, Alm P, Mohanty S, Wiklund L (2005). Neuroprotective effect of nitric oxide synthase inhibitors in spinal cord injury-induced pathophysiology and motor functions: An experimental study in the rat. *Ann. NY. Acad. Sci.* 1053: 422-434.
- Siani A, Pagano E, Iacone R, Iacoviello L, Scopacasa F, Strazzullo P (2000). Blood pressure and metabolic changes during dietary L-arginine supplementation in humans. *Am. J. Hyperten.* 13: 547-551.
- Shirazi M, Nilforoushan D, Alghasi H, Dehpour AR (2002). The role of nitric oxide in orthodontic tooth movement in rats. *Angle Orthod.* 72: 211-215.
- Sy BMC, Dweik EE, Dweik RA (2006). Arginine and nitric oxide. In: Shils ME, Shike M, Ross AC (Eds). *Modern nutrition in health and disease,* Lippincott Williams & Wilkins, Philadelphia, pp. 571-582.
- Tan BE, Li XG, Kong XF, Huang RL, Ruan Z, Deng ZY, Xie MY, Shinzato I, Yin YL, Wu G (2009). Dietary L-arginine supplementation enhances the immune status in early-weaned piglets. *Amino Acid.* 37: 323-331.
- Tan SD, Xie R, Klein-Nulend J, Van Rheden RE, Bronckers AL, Kuijpers-Jagtman AM, Von den Hoff JW, Maltha JC (2009). Orthodontic force stimulates eNOS and iNOS in rat osteocytes. *J. Dent. Res.* 88: 255-260.
- Tanne K, Yoshida S, Kawata T, Sasaki A, Knox J, Jones ML (1998). An evaluation of the biomechanical response of the tooth and periodontium to orthodontic forces in adolescent and adult subject. *Br. J. Orthod.* 25: 109-115.
- Tsubuku S, Hatayama K, Mawatari K, Smriga M, Kimura T (2004). Thirteen-week oral toxicity study of L-arginine in rats. *Int. J. Toxicol.* 23: 101-105.
- Wu X, Yin YL, Li TJ, Wang L, Ruan Z, Liu ZQ, Hou YQ (2010). Dietary supplementation with L-arginine or N-carbamylglutamate enhances intestinal growth and heat shock protein-70 expression in weanling pigs fed a corn- and soybean meal-based diet. *Amino Acid.* 39: 831-839.
- Yamasaki K, Shibata Y, Imai S, Tani Y, Shibasaki Y, Fukuhara T (1984). Clinical application of prostaglandin E1 (PGE1) upon orthodontic tooth movement. *Am. J. Orthod.* 85: 508-518.
- Yamasaki K (1989). Pharmacological control of tooth movement. In: Norton LA, Burstone CJ (Eds). *The Biology of tooth movement,* CRC press, Boca Raton, Florida, USA. pp. 287-320.
- Yao K, Guan S, Li TJ, Huang RL, Wu GY, Ruan Z, Yin YL (2011). Dietary L-arginine supplementation enhances intestinal development and expression of vascular endothelial growth factor in weanling piglets. *Br. J. Nutr.* 105: 703-709
- Yin YL, Tan BE (2010). Manipulation of dietary nitrogen, amino acids and phosphorus to reduce environmental impact of swine production and enhance animal health. *J. Food Agric. Environ.* 8: 447-462.