

www.ajbrui.net

Afr. J. Biomed. Res. Vol.16 (January 2013); 47 - 53

Full Length Research Paper

Attenuation of *Plasmodium Berghei berghei*-induced Bone Marrow Suppression by Bark Extract and fractions of *Chrysophyllum albidum* on bone marrow response to haemolytic conditions in male albino mice

¹Adewoye E.O, ¹Salami A.T and ³Jagun A.T

¹*Department of Physiology, University of Ibadan, Ibadan.*

³*Department of Veterinary Pathology, University of Ibadan, Ibadan*

ABSTRACT

Chrysophyllum albidum is a tropical plant in Southern Nigeria. Methanol extract of *Chrysophyllum albidum* bark (*MeCaB*) has been reported to have antiplasmodial, haematinic and membrane stabilizing properties. The mechanism by which *MeCaB* exerts these properties is yet to be elucidated. This study therefore seeks to investigate the probable mechanism of its haematinic potential on the bone marrow in two separate anaemic studies. Mice were inoculated with *plasmodium berghei berghei* while others were made anaemic by bleeding out 0.25mls of blood through the retro-orbital plexus. Smears of the femoral bone marrow from each group were prepared on days 3 and 7 of extract/drug treatment for blood precursor cell evaluation (myeloid: erythroid ratio; M:E). Groups parasitized and treated continuously for 3 days with *MeCaB* (*PMeCaB*) had significantly reduced M: E ratio (1.88 ± 0.03) compared with parasitized untreated group (*PUn*) (2.33 ± 0.00). Withdrawal of treatment with *MeCaB*, its fractions (*CFr 1,2,3*) and chloroquine from parasitized animals between days 4 to 7 showed significant increase in M:E ratio of *PUn* (5.13 ± 0.59) and *PCq* (4.11 ± 0.37) compared with *PCFr 1* and *3* (1.21 ± 0.08 and 1.69 ± 0.00 respectively). The *MeCaB* only treated and group bled and treated with *MeCaB* (*BMeCaB*) continuously for 7 days had significantly reduced M:E ratio (1.67 ± 0.00 and 1.35 ± 0.00 respectively) compared with control, bled and treated with haematinic (2.13 ± 0.19 , 2.25 ± 0.21 respectively). This study thus establishes that *Chrysophyllum albidum* exerts haematinic properties on bone marrow cells by stimulating the production of more erythroid series which reverted anaemia induced by *plasmodium berghei berghei* and bleeding out.

Keywords: *Chrysophyllum albidum*, bone marrow, anaemia, Mice.

INTRODUCTION

Anaemia is a haemolytic condition which develops when accelerated removal of erythrocytes is not compensated by production from the bone marrow (Lamikanra *et al*, 2007). The fact that some malaria patients develop

severe anaemia, whereas others retain normal or near normal haemoglobin (Hb) could be explained by the amount of erythrocyte destruction occurring during the period until return of normal bone marrow function (Ekvall, 2003). Suppression of bone marrow activity has been reported in all malaria patients (Kurtzhals *et al*

*Address for correspondence:

Email: elolade@yahoo.com

Tel: +234-802-340-5879

Date Received: November 1, 2012

Date Accepted: December 30, 2012

Abstracted by:

Bioline International, African Journals online (AJOL), Index Copernicus, African Index Medicus (WHO), Excerpta medica (EMBASE), CAB Abstracts, SCOPUS, Global Health Abstracts, Asian Science Index, Index Veterinarius, African Journals online

1997) and in asymptomatic *P. falciparum* infection (Kurtzhals *et al*, 1999). Data on the duration of bone marrow activity suppression after malaria attack are conflicting. Some studies reported hypo-proliferative erythropoiesis and dyserythropoiesis for weeks following treatment (Camacho *et al*, 1998, Phillips *et al*, 1986), while other studies have shown that suppression of bone marrow activity is reversed rapidly after treatment (Abdalla *et al*, 1980, Kurtzhals *et al*, 1997).

Bone marrow is a primary lymphoid tissue and a major haematopoietic organ responsible for the production of about 500 billion blood cells daily in form of Red blood cells, White blood cells, Platelets (Barbara, 2006, Abboud and Lichtman, 2001, Hoffman *et al*, 2000, Cline and Golde, 1979,). It accounts for approximately 3% of the body weight in adult rats (Schermer, 1967), 2% in dogs (Jain, 1986b) and 5% in humans (Picker and Siegelman, 1999).

Chrysophyllum albidum G.Don_Holl.(*Sapotaceae*) also known as African star apple is known for its various ethno-medicinal uses (Amusa *et al*, 2003, Pearson, 1976 and Dalziel,1937). The bark is used as a remedy for yellow fever and malaria while the leaves are used as emollients (Adisa, 2000, Adewusi *et al*, 1997). Eleagnine, an alkaloid isolated from *C. albidum* seed cotyledon has been reported to have antinociceptive, anti-inflammatory and antioxidant activities (Idowu *et al*, 2006). Previous studies on methanol extract of *C.albidum* stem bark have revealed its antiplasmodial (Adewoye *et al*, 2010), antimicrobial (Adewoye *et al*, 2011), haematinic (Adewoye *et al*, 2012) and membrane stabilizing potentials (Adewoye *et al*, 2013).

The mechanism by which *C. albidum* stem bark maintains a normal blood cell count in haemolytic conditions are yet to be elucidated. This study therefore seeks to evaluate the activity of the plant bark extract and its active chromatographic fractions on the bone marrow response in *plasmodium berghei berghei* infection and induced- anaemia in albino male mice.

MATERIALS AND METHODS

Reagents: All reagents are of analytical grade and were obtained from BDH chemicals LTD, Poole England.

Drug: Chloroquine was obtained from Sigma (UK) and was used as a positive control drug to evaluate the *in vivo* efficacy of methanolic extract of *C. albidum*.

Plant Materials: Collection and Identification: The fresh bark of *C. albidum* was collected between the months of November 2011 and April 2012 from its

natural habitat at Igbo Owe cash crop farm at Moniya, Akinyele Local Government Area of Oyo State, South-western Nigeria. The plant was identified and given a voucher number FHI 107514 at the Forestry Research Institute of Nigeria (FRIN), Ibadan, Nigeria. The plant bark were dusted and dried at room temperature for 3 weeks and then grounded to powder using a dry electric mill (Moulineux, UK).

Preparation of extract partition and fraction:

Chrysophyllum albidum bark (1.5kg) was soaked in 2.5 litres of n – hexane for 72 hours and the filtrate was collected as hexane partition. The residue was spread out evenly and allowed to dry for 24 hours. To this residue, 2.5 litres of dichloromethane was added and was well macerated for 72 hours after which the filtrate (dichloromethane partition) was decanted and the residue was again spread out to air dry for 24 hours. Methanol (2.5litres) was added to the dried residue stirred and left for 72 hours, after which the whole mixture was filtered and the filtrate was labelled methanol partition. The mixture was filtered with Whatman’s filter paper (No. 1). All the 3 different filtrates were evaporated to dryness *in-vacuo* and stored at 4°C until use.

The methanol partition (*MeCaB*) was then allowed to pass through a silica gel column chromatography using different solvent mixture of n-hexane, dichloromethane and methanol (in varying polarity gradient) to obtain 3 fractions (*CFr1,2 and 3*).

Parasites: Chloroquine-sensitive *P. berghei berghei* (NK 65) was obtained from the National Institute of Malaria Research (NIMAR) Yaba, Lagos State. The parasite was maintained in mice by serial passage of infected blood to uninfected mice in the animal house. Parasitized red blood cells used for inoculation in the experiment were obtained by cardiac puncture from an infected donor mouse. The blood was diluted to desired parasite density in 0.9% NaCl solution (Kendall McGraw, Laboratories, Inc, USA). Each mouse was inoculated with 1.0×10^6 parasitized red blood cells contained in 0.2mls of the NaCl solution.

In this study, the day of inoculation was defined as day zero D_0 while subsequent days were named D_1 , D_2 , D_3 etc.

Animal Grouping

Animals: One hundred and sixty five (165) Swiss male albino mice weighing between 19 - 22 g were used in this experiment. They were obtained from the Animal House of the College of Medicine, University of Ibadan,

Nigeria and were maintained under standard conditions (12 h light and 12 h dark) with free access to mice chow and clean water.

Extract administration

Drugs and extracts were administered orally using orogastric tube. Two separate studies were conducted: studies A and B

- Study A – investigated the effect of methanol extract of *C.albidum* bark (*MeCaB*) and its 3 fractions (*CFr 1, 2 and 3*) on bone marrow of *Plasmodium berghei berghei* infected mice.
- Study B – investigated the effect of methanol extract of *C.albidum* extract (*MeCaB*) on bone marrow of anaemia induced mice.

Study A – We investigated the effect of *C. albidum* bark extract and its 3 fractions (*CFr 1, 2 and 3*) on the bone marrow of *Plasmodium berghei berghei* infected mice. This study contained six groups of 15 mice each.

- Group 1 animals were infected and untreated, (*PUn*).
- Group 2 mice were infected and treated with 30mg/kg *b.w.* Chloroquine for 3 consecutive days, (*PCq*).
- Group 3 consist of mice infected and treated with 1000mg/kg *b.w* of *C. albidum* for 3 consecutive days, (*MeCaB*).
- Group 4 mice were infected and treated with 250mg/kg *b.w* of fraction 1 for 3 consecutive days, (*PCFr1*).
- Group 5 animals were infected and treated with 250mg/kg/day *b.w* of fraction 2 for 3 consecutive days, (*PCFr2*).
- Group 6 consists of mice infected and treated with 250mg/kg *b.w* of fraction 3 for 3 consecutive days, (*PCFr3*).

Study B – In this study, we investigated the effect of methanol extract of *C. albidum* (*MeCaB*) on bone marrow in anaemia-induced mice. The study contained five groups of 15 animals.

Drugs and extract administrations were given orally. Group 1 consists of *Normal* mice that were not bled and did not receive *C albidum* extract, (control). Group 2 consists of mice not bled but received methanolic extract of *C.albidum* (1000mg/kg *b.w*), (*MeCaB*). Group 3 animals were bled (0.25ml of blood) and administered ferrous sulphate (100mg/kg), (*BHaem*). Group 4 animals were bled (0.25ml of blood) and administered *MeCaB* (1000mg/kg *b.w*), (*BMeCaB*). Group 5 animals were bled (0.25ml of blood) and untreated, (*Bla*).

Morphological studies of Bone Marrow:- The air-dried bone marrow smears on glass slides were fixed

with methanol for 2-5 minutes at room temperature. After air-drying for 20- 30 minutes, the fixed smear was stained with Wright's-Giemsa stain and the morphology of hematopoietic cells was investigated under a light microscope. The bone marrow smear was differentially counted (at least 500 cells per slide) into erythroid, granulocytic, agranulocytic, and megakaryocytic series. The myeloid: erythroid ratio (M:E ratio = total myeloid cells/total erythroid cells) was calculated from the proportion of total myeloid lineage cells and total erythroid lineage cells.

Statistical analysis

Experimental data were analyzed using one way analysis of variance (ANOVA) and multiple range tests to determine significant differences between means. Difference between means were regarded as significant at $p < 0.05$.

RESULTS

Effect of *Chrysophyllum albidum* on myeloid-erythroid ratio in parasitized animals.

The effect of the methanolic extract of *Chrysophyllum albidum* (*MeCaB*) on myeloid-erythroid ratio in parasitized animals by days 3 and 9 is shown in figure 1. The myeloid-erythroid ratio of parasitized animals treated with fractions 1, 2 and 3 (*PCFr1, PCFr 2 and PCFr 3*) significantly increased ($p < 0.05$) by day 3 compared with parasitized untreated (*PUn*) and parasitized chloroquine treated (*PCq*) groups. By day 9 of the experiment, the myeloid –erythroid ratio of *PCFr1* and *PCFr 3* treated groups significantly decreased ($p < 0.05$) compared with *PUn* and *PCq* groups.

Histology of bone marrow smear in parasitized animals treated with *Chrysophyllum albidum* and its' fractions.

Plate 1 shows the histology of the bone marrow of *PUn*, *PCFr1* and *PCq* treated groups. The bone marrow showed more erythroid series than myeloid in *PCFr 1* compared with *PUn* and *PCq* which showed presence of malaria parasites.

Myeloid-Erythroid ratio of animals bled and treated with *Chrysophyllum albidum*.

The effect of methanolic extract of *Chrysophyllum albidum* bark (*MeCaB*) on myeloid-erythroid ratio in bled animals by day 7 is shown in Figure 2. The myeloid-erythroid ratio of *MeCaB* and bled mice treated with haematinic (*BMeCaB*) significantly decreased compared with bled untreated (*Bla*) and bled mice treated with haematinic (*BHaem*).




Figure 1: Myeloid-Erythroid ratio for parasitised groups treated with *C.albidum* extract and its' fractions (PCFr 1, PCFr 2 and PCFr 3)




Figure 2: Myeloid-Erythroid ratio for animals bled but treated with *C.albidum* extract (MeCaB).

Histology of bone marrow smear from animals bled and treated with *Chrysophyllum albidum*.

Plate 2 is the histology of bone marrow smear from normal mice, bled mice treated with haematinic

(BHaem) and bled mice treated with methanolic extract of *Chrysophyllum albidum* bark (BMeCaB). The Bhaem and BMeCaB groups revealed presence of more erythroid series compared with the control group.




Plate 1

Plate 1 showing the bone marrow of parasitized mice untreated and treated with PCFr1 and chloroquine. **A.** Parasitised group treated with fraction 1 of *C. albidum* (*PCFr 1*) bark extract. (Mag x 400 H&E). The erythroid cells(E) series are more than the myeloid cells (M). **B.** Parasitised untreated group, (PUn). (Mag x 1000 H&E). Note the presence of malaria parasite (P). **C.** Parasitized rats treated with chloroquine (Mag x 1000 H&E). Note the presence of malaria parasite (P).




Plate 2

Plate 2 showing bone marrow of normal mice and mice bled but treated with haematinics and *C.albidum* extract. **A** Normal picture of the cells in the bone marrow of a mice (Mag x 400 H&E) **B.** Group bled and treated with haematinic (*Bhaem*). (Mag x 400 H&E).Note the presence of more erythroid series (E and R) compared with myeloid series (M). **C.** Group bled treated with MeCaB (*BMeCaB*). (Mag x 1000), H&E.Note the presence of more erythroid series (E) than myeloid series (M).

DISCUSSION

This present study investigated the effects of methanolic extract of *C. albidum* stem bark on the bone marrow. Maggio-Price *et al*, 1985 reported the presence of granular leukocytes with or without little erythroblasts 1 to 3 days post plasmodium infection in mice. By day 5, immature erythroblasts were seen and by day 10, there was intense erythropoiesis and granulopoiesis in ratio 1:1. The anemic conditions observed in these studies might be the result of depletion of murine marrow BFU-E occurring in response to haemolysis or blood loss anaemia (Adamson *et al*, 1978, Hara and Ogawa 1977 and 1976).

Figures 1 and 2 show the significant difference in myeloid-erythroid ratio of *MeCaB*, *BMeCaB*, *PCFr1* and 3 compared with other groups (*PUn*, *PCq* and *BHaem*) by day 7 of the experiment. The increased granulopoiesis observed in bone marrow of *PCFr 2,3* treated and *PUn* between days 1-3 of treatment is similar to the reports of Maggio Price *et al*, 1985. Earlier studies (Adewoye *et al*, 2010, Jubbs *et al*, 1996) reported leucocytosis in infected mice as an indication of enhanced granulopoiesis and lymphocytosis as cellular and humoral responses respectively to the protozoan infection.

On treatment with *MeCaB* for 3 consecutive days and subsequent withdrawal of treatment between days 4 to 7 led to an increase in erythropoiesis in *PCFr1* and 3

treated groups compared with *PUn* and *PCq* (Figure 1, Plate 1). The ratio 1:1 reported by Maggio Price *et al*, 1985 was however not the same in this study. The ratio shifted from 1:1 to 1:2. after treatment. It is most likely that *PCFr 1* and *3* exhibited prolonged effects on the bone marrow thereby producing more erythroid series than *MeCaB*. This shift in ratio, which is a result of more erythroid series being produced in the bone marrow could be responsible for the resistance to developing anaemia in parasitized and extract treated animals (Adewoye *et al*, 2010) compared to chloroquine treated animals.

Adewoye *et al*, 2012 reported a significant haematinic (increased Red blood cell count, haemoglobin concentration and packed cell volume) and anti – inflammatory potential (a significant decrease in Neutrophil/Lymphocyte ratio) in the groups treated with this plant extract. Similarly, some plants have been reported to possess erythropoietic properties: Udut *et al*, 2005, reported the erythropoiesis stimulating effect of *Scutellaria baicalensis* extract through the activation of erythropoiesis precursor cells in the bone marrow of mice. George A Koffuor *et al*, 2012, reported the haematinic properties of ethanol root bark extract of *Carissa edulis* in phenylhydrazine-induced anaemic Sprague Dawley rats. Aimola *et al*, 2013, also reported the erythropoietic ability of *Terminalia catappa* extract in which it stimulated normal erythroid differentiation in phenylhydrazine-induced anaemic mice.

In this study, increased production of erythroid series observed in the extract treated groups have shown a possible mechanism by which *C. albidum* exerts its haematinic property. It may well be that one or more of the active ingredients (flavonoids, tannins, alkaloids and saponins) earlier reported by Adewoye *et al*, 2011 stimulated erythropoietic process (i.e. hyperplasia) during anaemia. Increase in the erythroid series observed in the extract treated groups corroborate the findings of Adewoye *et al*, 2010 where haematological values remained within normal range even with parasite infection when treated with the extract.

In conclusion, Erythrocyte hyperplasia (production of more erythroid cells) observed in this study, exhibited in the bone marrow of mice treated with methanolic extract of *Chrysophyllum albidum* bark (*MeCaB*), mice bled and treated with methanolic extract of *Chrysophyllum albidum* bark (*BMeCaB*), mice parasitized and treated with *Chrysophyllum albidum* fractions 1 and 3 (*PCFr 1* and *3*) could most likely be the mechanism by which the plant reverted or prevented anaemia here and also in previous studies. These observations confirms the potential of the methanolic

extract of *Chrysophyllum albidum* stem bark as a possible source for drug development

REFERENCES

- Aboud CN, Lichtman MA (2001):** Structure of the marrow and the hematopoietic microenvironment. Pages 29–58 in Williams' Hematology, E. Beutler, MA Lichtman, BS Collier, TJ Kipps, and U Seligsohn, 6th edition. McGraw-Hill, New York.
- Abdalla S, Weatherall DJ, Wickramasinghe SN, Hughes M (1980):** The anaemia of *P. falciparum* malaria. Br J Haematol Vol 46: pp. 171-183.
- Adamson JW, Torok-Stok B, Lin N (1978):** Analysis of erythropoiesis by erythroid colony formation in culture. Blood Cells 4:39.
- Adewoye EO, Salami AT, Taiwo VO (2010):** Antiplasmodial and Toxicological effects of methanolic bark extract of *Chrysophyllum albidum* in albino mice. *Journal of Physiology and Pathophysiology* Vol. 1 (1) pp. 1 – 9.
- Adewoye EO, Salami AT, Lawal TO, Adeniyi BA (2011):** The Antimicrobial and Kill Kinetics of *Chrysophyllum albidum* stem bark extract. *European Journal of Scientific Research* Vol 56 No 3 pp. 434 – 444.
- Adewoye EO, Salami AT, Emikpe BO (2012):** Effect of Methanolic Extract of *Chrysophyllum albidum* Bark on Hematological Indices in Mice with Experimental Hemorrhagic Anaemia. *Afr. J. Biomed. Res* 15: 85 – 91.
- Adewoye EO, Salami AT, Azeez OA, Moody JO (2013):** Effect of *Chrysophyllum albidum* stem bark extract on Erythrocyte Membrane Stability. *European Journal of Scientific Research* Vol 97 No 3 pp. 438 – 447.
- Adewusi HA 1997** The African Star Apple *Chrysophyllum albidum* Indigenous Knowledge from Ibadan, Southwestern Nigeria. In: Proceedings of a National Workshop on the Potentials of the Star Apple in Nigeria (Eds.) Denton OA, DO Ladipo, MA Adetoro, MB Sarumi: 25-33.
- Adisa SA 2000** Vitamin C, Protein and mineral contents of African Apple (*Chrysophyllum albidum*) IN: Proceedings of the 18th annual conference of NIST. (Eds.) Garba SA, Ijagbone IF, Iyagba AO, Iyamu AO, Kilani AS, Ufaruna N: 141-146.
- Aimola IA, Inuwa HM, Nok AJ, Mamman Ai, Habila N, Muhammad A, Agbaji AS, Igoche A, Ogungbemi K, Tighil M and Usman Y (2013):** Erythropoietic and Bone marrow stimulating activity of *Terminalia catappa* Extract: Possible Role of Nitric Oxide Signaling. *International Blood Research & Reviews* 1(1): 1-13.
- Amusa NA, Ashaye OA, Oladapo MO, (2003):** Biodeterioration of the African Star Apple (*Chrysophyllum albidum*) in storage and the effect on its food value. *Afr. J. Biotechnol* 2 : 56 – 57.
- Barbara Y, James SL, Alan S, John WH (2006):** Pages 46 – 59 in Wheeters functional histology, 5th edition. Churchill livingstone. Vol 3. China.
- Camacho LH, Gordeuk VR, Wilairatana P, Pootrakul P, Brittenham GM, Looareesuwan S (1998):** The course of

anaemia after the treatment of acute, *falciparum* malaria. *Ann Trop Med Parasitol* 92: 525-537.

Cline MJ, Golde DW (1979): Cellular interactions in haematopoiesis. *Nature* 277: 5693: 177- 181

De buryn PPH (1981): Structural substrates of bone marrow function *Semin Hematol* 18: 179 – 193.

Dalziel JM (1955): The Useful plants of Tropical West Africa, 3rd ed., London, Grown Agents.

Ekvall H (2003): Malaria and anemia. *Curr Opin Hematol* 10: 108-114.

George Asumeng Kuffour, George Henry Sam, Phyllis Elsie Dadzeasah, Emmanuel Owusu Owiafe and Asantewaa Asante – Gyapong (2012): Erythropoietic effect of ethanol root bark extract of *Carissa edulis* in phenylhydrazine – induced anaemic Sprague dawley rats. *Research Journal of Pharmacology* 6(2): 20 – 24.

Hara H, Ogawa M (1976): Erythropoietic precursors in mice with phenylhydrazine- induced anaemia. *Am J Hematol* 1: 453.

Hara H, Ogawa M (1977): Erythropoietic precursors in mice under erythropoietic stimulation and suppression. *Exp Hematol* 5: 141.

Hoffman R, Benz EJ, Shattil SJ, Furie B, Cohen HJ, Silberstein LE, McGlave P (2000): Hematology Basic Principles and Practice, 3rd edition. Churchill Livingstone, New York.

Idowu TO, Iwalewa EO, Aderogba MA, Akinpelu BA, Ogundaini AO (2006): Antinociceptive, Anti - inflammatory and Antioxidant activities of *Eleagnine*: an alkaloid isolated from seed cotyledon of *C. albidum*. *J. Biol. Sci* Vol 6 No 6 pp. 1029 - 1034.

Jain NC (1986b): The hematopoietic system. In Schalm's Veterinary Hematology, 4th edition, pp. 350—387. Lea and Febiger, Philadelphia, PA.

Jubb KVF, Kennedy PC, Palmer N (1995): Pathology of Domestic Animals 3rd edition, Academic Press Inc. New York.

Kurtzhals JA, Rodrigues O, Addae M, Commey JO, Nkrumah FK, Hviid L (1997): Reversible suppression of bone marrow response to erythropoietin in *Plasmodium falciparum* malaria. *Br J Haematol* 97: 169-174.

Kurtzhals JA, Addae MM, Akanmori BD, Dunyo S, Koram KA, Appawu MA, Nkrumah FK, Hviid L (1999): Anaemia caused by asymptomatic *Plasmodium falciparum* infection in semi-immune African schoolchildren. *Trans R Soc Trop Med Hyg* 93: 623-627.

Lamikanra AA, Brown D, Potocnik A, Casals-Pascual C, Langhorne J, Roberts DJ (2007): Malarial anaemia: of mice and men. *Blood* 110: 18-28.

Maggio-Price L, Brookoff D, Weiss L (1985): The haematopoietic stem cells in bone marrow of mice with *Plasmodium berghei berghei* malaria. *Blood journal* 66: 1080-1085.

Phillips RE, Pasvol G, (1992): Anaemia of *Plasmodium falciparum* malaria. *Baillieres Clin Haematol*, 5: 315-330.

Phillips RE, Looareesuwan S, Warrell DA, Lee SH, Karbwang J, Warrell MJ, White NJ, Swasdichai C, Weatherall DJ (1986): The importance of anaemia in cerebral and uncomplicated *falciparum* malaria: role of complications, dyserythropoiesis and iron sequestration. *Q J Med*, 58: 305-323.

Pearson D (1976): Page 54 in The Chemical Analysis of foods, 7th eds Churchill Livingstone New York

Schermer S (1967): Pages 43–49 in The Blood Morphology of Laboratory Animals, 3rd edition. Davis, Philadelphia, PA.

Udud EV, Zhdanov VV, Gur'iantseva LA, Minakova MI, Dygai AM (2005): Mechanisms of the erythropoiesis-stimulating effect of skull cap (*Scutellaria baicalensis*) extract. *Eksperimental'naia i Klinicheskaia Farmakologiya* 68: 43-45