

INVITED SHORT COMMUNICATION

This short communication was invited by the Editor-in-Chief of AJFAND after she read the author's contribution on FARA network on a matter that is becoming increasingly critical to the African farmer, in the wake of evolving climate change impacts on agricultural productivity and continued rising fertilizer prices which hurt smallholder famers.

There is also renewed acknowledgement that soils need biomass and why not source it from the so much waste that goes unutilised!

We would like to thank Prof Sabiiti's acceptance to prepare this piece and express our hope that it receives wide readership.

His contacts are: esabiiti@agric.mak.ac.ug

UTILISING AGRICULTURAL WASTE TO ENHANCE FOOD SECURITY AND CONSERVE THE ENVIRONMENT

Sabiiti EN¹*



Elly Sabiiti

*Corresponding author email: esabiiti@agric.mak.ac.ug

¹Department of Agricultural Production, School of Agricultural Sciences, College of Agricultural and Environmental Sciences, Makerere University, Kampala, Uganda.

ABSTRACT

Due to the increase in the world's population and most of it moving to urban cities, there is increased demand for food, and this has resulted in the production of large amounts of agricultural wastes, both at farmer, municipality and city levels. The bulk of the agricultural food in developing countries is transported to cities in its raw form, thus compounding the net effect on large deposits of waste in urban markets, around homes and in slums as well as in various dumping grounds. In Kampala alone, over 1000mt of waste accumulate in the city and only about 30% of it is collected by City Council leaving the rest to rot and pollute the environment. Although it is recognized that the accumulation of waste has enormous ill effects on humans and the environment, such wastes if properly managed could be considered a big bio-resource for enhancing food security in the smallholder farming communities that would not afford use of expensive inorganic fertilizers. These organic wastes contain high levels of Nitrogen, Phosphorus, Potassium and organic matter important for improving nutrient status of soils in urban agriculture. Various factors amplify the agricultural waste problem, especially in developing countries where there are limited facilities for recycling waste. A lot of the nutrients are leached from the damp fills and end up polluting rivers and water bodies, which have been associated with invasion of water weeds. Most importantly, there is lack of planning, weak public awareness, weak government policy and laws, and lack of or insufficient utilization of resources. In Kampala many small holder farmers have improved milk production by feeding various combination of agricultural wastes and others have increased nutrient supply in soils by applying organic compost leading to improved crop yields, especially vegetables, maize which fetch high prices for the farmers thus reducing poverty levels and enhancing food security of these farmers. This alternate method of removal of these wastes for agricultural production by farmers has also reduced the rate of accumulation with subsequent reduction on environmental pollution thus improving on environmental health. This paper briefly reviews how agricultural wastes can be used to enhance food security and conserve the environment.

Key Words: Agricultural waste, food security, environment

BACKGROUND

Food security (access for all, to sufficient and nutritious food and at all times) encompasses both food production and the ability to access or purchase food [7]. According to Sanchez et. al.[17], it is in Africa where the per capita food production continues to decrease, yet the population is growing at a rate higher than in any other region in the world. Also food insecurity is, most acute in Africa. If the attainment of food security is intrinsically linked with safeguarding the natural resource base, Africa has the highest rate of land degradation in the world [7]. Therefore, since the bulk of food in Africa is produced on a small-scale basis, to ensure food security, the following should be regarded as interlinked determinants: (i) an enabling environment for the smallholder farming sector, (ii) income generation, (iii) reversing soil fertility depletion, (iv) protecting the environment, and (v) intensifying and diversifying land use with high-value products. The importance of agricultural waste as an alternative resource base in food production systems has to be emphasised.

AGRICULTURAL WASTE

The by-products of agricultural activities are usually referred to as “agricultural waste” because they are not the primary products. These wastes chiefly take the form of crop residues (residual stalks, straw, leaves, roots, husks, shells etcetera) and animal waste (manure). Agricultural wastes are widely available, renewable and virtually free, hence they can be an important resource [15]. They can be converted into heat, steam, charcoal, methanol, ethanol, bio diesel as well as raw materials (animal feed, composting, energy and biogas construction and so on)(Fig 1.). However, many of the agricultural wastes are still largely underutilized, and left to rot or openly burned in the field, especially in developing countries. In the city of Kampala, Uganda, over 1000mt of organic waste accumulates daily and only about 30% of this is removed and dumped into a dumpfill in Kitezi .All these wastes are known to contain high nutrient levels of Nitrogen, Potassium, Phosphorus that would improve soil fertility and increase crop yields such as vegetables, maize that fetch high prices and hence enhance food security. This alternate method of utilisation by farmers for agricultural production has also reduced the rate of accumulation, with subsequent reduction on environmental pollution thus improving environmental health. This calls for a greater awareness of the public and farmers of the benefits of proper management and utilization of organic wastes in agriculture. This will lead to diminished fears and preconceived notions of nuisance problems that decrease land values and environmental degradation [20]. The goal should be to make the agricultural waste a resource that can be utilized and not just discarded. It is also very important to set up institutions that can harness the large potential of agricultural wastes as a resource in farming and in energy production. Using appropriate conversion technologies, animal and crop wastes can be turned into useful resources as below:




Figure 1: Conversion of Agricultural Wastes into Various Economic Resources

AGRICULTURAL WASTE AND THE ENVIRONMENT

The impact of agricultural waste on the environment depends not only on the amounts generated but also on the disposal methods used (Fig 2). Some of the disposal practices pollute the environment [16, 19]. For example, agricultural waste burning is a common practice in the undeveloped countries, but it is a source of atmospheric pollution. According to Ezcurra et al. [4], agricultural waste burning releases pollutants such as carbon monoxide, nitrous oxide, nitrogen dioxide and particles (smoke carbon). These pollutants are accompanied by the formation of ozone and nitric acid [6], hence contributing to acid deposition [10] thereby posing risk to human and ecological health.

Environmental pollution from animal waste (faeces, urine, and respiration and fermentation gases) is a global concern and is much more acute and serious in countries with high concentrations of animals on a limited land base for manure disposal. Animal wastes are excreted in solid, liquid, and gaseous forms. Respiration and fermentation gases are lost to the environment soon after being produced by the animal. After excretion, solid and liquid animal waste is subjected to microbial conversion (mainly anaerobic), which converts organic substrates into microbial biomass and soluble and gaseous products. Some of these products have an impact on the environment, as well as effects on water quality, soil deterioration, and air pollution. Odour pollution was reported to contribute highly to social tensions among urban livestock farmers in Kampala, Uganda [9].


Additionally, the application of excessive animal wastes on land as fertilizer and soil conditioner is subject to surface run-off and leaching that may contaminate ground or surface waters. For that reason, nitrate leaching is considered a major nitrogen (N) pollution concern on livestock farms [13]. When phosphorus (P) enters the surface waters from land application of excessive animal manure it can stimulate the growth of algae and other aquatic plants. Their subsequent decomposition results in increased oxygen demand that interferes with the welfare of fish. Manure decomposition can be a major source of methane (CH₄), ammonia (NH₃) and nitrogen oxides, which contribute to accumulation of greenhouse gases. Volatilization of ammonia causes acid deposition, which contributes to acid precipitation [11, 12]. Emissions of nitrous oxide (N₂O) during the nitrification-denitrification cycle contribute to ozone depletion [18].



Heaps of various agricultural wastes



Mode of transporting waste



Environmental pollution

Figure 2: Typical garbage situation in Kampala City and the surrounding environment

AGRICULTURAL WASTE AND FOOD SECURITY

The target should be to make agricultural wastes a resource that can be utilized and not just discarded. Agricultural wastes can be used to enhance food security mainly through their use as bio-fertilizer and soil amendment, use as animal feed, and energy

production. They contain large amounts of organic matter, and many of them can be directly added to the soil without any risk.

Turning these agricultural wastes (crop residues and animal manures) into organic fertilizers (through composting) is one of the waste treatment technologies that make it possible to use organic waste as a fertilizer even in populated areas. Technology plays a key role in soil fertility improvement, and hence crop productivity [1, 5]. The use of organic fertilizers is particularly important in most parts of Africa, where low availability of nutrients is a serious constraint for food production [2]. Composting also reduces the volume of the waste, hence solving serious environmental problems concerning disposal of large quantities of waste, kills pathogens that may be present, decreases the germination of weeds in agricultural fields, and reduces offensive odour [8]. The compost can be sold for additional revenue or used on the same farm. Besides, the production of composts for agricultural use is gaining popularity as a result of the rising interest in organic products such as goat and maize production (Fig. 3).



Figure 3: The wastes are eaten by goats for production of meat while compost applied on soil increases maize production.

Both crop residues and animal waste can be used as animal feed. However, the nutrient content of animal waste depends on the animal species, type of feed, and bedding material used [13]. The use of broiler litter in cattle feeding is a widely applied practice. It is worth noting that animals, especially ruminants are useful in converting crop residues into food, hence contributing substantially to reducing potential pollutants. The rumen contains the microbial enzyme cellulase, which is the only enzyme to digest the most abundant plant product, cellulose [3]. With ruminants, nutrients in by-products are utilized and do not become a waste-disposal problem [14].

The production of energy from agricultural waste has been utilized to varying degrees in different parts of the world [20]. According to Mackie et al.[13], besides generating revenue from the energy produced, waste-to-energy schemes offer an alternative and environmentally acceptable means of waste disposal. Additionally, the schemes also provide a valuable by-product: a good quality, agricultural fertilizer that is nearly odourless. With the concern over future energy shortages and increasing costs of conventional fuels and electricity derived from them, there is increasing interest in using anaerobic digestion as a source of renewable energy while providing acceptable

waste management. For instance, in Uganda, the basic source of fuel is wood used in the form of charcoal or firewood for heating and cooking. This dependence on traditional charcoal and firewood is responsible for the prevailing deforestation and soil degradation, the effects of which have manifested in irregular rainfall, floods and violent storms.

CONCLUSION

Agricultural wastes can be a valuable resource for improving food security; however, if not treated, kept or disposed of properly, agricultural wastes are likely to cause pollution to the environment or even harm to human health. This calls for increased public awareness on the benefits and potential hazards of agricultural wastes, especially in developing countries. There is also an urgent need to set up Centres of excellence in waste management in Africa; an initiative already underway at Makerere University.

ACKNOWLEDGMENTS

I wish to acknowledge the funding from Sida/SAREC Sweden that has enabled Makerere University since 2000 to train several PhD staff specializing in agricultural waste management. Some of the data has come from the research conducted over the years.

REFERENCES

1. **Amoding A** Supply potential and agronomic value of urban market crop wastes. PhD Thesis, Makerere University, 2007.
2. **Brouwer J and JM Powell** Increasing nutrient use efficiency in West African agriculture: The impact of micro-topography on nutrient leaching from cattle and sheep manure. *Agriculture, Ecosystems and Environment* .1998; **71**: 229 - 239.
3. **CAST** Ruminants as food producers: Now and for the future. Council for Agricultural Science and Technology. Special Publication. 1975; **4**: **1** - 13.
4. **Ezcurra AI, Ortiz de Zarate, Pham Vhan Dhin and JP Lacaux** Cereal waste burning pollution observed in the town of Vitoria (northern Spain). *Atmospheric Environment*. 2001; **35**: 1377 - 1386.
5. **Hargreaves JC, Adl MS and P R Warman** A review of the use of composted municipal solid waste in agriculture. *Agriculture, Ecosystems & Environment*. 2008: **123(1-3)**: 1 – 14.
6. **Hegg DA, Radke LF, Hobbs PV, Brock CA and P J Riggan** Nitrogen and Sulphur emissions from the burning of forest products near large urban areas. *J. of Geophysical Research*. 1987; **92**: 14701 - 14709.
7. **IFPRI**. Feeding the world, preventing poverty and protecting the Earth: a 2020 vision.: International Food Policy Research Institute, Washington DC. 1996.
8. **Jakobsen S** Aerobic decomposition of organic wastes 2. Value of compost as fertilizer. *Resources, Conservation and Recycling*. 1995; **13**: 57 - 71.
9. **Katongole CB** Developing rations for meat goats based on some urban market crop wastes PhD. Thesis, Makerere University, Uganda. 2009.
10. **Lacaux JP, Loemba-Ndembi J, Lefeivre B, Cros B and R Delmas** Biogenic emissions and biomass burning influences on the chemistry of the fogwater and stratiform precipitations in the African equatorial forest. *Atmospheric Environment*. 1992; **26(A/4)**: 541 - 551.
11. **Likens GE, Driscoll CT and DC Buso** Long-term effects of acid rain: Response and recovery of a forest ecosystem. *Science (Washington DC)* 272: 244 - 245.
12. **Lowe PD** Social issues and animal wastes: A European perspective. In: *Proceedings of International Livestock Odor Conference*. 1995; pp 168 - 171. Iowa State University College of Agriculture, America.

13. **Mackie RI, Stroot PG and VH Varel** Biochemical identification and biological origin of key odour components in livestock waste. *J. of Anim. Sci.* 1998; **76**: 1331 – 1342.
14. **Oltjen JW and JL Beckett** Role of ruminant livestock in sustainable agricultural systems. *J. Anim. Sci.* 1996; **74**:1406 - 1409.
15. **Sabiiti EN, Bareeba F, SpordlyE, Tenywa, JS , Ledin S, Ottabong E, Kyamanywa S, Ekbom B, Mugisha J and L Drake** Urban market garbage. A resource for sustainable crop/livestock production system and the environment in Uganda. A paper presented at the international conference. *Wastes-The Social Context*. Edmonton, Canada; 2005.
16. **Sabiiti EN, Bareeba F, Spordly E, Tenywa JS, Ledin S, Ottabong E, Kyamanywa S, Ekbom B, Mugisha J and L Drake** Urban market garbage: A hidden resource for sustainable urban/peri-urban agriculture and the environment in Uganda. *The Uganda J.* 2004; **50**: 102-109.
17. **Sanchez PA, Buresh RJ and RR Leakey** Trees, soils, and food security. *Philosophical Transactions of the Royal Society of London.* 1997; **352**: 949 - 961.
18. **Schulte DD** Critical parameters for emissions. **In:** JAM Voermans and GJ Monteny (eds.). *Proceedings. Ammonia and Odour Emissions from Animal Production Facilities*. Pp NVTL Publishing, Rosmalen, The Netherlands. 1997; pp 23-34.
19. **Tumuhairwe JB, Tenywa JS, Otabong E and S Ledin** Comparison of four low-technology composting methods for market crop wastes. *Waste Management.* 2009; **29**: 2274 – 2281.
20. **Westerman PW and JR Bicudo** Management considerations for organic waste use in agriculture. *Bioresource Technology.* 2005; **96**: 215 – 221.