

ORIGINAL RESEARCH ARTICLE

Community Factors Influencing Birth Spacing among Married Women in Uganda and Zimbabwe

¹* Courtney McGuire and ²Rob Stephenson

¹Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA; ²Health Behavior and Biological Sciences, School of Nursing, University of Michigan, Ann Arbor, MI.

*For Correspondence: E-mail: rbsteph@umich.edu: 001-734-615-0149

Abstract

Short birth spacing continues to be a problem in Uganda and Zimbabwe, resulting in negative infant, child, and maternal health outcomes. This study investigates community-level influences on birth spacing outcomes among women aged 15-49 in Uganda and Zimbabwe, using Demographic and Health Surveys conducted in 2011 (Uganda) and 2010-2011 (Zimbabwe). Women living in communities with higher mean maternal age, mean age at marriage, and mean parity were significantly more likely to have longer birth spacing. Women living in communities with higher levels of contraceptive use and low levels of unmet contraceptive need were more likely to have short birth spacing. The significance of community-level demographic and fertility norms, gender norms, economic prosperity, and family planning behaviors demonstrate the broad influence of community variables on birth spacing outcomes. This analysis highlights the importance of moving beyond individual and household-level interventions in order to harness the power of contextual influences on birth spacing. (*Afr J Reprod Health* 2015; 19[1]: 14-24).

Keywords: birth spacing, community, gender, social epidemiology

Résumé

L'espacement des naissances court continue d'être un problème en Ouganda et au Zimbabwe, ce qui entraîne des conséquences négatives chez le nourrisson, l'enfant et la santé maternelle. Cette étude examine les influences au niveau communautaire sur les résultats d'espacement des naissances chez les femmes de 15-49 ans en Ouganda et au Zimbabwe, en utilisant les enquêtes démographiques et de santé menées en 2011 (Ouganda) et 2010-2011 (Zimbabwe). Les femmes vivant dans les communautés ayant un âge maternel moyen plus élevé, l'âge moyen au moment du mariage, la parité moyenne et étaient significativement plus susceptibles d'avoir de plus long espacement des naissances. Les femmes vivant dans les communautés avec des niveaux plus élevés d'utilisation de la contraception et de faibles niveaux de besoins non satisfaits en matière de contraception étaient plus susceptibles d'avoir des naissances rapprochées. L'importance des normes démographique et de fécondité au niveau communautaire, des normes des sexes, la prospérité économique, et les comportements de planification familiale indiquent la grande influence des variables de la communauté sur les résultats d'espacement des naissances. Cette analyse met en évidence l'importance d'aller au-delà des interventions individuelles et au niveau des ménages dans le but d'exploiter la puissance des influences contextuelles sur l'espacement des naissances. (*Afr J Reprod Health* 2015; 19[1]: 14-24).

Mots-clés: espacement des naissances, communauté, sexe, épidémiologie sociale

Introduction

Increasing the proportion of women who are able to adequately space their births is an integral part of reducing under-five mortality rates¹⁻⁶. A 2002 global study of 456,889 pregnancies from low and middle income countries found that the neonatal death rate was 102% higher among children with birth intervals of 9-14 months and 27% higher among children with 15-20 months birth intervals

compared to children born 27-32 months after the previous birth⁷. Short birth intervals (<24 months) have been linked to health effects including low birth weight, pre-term birth, small-for-gestational age³⁻⁸, and childhood stunting⁹, and maternal morbidities, including uterine rupture and utero-placental bleeding disorders¹⁰. A 2011 study of Demographic and Health Surveys (DHS) from 72 countries representing 371,768 birth intervals, showed that an average of only 31% of births

occur in the lowest risk birth interval of 36-59 months¹¹. More than half (57%) of all births occur within an interval of less than 36 months with an additional 12% of births occurring more than 60 months¹¹.

There is increasing recognition of the importance of contextual influences on reproductive health outcomes¹²⁻¹⁹. Studies have shown residence in a community with higher levels of education is associated with higher likelihood of premarital sex¹² but a lower likelihood of risky sexual behaviors¹⁷. Community higher levels of education are also associated with lower demand for large families¹³, lower birth rates¹⁶, and greater contraceptive use¹⁸. This is perhaps due to increased access to reproductive health information in more educated communities, or the creation of alternative pathways to achieving social capital through education and employment that override the desire and expectation for large families. Studies have demonstrated that women who live in wealthier communities are more likely to use modern contraceptives¹⁸, have lowered demand for large families¹³, and are less likely to engage in premarital sex¹². While there are many studies of contextual influences on reproductive health outcomes, there are very few that have specifically examined community-level influences on birth intervals. Hung, et al analyzed DHS data from 11 sub-Saharan countries and found that community prevalence of intimate partner violence and sexual violence had a significant association with shorter birth intervals²¹. Women living in communities with high levels of intimate partner violence (IPV) have experienced pervasive “everyday violence” that contributes to acceptance of unequal power dynamics. This may contribute to an erosion of women’s self-efficacy and ability to control reproductive health outcomes. The limited number of studies on community-influences on birth intervals represents an important gap in the existing literature on birth spacing behaviours. This paper examines the influence of community-level factors on birth spacing behaviour in Uganda and Zimbabwe, to inform the ways through which interventions aimed at improving birth spacing can best target women within community environments.

Methods

Data for this analysis was taken from the most recent Demographic and Health Survey (DHS) for Uganda (2011) and Zimbabwe (2010-2011), available publicly from www.measuredhs.com. Data was downloaded in STATA format. In a 2011 analysis of the most recent DHS data for 70 countries, Zimbabwe was among the countries with the largest proportion of women (42.9%) reporting a birth space of 36-59 months between their most recent two births. Conversely, only 24.6% of women in Uganda reported a birth space of 36-59 months between their last two births¹⁰. Uganda and Zimbabwe were thus selected to represent two very different contexts of birth spacing.

The DHS collect data from women 15-49 and men 15-54 years. DHS employ a two-stage sampling strategy, with households randomly selected within Primary Sampling Units (PSUs). Primary Sampling Units typically comprise 20-30 households. For Uganda, a representative sample of 404 PSU, 10,086 households and 9,247 women aged 15-49 was surveyed. For Zimbabwe, a representative sample of 406 PSU, 10,828 households and 9,831 women aged 15-49 was surveyed. The original samples were 9,171 women in Zimbabwe and 8,674 women in Uganda. To identify community-level factors that influenced birth spacing, the samples were restricted to married women or women reported being in a union with a man, regardless of cohabitation status (to allow the inclusion of partner characteristics). This excluded 3,593 women in Zimbabwe and 3,183 women in Uganda. A further 1,586 women in Zimbabwe and 1,076 women in Uganda were excluded because they did not have two or more children. The final sample included 3,992 women from Zimbabwe and 4,415 women from Uganda.

DHS do not collect community-level data, and we approximate community-level factors by aggregating individual level data to the PSU level. This method has been used extensively in the analysis of community effects on sexual and reproductive health outcomes and behaviors¹²⁻¹⁴. We examine the following domains of community environments as potential influences on birth spacing behaviour.

Community Demographics and Fertility Norms

Previous studies have shown that living in communities with low mean age at marriage and low mean age at first childbirth may reduce a woman's use of contraception¹⁸ and uptake of maternal health services²². A young age at marriage or child birth may suggest that women's progress to womanhood is marked by expectations of early marriage and childbearing, and that there are fewer alternative opportunities – for example education or employment – available to women¹⁸. Similarly, community demographic profiles indicative of fewer opportunities for women may also lead women towards short birth spacing by reducing their ability to seek care and access to family planning services or by encouraging women to high fertility as a means of achieving social expectations. To measure demographic context we use the community mean age at marriage, the mean age at first sex, the mean age at childbearing, and the mean ideal number of children.

Community Economic Prosperity

Previous studies have demonstrated that higher socioeconomic status is associated with delayed onset of first marriage, sex, and childbirth²⁴ and with increased use of modern contraceptives²³. Residing in a wealthier community may result in greater access to health knowledge, family planning and maternal health care services. We measure community economic prosperity with the community mean wealth index factor score, a household level indicator of the ownership of household goods.

Community Gender Norms

Previous studies have shown that living in communities with higher education influence a woman's use of modern birth control^{12, 13, 18} and have lower levels of fertility¹⁶. Higher community levels of female employment and education may contribute to greater autonomy and its positive effect on reproductive health outcomes²⁵. These may be communities in which the rights to education and employment for women are viewed as the same for men, and therefore have more progressive gender norms that more freely allow

women to make reproductive health decisions. To measure community gender norms, we used ratio of women to men in the community with at least primary school education, the percentage of women in each community who were employed in the previous 12 months, and the community mean violence justification score. The violence justification index included questions of whether women felt a husband was justified in beating his wife in five situations, such as if she burns the food or neglects the children. Women who scored 5 on the index believed that husbands were justified in beating their wives in each scenario.

Community Family Planning Behaviour

Previous studies have shown that women who live in communities with family planning messaging, a marker of a successful reproductive health program, are four times as likely to be using a modern contraceptive method than those who are not exposed to family planning messages¹⁵. To measure community family planning behaviours, we use the percentage of women in the community with unmet need for spacing or limiting of births, the mean ideal birth space reported by women in the community, and the percentage of women in the community who are using modern contraceptive methods.

Individual and Household Measures

In addition to the community level variables, the analysis controlled for individual and household factors shown by previous studies to be associated with birth spacing behaviour. For individual variables we controlled for maternal age (15-24, 25-29, 30-34, 35-39, 40-49), parity (2, 3-4, 5-6, 7+ children), age at marriage (less than 14, 15-19, 20-39), and spousal age difference (20 years younger to 5 years older, 6 years older and above). We also controlled for current use of birth control (modern, no method/folkloric/traditional), women's and husband education (no education/primary, secondary/higher education). Individual variables relating to the index child included the sex of the index child and if it was alive. At the household level we controlled for household wealth, measured by the wealth quintile.

Analysis was conducted in STATA 12²⁶. Data were first cleaned, including checks for missing data and re-categorization of variables to create either categorical variables (transfer of continuous age at marriage into categories) or to reduce the number of categories (e.g. combining secondary and higher education into one category). The analysis considers two outcomes. The first is a continuous variable measuring the self-reported length of time in months between the most recent birth (index birth) and the previous birth. The second outcome is a categorical variable measuring the number of months between the index birth and the previous birth, categorized as: <24 months, 25-38 months, 39-59 months, and >60 months). The reference category was taken as 25-38 months. The second stage of analysis involved bivariate analysis to test for significant associations between the measures of birth spacing and the individual, household and community variables. T-tests for variations in means were conducted to examine variations in birth spacing length across the variables. The third stage of analysis involved multivariate models. Two regression models were fitted. A linear regression model was fitted to the first outcome, the continuous measure of self-reported birth spacing.

An ordered logistic model was fitted for the second outcome, the categorical variable measuring the length of the previous birth space. The same covariates were included in each model. Each model also controlled for the sampling strategy used in the collected of the DHS data (the survey strata, clustering and weighting). Interaction terms were tested between key covariates, but no significant interactions were found.

Results

Community characteristics were only significantly associated with the length of the preceding birth interval in Uganda (Table 1). In Uganda, living in a community with a higher mean age at first birth was significantly negatively associated with the length of the preceding birth interval (beta -1.26, SE 0.39, p=0.001). Living in a community with a higher mean age at first cohabitation in years was significantly positively associated with the length of the preceding birth interval (beta 1.13, SE 0.36, p=0.002). Community wealth index score was associated with birth intervals in Uganda (beta 0.00, SE 0.00, p=0.004).

Table 1: Linear Regression for Length of Preceding Birth Interval in Months

Figures shown are beta coefficients, standard error and associated P-Value.

	Zimbabwe		Uganda	
	Beta coefficient (SE)	P value	Beta coefficient (SE)	P value
Age				
15-24 (ref)				
25-29	13.69 (2.02)*	0.000	8.21 (1.07)*	0.000
30-34	25.51 (2.20)*	0.000	16.47 (1.29)*	0.000
35-39	39.35 (2.43)*	0.000	22.42 (1.41)*	0.000
40-49	40.06 (2.51)*	0.000	26.16 (1.45)*	0.000
Women's Education				
No education/Primary (ref)				
Secondary/Higher	-2.76 (1.38)*	0.046	1.51 (0.99)	0.128
Husband's Education				
No education/Primary (ref)				
Secondary/Higher	-1.98 (1.43)	0.166	0.69 (0.64)	0.377
Age at Marriage				
5-14 (ref)				
15-19	-0.66 (2.24)	0.768	-4.32 (0.87)*	0.000
20-39	-5.35 (2.41)*	0.026	-7.55 (1.12)*	0.000
Spousal Age Difference				
20 years younger to 5 years older (ref)				
6 years older and above	-0.37 (1.17)	0.753	0.19 (0.64)	0.770
Parity				
2 children (ref)				

3-4 children	-7.11 (1.48)*	0.000	-5.23 (1.01)*	0.000
5-6 children	-20.00 (2.15)*	0.000	-10.44 (1.25)*	0.000
7+ children	-30.33 (2.89)*	0.000	-19.07 (1.40)*	0.000
Current use of birth control by method type				
No method/Folkloric/Traditional (ref)				
Modern Method	6.63 (1.29)*	0.000	-0.71 (0.80)	0.376
Sex of First Child				
Male (ref)				
Female	0.61 (1.15)	0.593	0.87 (0.63)	0.171
Alive first child				
Yes (ref)				
No	-12.67 (2.18)*	0.000	-5.04 (1.11)*	0.000
Wealth Quintiles				
Poorest (ref)				
Poorer	5.61 (1.85)*	0.003	-0.15 (1.03)	0.884
Middle	3.59 (1.93)	0.063	-1.07 (1.11)	0.336
Richer	4.75 (1.98)*	0.017	0.67 (1.19)	0.574
Richest	4.52 (2.18)*	0.039	-0.12 (1.58)	0.941
Community Level Variables				
Wealth Index	0.00 (0.00)	0.673	0.00 (0.00)*	0.004
Violence Justification Index	-0.81 (1.15)	0.480	-0.87 (0.45)	0.052
Women working	-0.88 (2.65)	0.741	-1.36 (1.80)	0.450
Ideal birth spacing	-0.34 (0.04)	0.378	-0.05 (0.03)	0.073
Ideal number of children	0.22 (0.75)	0.770	-0.19 (0.38)	0.612
Age at first birth	0.44 (0.67)	0.511	-1.26 (0.39)*	0.001
Age at first sex	-0.33 (0.33)	0.314	-0.04 (0.31)	0.893
Age at first cohabitation	-0.42 (0.61)	0.485	1.13 (0.36)*	0.002
Unmet need	3.32 (6.02)	0.581	3.99 (2.29)	0.082
Birth control	3.43 (4.35)	0.431	-0.74 (2.55)	0.772
Ratio of education (women to men)	0.26 (6.55)	0.968	2.40 (1.45)	0.097

In Uganda only, women residing in wealthier communities were more likely to report all birth intervals relative to birth intervals of 25-38 months (Table 2). Zimbabwean women who resided in a community with greater justification for violence against women were more likely to report birth intervals of 39-59 months than 25-38 months (Relative Risk Ratio (RRR) 1.40, 95% CI 1.10-1.78, $p=0.007$). Women living in communities in Uganda in which women reported wanting longer birth spaces, measured in months, were more likely to report birth intervals of 39-59 months than 25-38 months (RRR 0.99, 95% CI 0.98-1.00, $p=0.016$). Women living in communities in Zimbabwe in which there was a reported higher mean age at first birth in years were significantly more likely to report birth intervals of less than 24 months than 25-38 months (RRR 1.24, 95% CI 1.01-1.51, $p=0.037$). Ugandan women living in communities in which there was a higher reported mean age at first birth in years were significantly less likely to report a birth of greater than 60 months relative to 25-38 months (RRR 0.80, 95%

CI 0.70-0.91, $p=0.001$). Ugandan women in communities with lower levels of reported unmet need were significantly less likely to report birth intervals of less than 24 months than 25-38 months (RRR 0.32, 95% CI 0.18-0.55, $p=0.000$). Zimbabwean women residing in communities with higher levels of reported contraceptive use were significantly less likely to report birth intervals of less than 24 months than 25-38 months (RRR 0.28, 95% CI 0.08-0.94, $p=0.039$). Community-level characteristics of education, age at first cohabitation and sex, women working, and ideal number of children were not significantly associated with birth intervals in Uganda or Zimbabwe.

Individual and household factors found to be significantly associated with birth spacing behavior were generally in line with those identified in previous studies. Age had a significant positive association with the length of the preceding birth interval in both countries and among all ages. Relative to women who married at less than 14, older age at marriage (20-39) was

negatively associated with the length of the preceding birth interval (Uganda: beta -7.55, SE 1.12, $p=0.000$; Zimbabwe: beta -5.35, SE 2.41, $p=0.026$). Parity was significantly associated with the length of the preceding birth interval in both Uganda and Zimbabwe. Relative to women whose first child was alive, women whose index child had died were more likely to have shorter birth spaces. There was evidence of an association between contraceptive use and birth spacing: In Zimbabwe, compared to women who utilized no method, folkloric or traditional methods, women who utilized modern contraceptive methods were significantly more likely to have a birth interval greater than 60 months than a birth interval of 25-38 months (RRR2.00, 95% CI 1.50-2.66, $p=0.000$). Women's education was significantly associated with the length of the preceding birth interval in Zimbabwe, with women reporting secondary or higher education reporting shorter birth intervals (beta -2.76, SE 1.38, $p=0.046$). Wealth was only significantly associated with the length of the preceding birth interval in Zimbabwe. Relative to the poorest quintile, the middle quintile had a significant positive association with the length of the preceding birth interval.

Discussion

The results point to the different pathways through which the community environment may influence birth spacing behaviour. Interestingly, the analysis found more significant community-level factors associated with birth spacing in Uganda than in Zimbabwe.

Several of the results found here suggest the role of community-level gender norms in shaping birth spacing behavior. Women living in communities with higher age at marriage and higher age at first birth had shorter birth intervals, indicating that demographic patterns in which women marry and begin childbearing later do not necessarily result in healthy child spacing behaviors. Delayed marriage and childbirth may be indicative of more liberal community norms but subsequent short birth intervals are reflective of gender inequities that emphasize the importance of large family sizes. Upadhyay found that women in Cebu, Philippines reported that they had short

birth intervals after delaying childbearing in order to "catch up" with their peers who had longer periods to plan their birth intervals²⁷. This community emphasis on achieving fertility expectations in a shorter amount of time may be the pathway to short birth intervals.

Women in Zimbabwe living in communities with higher violence justification rates were more likely to have optimal birth spacing intervals (Table 2). This surprising result is consistent with Elfstrom's (2012) findings that women in Zimbabwe who reported higher levels of violence justification were more likely to use modern contraceptives¹⁴. Both studies utilized cross-sectional DHS data and thus cannot draw conclusions on causality. This surprising result warrants further research: it is possible that higher levels of violence justification may be discouraging women from childbearing.

The study's results point to the impact of living in a community with limited access to reproductive health services on achieving optimal birth intervals. Women living in communities with greater levels of unmet need for modern contraception and lower rates of contraceptive use are more likely to have shorter birth intervals (Table 2), demonstrating an important synergy in reproductive health outcomes. Kaggwa et al. showed that women living in Malian communities with high levels of exposure to family planning messaging were more likely to adopt modern contraceptives¹⁵. Communities without physical or financial access to reproductive health services are left without a crucial source of reproductive knowledge and care, hence limiting options for family planning and increasing the likelihood of short birth spaces.

There are a number of limitations to the current study. DHS does not collect community-level data. For this analysis, we aggregated individual data to create proxies for community-level variables. Many other studies have utilized similar methodology¹²⁻¹⁹ to establish associations between community level proxies and sexual and reproductive health outcomes. Since the data used were cross-sectional, it is not possible to draw conclusions regarding causality of the associations identified. Another limitation is the absence of data on the presence of reproductive health

services within the community. The results do not consider the availability or quality of sexual and reproductive health services in the community, although we use two proxies (mean unmet need for

contraception and mean contraceptive use) as measures of likely exposure to sexual and reproductive health services.

Table 2: Ordinal Regression Model for Length of Preceding Birth Interval.

Outcome variable is: <24 months, 25-38 months (reference), 39-59 months, and > 60 months.

Figures presented are Relative Risk Ratio of birth interval category relative to 25-38 months, associated 95% Confidence Intervals, and P-values.

	<u>Zimbabwe</u>			<u>Uganda</u>								
	<24 Months RRR (95% CI)	P value	39-59 months RRR (95% CI)	P value	>60 months RRR (95% CI)	P value	<24 Months RRR (95% CI)	P value	39-59 months RRR (95% CI)	P value	>60 months RRR (95% CI)	P value
Age												
15-24 (ref)												
25-29	0.62 (0.37-1.03)	0.066	2.59 (1.75-3.83)*	0.000	4.55 (2.79-7.43)*	0.000	0.65 (0.50-0.83)*	0.001	1.72 (1.26-2.33)*	0.001	3.18 (1.96-5.15)*	0.000
30-34	0.62 (0.33-1.14)	0.125	4.18 (2.64-6.63)*	0.000	13.53 (7.89-23.20)*	0.000	0.72 (0.53-0.99)*	0.045	2.43 (1.69-3.50)*	0.000	9.70 (5.76-16.33)*	0.000
35-39	0.60 (0.29-1.22)	0.156	5.19 (3.06-8.80)*	0.000	29.98 (14.90-48.84)*	0.000	0.65 (0.45-0.92)*	0.016	2.74 (1.84-4.08)*	0.000	19.07 (11.04-32.96)*	0.000
40-49	0.97 (0.47-2.01)	0.945	6.30 (3.59-11.06)*	0.000	37.95 (20.37-70.72)*	0.000	0.86 (0.60-1.25)	0.438	3.66 (2.42-5.53)*	0.000	35.56 (20.37-62.07)*	0.000
Women's Education												
No education/Primary (ref)												
Secondary/Higher	1.07 (0.72-1.59)	0.730	0.95 (0.711-1.27)	0.711	0.83 (0.61-1.12)	0.223	0.90 (0.71-1.15)	0.410	1.02 (0.78-1.34)	0.873	1.11 (0.80-1.53)	0.535
Husband's Education												
No education/Primary (ref)												
Secondary/Higher	1.23 (0.81-1.87)	0.324	1.02 (0.75-1.38)	0.920	0.97 (0.71-1.33)	0.854	0.91 (0.75-1.10)	0.319	0.95 (0.77-1.17)	0.601	1.16 (0.89-1.52)	0.260
Age at Marriage												
5-14 (ref)												
15-19	0.42 (0.23-0.79)*	0.007	0.52 (0.31-0.87)*	0.013	0.58 (0.34-0.99)*	0.046	1.20 (0.96-1.49)	0.103	0.79 (0.63-1.00)*	0.047	0.64 (0.47-0.87)*	0.004
20-39	0.40 (0.20-0.80)*	0.009	0.43 (0.25-0.74)*	0.002	0.39 (0.22-0.69)*	0.001	1.47 (1.11-1.93)*	0.007	0.73 (0.54-0.98)*	0.038	0.53 (0.36-0.77)*	0.001
Spousal Age Difference												
20 years younger to 5 years older (ref)												
6 years older and above	0.74 (0.53-1.05)	0.088	1.06 (0.83-1.36)	0.654	0.97 (0.75-1.26)	0.833	1.04 (0.89-1.21)	0.623	1.07 (0.90-1.28)	0.415	1.13 (0.90-1.41)	0.299
Parity												
2 children (ref)												

<i>McGuire & Stephenson</i>		<i>Community Contexts and Birth Spacing</i>										
3-4 children	1.32 (0.84-2.09)	0.226	0.52 (0.38-0.72)*	0.000	0.51 (0.37-0.72)*	0.000	0.95 (0.74-1.21)	0.659	0.78 (0.59-1.04)	0.087	0.40 (0.28-0.57)*	0.000
5-6 children	1.53 (0.79-2.97)	0.211	0.30 (0.18-0.47)*	0.000	0.21 (0.13-0.33)*	0.000	1.05 (0.76-1.43)	0.775	0.63 (0.45-0.89)*	0.010	0.25 (0.17-0.38)*	0.000
7+ children	1.76 (0.78-3.98)	0.171	0.18 (0.10-0.34)*	0.000	0.08 (0.04-0.16)*	0.000	1.13 (0.79-1.61)*	0.505	0.60 (0.40-0.88)*	0.009	0.08 (0.05-0.13)*	0.000
Current use of birth control by method type												
No												
method/Folkloric/Traditional (ref)												
Modern Method	0.96 (0.67-1.38)	0.824	1.20 (0.92-1.58)	0.180	2.00 (1.50-2.66)*	0.000	1.05 (0.87-1.28)	0.605	0.88 (0.71-1.09)	0.241	1.01 (0.77-1.32)	0.955
Sex of First Child												
Male (ref)												
Female	0.89 (0.64-1.25)	0.501	1.08 (0.84-1.37)	0.550	1.16 (0.90-1.49)	0.240	0.91 (0.78-1.06)	0.240	1.05 (0.88-1.24)	0.590	1.08 (0.86-1.34)	0.504
Alive first child												
Yes (ref)												
No	4.39 (2.80-6.88)*	0.000	0.43 (0.26-0.73)*	0.002	0.44 (0.25-0.75)*	0.003	3.20 (2.45-4.18)*	0.000	1.69 (1.22-2.34)*	0.002	1.34 (0.86-2.08)	0.197
Wealth Quintiles												
Poorest (ref)												
Poorer	0.86 (0.51-1.46)	0.584	1.05 (0.72-1.54)	0.785	1.47 (0.98-2.20)	0.064	9.83 (0.65-1.06)	0.142	0.82 (0.62-1.08)	0.149	1.11 (0.73-1.69)	0.717
Middle	0.92 (0.53-1.60)	0.771	1.20 (0.80-1.79)	0.376	1.37 (0.90-2.10)	0.146	1.07 (0.83-1.40)	0.592	0.87 (0.64-1.17)	0.354	1.18 (0.77-1.82)	0.452
Richer	0.99 (0.56-1.74)	0.965	1.14 (0.76-1.73)	0.510	1.43 (0.93-2.20)	0.107	1.03 (0.77-1.37)	0.861	1.17 (0.85-1.59)	0.335	1.51 (0.97-2.33)	0.065
Richest	1.13 (0.60-2.10)	0.711	1.18 (0.74-1.89)	0.475	1.59 (0.98-2.58)	0.058	0.91 (0.62-1.35)	0.651	1.06 (0.70-1.60)	0.799	0.93 (0.53-1.63)	0.791
Community Level Variables												
Wealth Index	1.00 (1.00-1.00)	0.987	1.00 (1.00-1.00)	0.289	1.00 (1.00-1.00)	0.959	1.00 (1.00-1.00)*	0.007	1.00 (1.00-1.00)*	0.016	1.00 (1.00-1.00)*	0.000
Violence Justification Index	0.91 (0.65-1.28)	0.600	1.40 (1.10-1.78)*	0.007	1.07 (0.83-1.38)	0.622	0.99 (0.89-1.10)	0.865	0.95 (0.84-1.07)	0.364	0.91 (0.78-1.06)	0.232
Women working	1.29 (0.59-2.82)	0.527	0.81 (0.46-1.43)	0.471	1.00 (0.56-1.79)	0.993	0.85 (0.55-1.32)	0.468	0.73 (0.45-1.18)	0.201	0.66 (0.35-1.25)	0.203
Ideal birth spacing	0.99 (0.98-1.00)	0.213	1.01 (1.00-1.02)	0.120	1.00 (0.99-1.01)	0.688	1.00 (0.99-1.01)	0.827	0.99 (0.98-1.00)*	0.016	0.99 (0.99-1.00)	0.213
Ideal number of children	0.91 (0.73-1.14)	0.407	0.86 (0.87-1.16)	0.943	0.91 (0.77-1.07)	0.242	1.02 (0.93-1.12)	9.640	0.95 (0.86-1.05)	0.350	0.97 (0.84-1.12)	0.683
Age at first birth	1.24 (1.01-1.51)*	0.037	1.01 (0.87-1.16)	0.943	.08 (0.93-1.26)	0.301	0.94 (0.85-1.03)	0.181	0.93 (0.83-1.02)	0.160	0.80 (0.70-0.91)*	0.001
Age at first sex	0.96 (0.88-1.05)	0.402	0.98 (0.91-1.04)	0.469	0.94 (0.88-1.01)	0.119	0.97 (0.90-1.05)	0.469	1.01 (0.93-1.09)	0.837	1.02 (0.91-1.13)	0.771
Age at first cohabitation	0.88 (0.83-1.06)	0.165	1.09 (0.96-1.25)	0.174	0.96 (0.84-1.10)	0.604	1.00 (0.92-1.10)	0.969	1.02 (0.93-1.13)	0.682	1.06 (0.94-1.19)	0.381
Unmet need	0.42 (0.09-2.46)	0.368	0.87 (0.24-3.12)	0.831	0.84 (0.23-3.07)	0.787	0.32 (0.18-0.55)*	0.000	0.69 (0.37-1.28)	0.236	0.88 (0.39-2.00)	0.757
Birth control	0.28 (0.08-0.94)*	0.039	1.12 (0.44-2.84)	0.814	0.86 (0.33-2.22)	0.752	0.85 (0.45-1.58)	0.600	0.88 (0.45-1.74)	0.718	0.85 (0.36-2.03)	0.718
Ratio of education (women to men)	2.65 (0.46-15.41)	0.277	-0.65 (0.17-2.51)	0.530	0.82 (0.19-3.50)	0.794	0.78 (0.56-1.10)	0.157	0.98 (9.66-1.46)	0.936	1.65 (0.90-3.00)	0.103

Conclusion

This study is the first of its kind to investigate multiple community-level influences on birth spacing in two different resource-poor settings. The results add to a growing body of literature on the importance of moving beyond individual and household-level variables and exploring contextual influences on reproductive health. The findings highlight the importance of the community demographic profile and related prevailing gender norms and expectations in shaping birth spacing behaviour. The results indicate the need to move beyond traditional individual and household level factors when designing interventions to improve birth spacing behaviour, and to consider the importance of the community environment as a target for intervention.

Contribution of Authors

CM and RS conceptualized and designed the study. CM conducted the data analysis, with supervision from RS. CM and RS both contributed to the writing of the manuscript. CM and RS both approved the final content of the manuscript.

References

- Houle, B.; Stein, A.; Kahn, K.; Madhavan, S.; Collinson, M.; Tollman, S.M.; Clark, S.J.: 2013: Household context and child mortality in rural south africa: The effects of birth spacing, shared mortality, household composition and socio-economic status. *International journal of epidemiology*, 42, 1444-1454.
- Ayiko, R.; Antai, D.; Kulane, A.: 2009: Trends and determinants of under-five mortality in uganda. *East African journal of public health*, 6, 136-140.
- Kozuki, N.; Lee, A.C.; Silveira, M.F.; Victora, C.G.; Adair, L.; Humphrey, J.; Ntozini, R.; Black, R.E.; Katz, J.: 2013: The associations of birth intervals with small-for-gestational-age, preterm, and neonatal and infant mortality: A meta-analysis. *BMC public health*, 13, S3.
- Kozuki, N and Walker, N: 2013: Exploring the association between short/long preceding birth intervals and child mortality: Using reference birth interval children of the same mother as comparison. *BMC public health*, 13 Suppl 3, S6.
- Fotso, J.C.; Cleland, J.; Mberu, B.; Mutua, M.; Elungata, P.: 2013: Birth spacing and child mortality: An analysis of prospective data from the nairobi urban health and demographic surveillance system. *Journal of biosocial science*, 45, 779-798.
- DaVanzo, J.; Hale, L.; Razzaque, A.; Rahman, M.: 2008: The effects of pregnancy spacing on infant and child mortality in matlab, bangladesh: How they vary by the type of pregnancy outcome that began the interval. *Population studies*, 62, 131-154.
- Rustein S: 2002: Effects of birth interval on mortality and health: multivariable cross-country analysis, MACRO International Presentation at USAID, July 2000. In: Setty-Venugopal V, Upadhyay UD: Birth Spacing: Three to Five Saves Lives. Baltimore, Johns Hopkins Bloomberg School of Public Health, Population Information Program, (Population Reports, Series L, Number 13).
- Conde-Agudelo, A.; Rosas-Bermudez, A.; Kafury-Goeta, A.C.: 2006: Birth spacing and risk of adverse perinatal outcomes: A meta-analysis. *JAMA : the journal of the American Medical Association*, 295, 1809-1823.
- Adekanmbi, V.T.; Kayode, G.A.: 2012: Uthman, O.A., Individual and contextual factors associated with childhood stunting in nigeria: A multilevel analysis. *Maternal & child nutrition*, 9, 244-259.
- Conde-Agudelo, A.; Rosas-Bermudez, A.; Kafury-Goeta, A.C.: 2007: Effects of birth spacing on maternal health: A systematic review. *American journal of obstetrics and gynecology*, 196, 297-308.
- Rutstein, S O: 2011. Trends in Birth Spacing. DHS Comparative Reports No. 28. Calverton, Maryland, USA: ICF Macro.
- Setty-Venugopal V, Upadhyay UD: 2002: Birth Spacing: Three to Five Saves Lives. Baltimore, Johns Hopkins Bloomberg School of Public Health, Population Information Program, (Population Reports, Series L, Number 13).
- Chiao, C: 2010: Community context and the prevalence of premarital sex among young women in kenya and the philippines: Trends and differences from 1993 to 2003. *Health & place*, 16, 512-522.
- Chiao, C.; Yi, C.C.; Ksobiech, K.: 2012: Community effects on pregnancy intention among cohabiting women in the philippines: Implications for maternal and child health. *Maternal and child health journal*, 16, 1293-1303.
- Elfstrom, K.M.; Stephenson, R.: 2012: The role of place in shaping contraceptive use among women in africa. *PloS one*, 7, e40670.
- Kaggwa, E.B.; Diop, N.; Storey, J.D.: 2008: The role of individual and community normative factors: A multilevel analysis of contraceptive use among women in union in mali. *International family planning perspectives*, 34, 79-88.
- Kravdal, O: 2002: Education and fertility in sub-saharan africa: Individual and community effects. *Demography*, 39, 233-250.
- Stephenson, R: 2009: Community influences on young people's sexual behavior in 3 african countries. *American journal of public health*, 99, 102-109.
- Stephenson, R.; Baschieri, A.; Clements, S.; Hennink, M.; Madise, N: 2007: Contextual influences on modern

- contraceptive use in sub-saharan africa. *American journal of public health*, 97, 1233-1240.
20. Stephenson, R.; Beke, A.; Tshibangu, D: 2008: Community and health facility influences on contraceptive method choice in the eastern cape, south africa. *International family planning perspectives*, 34, 62-70.
 21. Stephenson, R.; Tsui, A.O: 2002: Contextual influences on reproductive health service use in uttar pradesh, india. *Studies in family planning*, 33, 309-320.
 22. Hung, K.J.; Scott, J.; Ricciotti, H.A.; Johnson, T.R.; Tsai, A.C: 2012: Community-level and individual-level influences of intimate partner violence on birth spacing in sub-saharan africa. *Obstetrics and gynecology*, 119, 975-982.
 23. Nasrullah, M.; Zakar, R.; Kramer, A: 2013: Effect of child marriage on use of maternal health care services in pakistan. *Obstetrics and gynecology*, 122, 517-524.
 24. Chakraborty, N.M.; Firestone, R.; Bellows, N: 2013: Equity monitoring for social marketing: Use of wealth quintiles and the concentration index for decision making in hiv prevention, family planning, and malaria programs. *BMC public health*, 13 Suppl 2, S6.
 25. Mistry, R.; Galal, O.; Lu, M: 2009: Women's autonomy and pregnancy care in rural india: A contextual analysis. *Social science & medicine*, 69, 926-933.
 26. StataCorp: 2013: Stata statistical software: Release 13.
 27. Upadhyay UD and Hindin MJ: 2005: Do higher status and more autonomous women have longer birth intervals? Results from Cebu, Philippines. *Social Science and Medicine*, volume 60, issue 11, 2641-2655