

ORIGINAL RESEARCH ARTICLE

Analysis of maternal serum vitamin D concentrations at birth in women presenting with spontaneous preterm birth: A case-control study

DOI: 10.29063/ajrh2021/v25i2.10

Grace Chiudzu^{1*}, Alfred Maluwa², Augustine Choko³ and Jon Odland^{4,5}

Department of Obstetrics and Gynaecology, University of Malawi, College of Medicine, Lilongwe, Malawi; Kamuzu Central Hospital, Lilongwe, Malawi; Bwaila Hospital, Lilongwe, Malawi¹; Malawi University of Science and Technology, Limbe²; Malawi Liverpool Wellcome Trust Clinical Research Programme, TB/HIV Department, Blantyre, Malawi³. The Norwegian University of Science and Technology, Trondheim, Norway⁴; School of Health Systems and Public Health, University of Pretoria, South Africa⁵

*For Correspondence: Email: m201580013239@medcol.mw; Phone: 265 (0) 888962561

Abstract

Vitamin D is a potent immune system modulator; its deficiency correlates with increased susceptibility to infections. We evaluated the status of maternal serum vitamin D in women with spontaneous preterm birth. In this case-control study, the maternal serum concentration of vitamin D (25OH D) was measured in 95 women delivering preterm and 92 women having a term birth. Vitamin D sufficiency was found in 79% of the mothers who delivered preterm and 80.4% of the mothers who had term birth ($p=0.822$). There was a negative correlation between maternal serum concentration of 25 Hydroxyvitamin D and maternal age in the preterm birth group ($p=0.043$). In conclusion, there was no difference in maternal serum concentrations of 25 Hydroxyvitamin D between women delivering preterm compared to those having term birth. Maternal serum concentration of 25 Hydroxyvitamin D is not associated with occurrence of preterm birth. (*Afr J Reprod Health* 2021; 25[2]: 103-109).

Keywords: Preterm birth, vitamin D, preterm delivery, 25 Hydroxyvitamin D, premature

Résumé

La vitamine D est un puissant modulateur du système immunitaire; sa carence est corrélée à une sensibilité accrue aux infections. Nous avons évalué le statut de la vitamine D sérique maternelle chez les femmes ayant une naissance prématurée spontanée. Dans cette étude cas-témoins, la concentration sérique maternelle de vitamine D (25OH D) a été mesurée chez 95 femmes ayant accouché avant terme et 92 femmes ayant une naissance à terme. Une suffisance en vitamine D a été trouvée chez 79% des mères qui ont accouché prématurément et 80,4% des mères qui ont eu un accouchement à terme ($p = 0,822$). Il y avait une corrélation négative entre la concentration sérique maternelle de 25 hydroxyvitamine D et l'âge maternel dans le groupe des naissances prématurées ($p = 0,043$). En conclusion, il n'y avait aucune différence dans les concentrations sériques maternelles de 25 Hydroxyvitamine D entre les femmes ayant accouché avant terme et celles ayant une naissance à terme. La concentration sérique maternelle de 25 hydroxyvitamine D n'est pas associée à la survenue d'un accouchement prématuré. (*Afr J Reprod Health* 2021; 25[2]: 103-109).

Mots-clés: Naissance prématurée, vitamine D, accouchement prématuré, 25 hydroxyvitamine D, prématurée

Introduction

Vitamin D is a fat-soluble secosteroid which has multiple biological effects. It plays a significant role in calcium and phosphate homeostasis and metabolism¹. Other non-calcium functions of vitamin D have been established. These include modulation of immunity^{2,3}, stimulation of insulin production⁴, effects on myocardial contractility^{5,6}, promotion of thyroid-stimulating hormone

secretion⁷⁻⁹, and prevention of inflammatory bowel disease¹⁰. The synthesis of vitamin D involves two hydroxylation processes converting cholecalciferol (vitamin D3) and ergocalciferol (vitamin D2) to the biologically active form, $1\alpha, 25$ dihydroxyvitamin D ($1,25(\text{O.H.})_2\text{D}$) in the liver and kidney¹¹. Vitamin D3 is in the majority and is synthesised from cholesterol through a chemical reaction which depends on ultraviolet B-rays^{12,13} while vitamin D2 is ingested from diet and supplements¹⁴. During

pregnancy, 1,25(O.H.)₂D is also synthesised by human decidua and placenta tissues¹⁵. At the same time, the synthesis of 1, 25(OH)₂D by the kidneys is increased. Thus, 1,25(O.H.)₂D concentration increases during pregnancy with maximum levels reached during the third trimester¹⁶.

Worldwide, vitamin D deficiency has been reported in women of reproductive age¹⁷. Despite the sufficient and stable sunny conditions across the tropical countries enabling endogenous production of vitamin D, high prevalence of vitamin D deficiencies in pregnancy and lactating women have been reported in the literature¹⁸⁻²¹. This has been attributed to dark skin pigmentation which is prevalent in these areas and a range of lifestyle factors including the increased time they spend indoors, covered clothing due to the religious customs, liberal use of sun creams for fear of cancer²². The deficiency of vitamin D in pregnancy has been associated with various maternal and fetal complications. Maternal adverse effects include preeclampsia²³, gestational diabetes mellitus²⁴, and preterm birth²⁵, and fetal complications of small for gestation age, low birth weight, and neonatal hypocalcemia²⁶.

Preterm birth remains a global health problem with an average preterm birth rate of 11.1% every year²⁷. Malawi is one of the countries with a very high preterm birth rate of 18.1%²⁷. Various pathological processes have been identified as precursors leading to a common pathway of increased myometrial contractility, cervical dilatation and chorioamnionic membrane weakening and rupture resulting in spontaneous preterm birth (spPTB)²⁸. Infections have been causally linked to spPTB²⁹. Vitamin D is a potent immune system modulator that its deficiency correlates with increased susceptibility to infections due to impaired localised innate immunity and defective antigen-specific cellular immune response³⁰.

Considering the high preterm birth rate in Malawi and the vital role which vitamin D play in immune modulation in humans, we evaluated the status of vitamin D in spPTB in a low resource tropical country. The null hypothesis was that there is no relationship between serum concentration of vitamin D and the prevalence of spPTB.

Methods

A case-control study was conducted between June 2016 and March 2017 at Kamuzu Central Hospital (KCH) and Bwaila Hospital (B.H.), public and government-funded tertiary and general hospitals respectively. These two hospitals are 4 kilometres from each other within the city of Lilongwe, in the central region of Malawi. Neonatal services are only available at KCH. The target population was all pregnant women presenting with spPTB at the hospitals. Women presenting with spontaneous preterm birth (gestation 28 - <37 weeks) were recruited as cases and controls were women presenting with spontaneous labour at term (gestation ≥37-41 weeks). The measured outcome was maternal serum concentrations of vitamin D. Women with comorbidities (preeclampsia, gestational diabetes, and heart disease), multiple pregnancies, intrauterine growth restriction (IUGR), fetal anomaly, preterm premature rupture of membrane (PPROM) were excluded from the study. Systematic random sampling was done by allocating numbers to all eligible preterm cases and all with odd numbers recruited in the study. Controls were recruited as they presented in the labour ward and matched to the living area. This study was approved by the ethics committee of the University of Malawi, College of Medicine Research Committee (COMREC) P.05/15/1738.

All data and samples were obtained soon after delivery. After obtaining written informed consent, 5 millilitres of blood was collected from the woman using a regular vacutainer technique. The blood sample was centrifuged at 3000g for 10 minutes within 30 minutes of collection. The serum was stored at -80°C in a screw-top vial until further analysis. Vitamin D metabolites (25 Hydroxyvitamin D₂ and Hydroxyvitamin D₃) were analysed by Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) using Waters Acquity™ I-class UPLC system with an autosampler, and a binary solvent delivery system (Waters, Milford, MA) interfaced to Waters Xevo TQ-S benchtop tandem quadrupole mass spectrometer (Waters, Manchester, U.K.). Data was collected in a cross-section way on all maternal demographics, race, wearing of concealed clothing,

education status, HIV status and use of HAART. Gestation age was based on early fetal ultrasound or modified Ballard examination. Determination of haemoglobin was done using HemoCue Hb 301 (HemoCue AB, Angelhalm, Sweden).

Statistical analysis

A value of less than ≤ 50 nmol/L was taken as a threshold to define deficiency of vitamin D; insufficiency was 51 -74 nmol/L and sufficiency were 75-250 nmol/L³¹. A sample size of 90 was reached through the power of test calculation. This sample size gives 80% power to detect a significant difference at 5% level of significance. Data were processed using IBM SPSS version 20.0 software and imported into STATA 14.0 for analysis. A binary variable was generated for the primary outcome of spPTB (cases) coded as 1 and term birth (controls) coded 0. One-way frequencies were computed for categorical variables, while mean and standard deviation were calculated for continuous variables that were normally distributed. Log transformation was done where there was a presence of highly skewed data. Two-way associations between independent variables and the outcome variable of preterm birth were investigated using the Student t-test for continuous independent variables and the Chi-squared test for categorical variables. Pearson correlation was used to determine the effect of two values on each other. P-values from all tests of less than 0.05 were considered statistically significant.

Results

There were a total of 187 mothers, 92 were term birth group with mean maternal age of 26.2 years (SD: 5.9), while 95 were preterm births with mean maternal age of 23.9 years (SD: 6.1) participating in the present study (Table 1). All participating women were of the black race and wore no protective clothing. Term births group had significantly higher maternal age compared to preterm birth group ($p=0.008$). The preterm birth group were more likely to have a haemoglobin of <11 g/dl (62/95, 65.3%) compared to term birth group (42/92, 45.7%) ($p=0.007$). Gravidity, education, Human immunoglobulin Virus (HIV) status and use of HAART were not significantly

different between the two groups. Similarly, there was no difference in the concentration of 25(O.H.) D between women delivering preterm and those having a term birth, $p=0.822$.

The data was further analysed by Pearson correlation to determine the effect of maternal age, gravidity, and birth weight on micronutrients concentrations in the preterm birth group and in the term group. There was a negative correlation between maternal serum concentration of 25(O.H.) D and maternal age ($p=0.043$), as shown in Figure 1. No correlation was observed between maternal serum concentration of 25(O.H.)D and gravidity ($p=0.147$) or birth weight ($p=0.661$) in the preterm birth group.

Similarly there was no correlation between maternal serum concentration of 25(O.H.)D and maternal age ($p=0.190$) gravidity ($p=0.510$), and birth weight ($p=0.064$) in the term birth group, (Figure 2).

Discussion

The present study indicates that there is no difference in maternal concentrations of 25(OH)D between women who delivered preterm and those who had term birth. However, the study found a negative correlation between the maternal concentration of 25(O.H.) D and maternal age in the preterm birth group. Vitamin D sufficiency was found in 79.0% of the mothers who delivered preterm and 80.4% of the mothers who delivered at term, and this was not statistically different. There are no data on serum 25(OH) D concentrations in pregnant women in Malawi. Other published studies from other regions indicate lower concentrations of vitamin D in women delivering preterm. A study in Brazil looking at 92 women with preterm birth, the mean vitamin D concentrations was significantly lower in women having a preterm birth, $p=0.011$ ³². Another study in Japan found concentrations of 25(O.H.) D significantly different in preterm birth compared to term birth group³³. The study population in these studies are culturally, environmentally and, nutritionally diverse from the Malawi population.

This study design did not measure the cause-effect association. However, vitamin D deficiency is known to be associated with multiple adverse pregnancy outcomes, including preterm birth^{30,34}.

Table 1: Maternal clinical characteristics by gestation age (preterm versus term births)

Characteristic	Term births n=92	Preterm births n=95	p-value
Maternal age, Mean(SD) [†] , (years)	26.2(5.9)	23.9(6.1)	0.008*
Gestation, Mean(SD), (weeks)	38.0(1.0)	33.4(2.5)	<0.001*
Black race, N [‡] (%)	92(100)	95.0(100)	-
Wearing concealed clothing, N(%)	0 (0)	0(0)	-
Birth weight, Mean(SD) , (kg)	3.1(0.4)	2.1(0.5)	<0.001*
Gravidity 1-2, N(%)	48(52.2)	64(67.4)	0.085
At least secondary education, N(%)	48(52.2)	36(37.9)	0.075
HIV [§] positive, N(%)	15(16.3)	13(13.7)	0.300
Use of HAART [¶] , N(%)	15(16.3)	9(9.5)	0.163
Haemoglobin <11g/dl, N(%)	42(45.7)	62(65.3)	0.007*
25(O.H.)D ^{††} , N(%) , nmol/L			
≤50 (deficiency)	2(1.1)	4(4.3)	0.822
51-74 (insufficiency)	16(17.4)	16(16.8)	
75-250 (sufficiency)	74(80.4)	75(79.0)	

[†] standard deviation, [‡] number, [§] human immunoglobulin virus, [¶] Highly active antiretroviral treatment, * significant values, ^{††} 25 Hydroxyvitamin D
t-test used for continuous variables, Chi-square or Fisher’s Exact test used for categorical variables




Figure 9: Correlation between maternal serum concentrations 25(O.H.)D and maternal age, gravidity, birth weight respectively in the preterm group




Figure 2: Correlation between maternal serum concentrations 25(OH)D and maternal age, gravidity, birth weight respectively in the term birth group

Previous studies had indicated that vitamin D reduces pro-inflammatory cytokines in conditions characterised by inflammation³⁵. Inflammation is considered a precursor leading to preterm birth²⁸. The difference between term and preterm birth was not seen in our study.

In this study, we observed a significant negative correlation between maternal age and maternal serum concentrations of 25(OH) D in the preterm group which was not present in term birth group. This finding is similar to what previous studies have reported^{22,36}. Following changes in social demographics and increasing immobility associated with increasing age, women tend to stay more indoors reducing sun exposure required for vitamin D synthesis²². Deficiency of vitamin D is

also more prevalent in old age due to endocrine changes accompanying increasing age. These include the development of intestinal resistance to 1,25(OH)₂D hampering intestinal calcium uptake, decrease in the number of vitamin D receptor in various organs involved in calcium metabolism and reduced activity of 1 α -hydroxylase due to decrease in renal function, reducing the activation of vitamin D³⁷.

When interpreting these results, it should be considered that there was no information on nutrition status and body mass index for the study population. Previous studies had indicated a higher body index to be associated with decreased levels of vitamin D³⁸. The adipose tissue sequesters vitamin D. Despite this weakness, the results from

this study add on a body of evidence regarding maternal serum concentrations of vitamin D in term and preterm pregnancies in Malawi women. This study population was from two main hospitals in the central region of Malawi. We don't know if this reflects the general pregnant population in the country.

Conclusion

The study demonstrated the serum concentrations of vitamin D in pregnant women delivering at the two urban hospitals in the central region of Malawi. Serum concentrations of 25(O.H.) D were not significantly different between the preterm and term births. There was a negative correlation between maternal age and serum concentrations of vitamin D in the spPTB group. Considering that there was no correlation between increasing maternal age and maternal serum concentrations of 25(O.H.)D in the term birth group, the study recommends further research confirming this finding in other pregnant women communities' outside the Lilongwe urban, central Malawi and establish a causal relationship if indeed it exists.

Acknowledgements

The authors heartily thank Julie Sophie Kleppe Strømberg and Christina Ripman Hansen from the Department of Laboratory Medicine, University Hospital of North Norway, for facilitation during sample preparation and the Helse Nord Malawi Fund for the generous assistance of the analytical work. Ole-Martin Fuskevåg PhD did the blood serum analysis of vitamin D.

Conflict of interest

The authors have no conflict of interest.

References

- Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr.* 2004;80(6):1678S-1688S. doi:10.1093/ajcn/80.6.1678S
- Tsoukas C, Provedini D and Manolagas S. 1,25-dihydroxyvitamin D₃: a novel immunoregulatory hormone. *Science.* 1984;224(4656):1438-1440. doi:10.1126/science.6427926
- Bhalla AshokK, Amento EP, Clemens TL, Holick MF and Krane SM. Specific high-affinity receptors for 1,25-dihydroxyvitamin D₃ in human peripheral blood mononuclear cells: presence in monocytes and induction in T lymphocytes following activation. *J Clin Endocrinol Metab.* 1983;57(6):1308-1310. doi:10.1210/jcem-57-6-1308
- Lee S. 1,25-Dihydroxyvitamin D₃ and pancreatic beta-cell function: vitamin D receptors, gene expression, and insulin secretion. *Endocrinology.* 1994;134(4):1602-1610. doi:10.1210/en.134.4.1602
- Weishaar RE and Simpson RU. The Involvement of the Endocrine System in Regulating Cardiovascular Function: Emphasis on Vitamin D₃ *. *Endocr Rev.* 1989;10(3):351-365. doi:10.1210/edrv-10-3-351
- Weishaar RE, Kim SN, Saunders DE and Simpson RU. Involvement of vitamin D₃ with cardiovascular function. III. Effects on physical and morphological properties. *Am J Physiol-Endocrinol Metab.* 1990;258(1):E134-E142. doi:10.1152/ajpendo.1990.258.1.E134
- Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr.* 2004;79(3):362-371. doi:10.1093/ajcn/79.3.362
- Holick MF. McCollum Award Lecture, 1994: Vitamin D—new horizons for the 21st century. *Am J Clin Nutr.* 1994;60(4):619-630. doi:10.1093/ajcn/60.4.619
- Holick MF. Vitamin D: A millenium perspective: Vitamin D. *J Cell Biochem.* 2003;88(2):296-307. doi:10.1002/jcb.10338
- Cantorna MT, Munsick C, Bemiss C and Mahon BD. 1,25-Dihydroxycholecalciferol Prevents and Ameliorates Symptoms of Experimental Murine Inflammatory Bowel Disease. *J Nutr.* 2000;130(11):2648-2652. doi:10.1093/jn/130.11.2648
- Jones G. Pharmacokinetics of vitamin D toxicity. *Am J Clin Nutr.* 2008;88(2):582S-586S. doi:10.1093/ajcn/88.2.582S
- Holick M, MacLaughlin J, Clark M, Holick SA, Potts JT, Anderson RR, Blank IH, Parrish JA and Elias P. Photosynthesis of previtamin D₃ in human skin and the physiologic consequences. *Science.* 1980;210(4466):203-205. doi:10.1126/science.6251551
- Holick MF, Chen TC, Lu Z and Sauter E. Vitamin D and Skin Physiology: A D-Lightful Story. *J Bone Miner Res.* 2007;22(S2):V28-V33. doi:10.1359/jbmr.07s211
- Calvo MS, Whiting SJ and Barton CN. Vitamin D Intake: A Global Perspective of Current Status. *J Nutr.* 2005;135(2):310-316. doi:10.1093/jn/135.2.310
- Weisman Y, Harell A, Edelstein S, David M, Spirer Z and Golander A. *Alpha, 25-Dihydroxyvitamin D₃ and 24, 25-Dihydroxyvitamin D₃ in Vitro Synthesis by Human Decidua and Placenta Nature 1979 281. N; 1.*
- Kovacs CS and Kronenberg HM. Maternal-fetal calcium

- and bone metabolism during pregnancy, puerperium, and lactation. *Endocr Rev.* 1997;18(6):832-872. doi:10.1210/er.18.6.832
17. Mithal A, Wahl DA, Bonjour J-P, P Burckhardt, B Dawson-Hughes, Eisman JA, El-Hajj Fuleihan, RG Josse, P Lips and J Morales-Torres. Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int.* 2009;20(11):1807-1820.
 18. Nair R and Maseeh A. Vitamin D: The “sunshine” vitamin. *J Pharmacol Pharmacother.* 2012;3(2):118-126. doi:10.4103/0976-500X.95506
 19. Toko EN, Sumba OP, Daud II, Ogolo S, Majiwa M, Krisher JT, Ouma C, Dent AE, Rochford R and Mehta S. Maternal Vitamin D Status and Adverse Birth Outcomes in Children from Rural Western Kenya. *Nutrients.* 2016;8(12):794. doi:10.3390/nu8120794
 20. Nimitphong H and Holick MF. Vitamin D status and sun exposure in southeast Asia. *Dermatoendocrinol.* 2013;5(1):34-37. doi:10.4161/derm.24054
 21. Mehta S, Spiegelman D, Aboud S, Aboud S, Giovannucci EL, Msamanga GI, Hertzmark E, Mugusi FM, Hunter DJ and Fawzi WW. Lipid-soluble vitamins A, D and E in HIV-infected pregnant women in Tanzania. *Eur J Clin Nutr.* 2010;64(8):808-817. doi:10.1038/ejcn.2010.76
 22. Lips P, Schoor NM van and Jongh RT de. Diet, sun, and lifestyle as determinants of vitamin D status. *Ann N Y Acad Sci.* 2014;1317(1):92-98. doi:10.1111/nyas.12443
 23. Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW and Roberts JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab.* 2007;92(9):3517-3522.
 24. Poel YHM, Hummel P, Lips P, Stam F, Van Der Ploeg T and Simsek S. Vitamin D and gestational diabetes: a systematic review and meta-analysis. *Eur J Intern Med.* 2012;23(5):465-469.
 25. Bodnar LM, Platt RW and Simhan HN. Early-Pregnancy Vitamin D Deficiency and Risk of Preterm Birth Subtypes. *Obstet Gynecol.* 2015;125(2):439-447. doi:10.1097/AOG.0000000000000621
 26. van der Pligt P, Willcox J, Szymlek-Gay E, Murray E, Worsley A and Daly R. Associations of Maternal Vitamin D Deficiency with Pregnancy and Neonatal Complications in Developing Countries: A Systematic Review. *Nutrients.* 2018;10(5):640. doi:10.3390/nu10050640
 27. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A, Narwal R, Adler A, Garcia CV, Rohde S, Say L and Lawn JE. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *The Lancet.* 2012;379(9832):2162-2172. doi:10.1016/S0140-6736(12)60820-4
 28. Romero R, Espinoza J, Kusanovic J, Gotsch F, Hassan S, Erez O, Chaiworapongsa T and Mazor M. The preterm parturition syndrome. *BJOG Int J Obstet Gynaecol.* 2006;113:17-42. doi:10.1111/j.1471-0528.2006.01120.x
 29. Gonçalves LF, Chaiworapongsa T and Romero R. Intrauterine infection and prematurity. *Ment Retard Dev Disabil Res Rev.* 2002;8(1):3-13. doi:10.1002/mrdd.10008
 30. Adams JS and Hewison M. Unexpected actions of vitamin D: new perspectives on the regulation of innate and adaptive immunity. *Nat Clin Pract Endocrinol Metab.* 2008;4(2):80-90. doi:10.1038/ncpendmet0716
 31. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH and Weaver CM. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2011;96(7):1911-1930. doi:10.1210/jc.2011-0385
 32. Kassai MS, Cafeo FR, Affonso-Kaufman FA, Suano-Souza FI and Sarni ROS. Vitamin D plasma concentrations in pregnant women and their preterm newborns. *BMC Pregnancy Childbirth.* 2018;18(1):412. doi:10.1186/s12884-018-2045-1
 33. Shibata M, Suzuki A, Sekiya T, Sekiguchi S, Asano S, Udagawa Y, and Itoh M. High prevalence of hypovitaminosis D in pregnant Japanese women with threatened premature delivery. *J Bone Miner Metab.* 2011;29(5):615-620. doi:10.1007/s00774-011-0264-x
 34. Zhou S-S, Tao Y-H, Huang K, Zhu B-B and Tao F-B. Vitamin D and risk of preterm birth: Up-to-date meta-analysis of randomized controlled trials and observational studies. *J Obstet Gynaecol Res.* 2017;43(2):247-256. doi:10.1111/jog.13239
 35. Yin K and Agrawal DK. Vitamin D and inflammatory diseases. *J Inflamm Res.* 2014;7:69.
 36. Baker MR, Peacock M and Nordin BEC. The decline in vitamin d status with age. *Age Ageing.* 1980;9(4):249-252. doi:10.1093/ageing/9.4.249
 37. Lips P. Vitamin D physiology. *Prog Biophys Mol Biol.* 2006;92(1):4-8. doi:10.1016/j.pbiomolbio.2006.02.016
 38. Konradsen S, Ag H, Lindberg F, Hexeberg S and Jorde R. Serum 1, 25-dihydroxy vitamin D is inversely associated with body mass index. *Eur J Nutr.* 2008;47(2):87-91.