

Nurten Ozsoy<sup>1\*</sup> Alper Okyar<sup>2</sup> Pelin Arda-Pirincci<sup>3</sup> Ayse Can<sup>1</sup> Sehnaz Bolkent<sup>3</sup> Nuriye Akev<sup>1</sup>

<sup>1</sup>Department Of Biochemistry, Faculty Of Pharmacy, Istanbul University, Istanbul 34116, Turkey

<sup>2</sup>Department Of Pharmacology, Faculty Of Pharmacy, Istanbul University, Istanbul 34116, Turkey

<sup>3</sup>Department Of Biology, Faculty Of Science, Istanbul University, Istanbul 34116, Turkey

\*corresponding auhtor E-Mail: [nozsoy@istanbul.edu.tr](mailto:nozsoy@istanbul.edu.tr)

## Abstract

**Background:** Because reactive oxygen species (ros) contribute to the pathogenesis of various acute and chronic liver diseases, dietary antioxidants and drugs from herbal origins have been proved to be beneficial as therapeutic agents in reversing hepatotoxicity and oxidative stress. The objective of this study was to investigate the protective effect of an aqueous extract from *smilax excelsa* l. Shoots and leaves against acute ccl<sub>4</sub>-induced liver injury as well as the changes in antioxidative defense system in female wistar albino rats.

**Materials and Methods:** *S. Excelsa* extract was administered orally in doses of 100, 200 and 400 mg/kg body weight, once daily for 9 days. Acute hepatic toxicity was induced by intraperitoneal injection of ccl<sub>4</sub> (1 ml/kg) on the 10<sup>th</sup> day. 24 h after ccl<sub>4</sub> intoxication, biochemical and histopathological analyses were undertaken on sera and liver tissues.

**Results:** Ccl<sub>4</sub> challenge caused significant increases in the activities of liver enzymes as well as the levels of bilirubin, malondialdehyde and nitric oxide, while total serum protein levels and antioxidant defense system parameters were reduced significantly compared to the normal group. Administration of *s. Excelsa* extract at a dose of 400 mg/kg resulted in a suppression of ccl<sub>4</sub>-induced lipid peroxidation and altered oxidative stress parameters to nearly normal values in comparison to ccl<sub>4</sub>-treated rats. Nevertheless the extract did not reduce the extent of ccl<sub>4</sub>-induced mild liver injury, as seen by the histopathology of liver damage.

**Conclusion:** The results of this study suggest that *s. Excelsa* could protect the liver tissues against ccl<sub>4</sub>-induced oxidative stress probably by increasing antioxidative defense activities.

**Key words:** Antioxidant enzymes, carbon tetrachloride, liver injury, *smilax excelsa*, hepatoprotective activity

## Introduction

For centuries, medicinal plants have been used as remedies for human diseases because they contain phytoconstituents of therapeutic value. These phytochemicals may be useful in preventing oxidative stress-related diseases by fortifying antioxidant defense mechanisms or by acting as protective factors.

Liver diseases remain one of the serious health problems, but medical treatments for these diseases are often difficult to handle and have limited efficacy. For these reasons, developing drugs for liver diseases from plants used in traditional medicine, may lead to improved therapies (kim et al., 2009).

Based on reports regarding the presence of polyphenols in shoots and leaves of *smilax excelsa*, a plant species from the black sea region of turkey, and the assumption that the traditional uses of *s. Excelsa* might be partially attributed to its antioxidant activity (ozsoy et al., 2008), the aim of this study was to evaluate the potential protective effect of *s. Excelsa* against the oxidative stress induced by carbon tetrachloride (ccl<sub>4</sub>) in the rat liver.

## Material and Methods

### Plant Material

*S. Excelsa* l. Shoots and leaves were collected in september from istanbul in turkey and identified by prof. Dr. Kerim alpinar from the faculty of pharmacy, istanbul university. The leaves were separated from the other parts, washed in running tap water and dried at room temperature. Voucher specimens are deposited in the herbarium of the faculty of pharmacy, istanbul university (iste); herbarium code number: iste 81928. The dried shoots and leaves were manually ground to a fine powder before extraction.

### Preparation of the Extract

80 g of ground dried shoots and leaves were extracted with boiling water (2000 ml) for 15 min while stirring. The extracts were filtered and evaporated to dryness under reduced pressure at 40°C in a rotary evaporator (buchi r210). The yield was 25 g. Appropriate dilutions were made before each experiment.

### Animals

The experimental protocol described in the present study was approved by the animal assays ethics committee of istanbul university (hadyek). Female wistar rats weighing 200-240 g were supplied from istanbul university institute of experimental medicine (detae). Animals were acclimatized to their environment for one week prior to experimentation, housed in a room with a 12 h light/dark cycle at about 22°C and fed on standard diet with *ad libitum* access to drinking water.

### Induction of Hepatic Injury

The animals were divided into 5 groups each containing 6 animals. Group 1 (control), which served as normal control, received water and basal diet for 10 days. Group 2 (ccl<sub>4</sub>) which served as toxin control, received water and basal diet for 10 days and was treated with ccl<sub>4</sub> (1 ml/kg body weight in 20 % olive oil, v/v) on the 10 th day. Three different groups of rats, groups 3, 4 and 5 were separately treated with three different doses of the leaf extract of *s. Excelsa* – 100, 200 and 400 mg/kg body weight respectively, once daily for 9 days, 24 h prior to intraperitoneal administration of ccl<sub>4</sub> on the 10th day.

24 h after ccl<sub>4</sub> intoxication, the rats were anesthetized and blood samples were taken from each rat through direct intracardiac intervention and centrifuged at 3.000 x g for 10 min to separate the serum. Immediately after collecting the blood samples, the livers were excised, rinsed in ice-cold normal saline solution followed by ice-cold 0.15 m potassium phosphate buffer (ph 7.4), blotted, dried, and weighed. Part of liver tissues were used for histopathological examination. With another part of liver tissues, 10 % w/v homogenates were prepared in ice-cold 0.15 m potassium phosphate buffer using a homogenator (art-micra d-1), and centrifuged at 13.000 rpm for 5 min at 4°C (megafuge hereaus1.0r). The supernatants, thus obtained were used for biochemical studies.

### Biochemical Studies

#### Assessment of Serum Biochemical Parameters and Lipid Peroxidation

Serum aspartate aminotransferase (ast) and alanine aminotransferase (alt) activities were estimated as prescribed by the international federation of clinical chemistry (bergmeyer, 1980; bergmeyer et al., 1986). A unit of enzyme activity was defined as 1 μmol of nadh or nad formed per minute under the assay conditions. Alkaline phosphatase (alp) activity was determined in serum using p-nitrophenyl phosphate as a substrate and one unit of enzyme activity was defined as μmol p-nitrophenol formed per minute (walter and schütt, 1974). Serum bilirubin was estimated by the diazo method (balistreri and rej, 1994). The formation of lpo products was assayed by the measurement of thiobarbituric acid reactive substances (tbars) levels on the basis of malondialdehyde (mda) reaction with thiobarbituric acid at 532 nm, according to buege and aust (1978). The values of thiobarbituric acid related substances (tbars) were calculated using an extinction coefficient of 1.56x10<sup>5</sup> m<sup>-1</sup>cm<sup>-1</sup> and expressed as nmol of mda/g wet weight.

Serum nitric oxide (no) was measured in terms of its products, nitrite and nitrate, by the method of griess modified by (fiddler, 1977) after samples, after deproteination, were incubated with cadmium beads in order to convert nitrate to nitrite, and nitrite levels were determined spectrophotometrically at 540 nm by using a colorimetric assay based on the griess reaction, no products were expressed as μmoles/l serum.

#### Assessment of Antioxidant Status via Antioxidant Enzymes in Liver Homogenates

Superoxide dismutase (sod) activity was assayed by the method described by aruoma et al. (1989). Results were expressed as u/mg protein. One unit of sod inhibits the rate of increase in absorbance at 560 nm by 50 %, under the conditions of the assay. The protein contents were determined by the method of lowry et al. (1951) using bovine serum albumin (bsa) as standard. Catalase (cat) activity was measured by the method of aebi (1974) and expressed as μmol h<sub>2</sub>O<sub>2</sub>/mg protein. Glutathione reductase (gr) activity was determined following the oxidation of nadph at 340 nm as described by carlberg and mannervik (1985). The activity of glutathione peroxidase (gpx) was measured using a coupled enzyme assay system linked with gr as described by lawrence and burk (1976). Gr and gpx activities were expressed as nmol of nadph oxidized per min per mg protein using the molar extinction coefficient for nadph at 340 nm of 6.22x10<sup>6</sup>m<sup>-1</sup>cm<sup>-1</sup>. Glutathione-s-transferase (gst) activity was assayed spectrophotometrically using 1-chloro-2,4-dinitrobenzene as substrate as described by habig and jacob (1981). Specific activity was expressed as nmol conjugate formed/min/mg protein using a molar extinction coefficient of 9.6 m<sup>-1</sup>cm<sup>-1</sup>.

Tissue myeloperoxidase (mpo) levels were measured using a method described by singbartl et al. (2000) with some modifications. 1 u of activity was defined as the amount of enzyme that consumes 1 μmol h<sub>2</sub>O<sub>2</sub>/min at 25°C. Results were expressed as units of mpo per g wet tissue.

The protein carbonyl content (pcc) was assayed by the modification of the procedure described by reznick and packer (1994), using dinitrophenylhydrazine (dnph). Results were expressed as nmol of protein carbonyl per mg of protein using a molar extinction coefficient of 22.000 m<sup>-1</sup>cm<sup>-1</sup> for dnph.

The level of the reactive oxygen species (ros) scavenger, reduced glutathione (gsh) was measured in both liver and tissues by the enzymic recycling procedure using 5,5'-dithiobis-(2-nitrobenzoic acid) (dtNB) and glutathione reductase (dtNB-gssg reductase recycling assay) as described by anderson (1985). The oxidized glutathione (gssg) level was measured using the same method but in presence of 2-vinylpyridine. The results were expressed as μmol of gsh (gssg)/g of wet weight.

### Histopathological Assay

Dissected liver tissues were taken immediately after sacrifice and fixed in bouin's solution for histopathological examinations. The tissues were dehydrated and embedded in paraffin. The paraffin sections of 5 μm thickness were stained with hematoxylin-eosin (h&e) and masson's trichrome stain (masson). All sections were examined under olympus-cx 41 light microscope.

### Statistical Analysis

Biochemical results were evaluated using an unpaired *t*-test and anova variance analysis using the ncss statistical computer package. The values were expressed as mean ± sd. Analysis between control and experimental groups was performed using the mann-whitney test. *P* < 0.05 was considered as significant.

**Results**

**Assessment of Serum Biochemical Parameters and Lipid Peroxidation**

Results showed in Table 1, revealed that pretreatment of rats with an aqueous extract of *s. Excelsa* at 100, 200 and 400 mg/kg doses, significantly attenuated the rise in serum level of ast, alt and total bilirubin, provoked by ccl<sub>4</sub>. Treatment with *s. Excelsa* aqueous extract with three doses, prior to the ccl<sub>4</sub> administration decreased also the elevation of serum lipid peroxidation (as shown by the reduction of the enhanced mda levels). Increased no levels in the ccl<sub>4</sub> group were significantly reduced by *s. Excelsa* at 200 and 400 mg/kg doses.

**Table 1:** Effect of pretreatment with *Smilax excelsa* L. leaf aqueous extract after CCl<sub>4</sub> intoxication, on serum biochemical parameters and lipid peroxidation in rats

Group	Protein	AST	ALT	ALP	Bilirubin	MDA	NO
	g/dl	U/L	U/L	U/L	mg/dl	µmol/L	µmol/L
Control	7.5 ± 0.9	44.8 ± 16.7	17.5 ± 5.70	261.6 ± 50.0	0.2 ± 0.04	1.4 ± 0.3	1.8 ± 0.8
CCl <sub>4</sub>	7.2 ± 1.2	1616.5 ± 53.3 <sup>a</sup>	613.6 ± 105.9 <sup>a</sup>	226.7 ± 80.7	1.6 ± 0.3 <sup>a</sup>	3.4 ± 0.4 <sup>a</sup>	34.1 ± 14.6 <sup>d</sup>
<i>S.excelsa</i> <sub>100</sub> +CCl <sub>4</sub>	7.2 ± 0.8	1026.2 ± 87.8 <sup>a,b</sup>	386.0 ± 83.7 <sup>a,c</sup>	234.1 ± 22.0	0.9 ± 0.5 <sup>d,e</sup>	2.5 ± 0.4 <sup>a,c</sup>	21.1 ± 4.7 <sup>a</sup>
<i>S.excelsa</i> <sub>200</sub> +CCl <sub>4</sub>	6.9 ± 1.1	1033.7 ± 99.0 <sup>a,b</sup>	252.9 ± 13.6 <sup>a,b</sup>	214.1 ± 28.3	1.1 ± 0.3 <sup>a,e</sup>	2.7 ± 0.6 <sup>c,d</sup>	4.8 ± 3.5 <sup>e</sup>
<i>S.excelsa</i> <sub>400</sub> +CCl <sub>4</sub>	7.3 ± 0.6	203.9 ± 55.4 <sup>a,b</sup>	64.5 ± 12.9 <sup>a,b</sup>	231.7 ± 23.9	0.3 ± 0.1 <sup>b</sup>	1.6 ± 0.4 <sup>b</sup>	3.2 ± 1.1 <sup>b,d</sup>
<i>P</i> <sub>ANOVA</sub>	0.811	0.0001	0.0001	0.470	0.0001	0.0001	0.0001

<sup>a</sup>p<0.0001 vs Control; <sup>b</sup>p<0.0001 vs CCl<sub>4</sub>; <sup>c</sup>p<0.005 vs CCl<sub>4</sub>; <sup>d</sup>p<0.05 vs Control; <sup>e</sup>p<0.05 vs CCl<sub>4</sub>; Results are given as mean ± SD.

**KEY**

AST = aspartate aminotransferase; ALT = alanine aminotransferase; ALP = alkaline phosphatase; MDA = malondialdehyde  
NO = nitric oxide

**Assessment of Antioxidant Status via Antioxidant Enzymes in Liver Homogenates**

Pretreatment with *s. Excelsa* leaf extract restored the impaired parameters of sod, cat, gr, gp<sub>x</sub>, gst, mpo activities as well as the levels of pcc in liver tissue (Table 2). Tissue lipid peroxidation levels measured as mda and the levels of the glutathione system (gsh, gssg and gssg/gsh ratio) were also significantly reversed by pretreatment with *s. Excelsa* extracts at the three doses given (Table 3).

**Table 2.** Effect of pretreatment with *Smilax excelsa* L. leaf aqueous extract after CCl<sub>4</sub> intoxication, on liver tissue antioxidant enzymes in rats

Group	Protein	SOD	CAT	GR	GPx	GST	MPO	PCC
	mg/ml	U/mg protein	U/mg protein	U/mg protein	U/mg protein	U/mg protein	U/g tissue	nmol PC/mg protein
Control	9.0 ± 1.2	1.65 ± 0.2	85.9 ± 15.1	52.4 ± 9.4	1120.8 ± 99.8	785.9 ± 75.9	1.1 ± 0.4	2.5 ± 0.5
CCl <sub>4</sub>	7.6 ± 1.1 <sup>a</sup>	2.46 ± 0.7 <sup>a</sup>	60.0 ± 8.8 <sup>a</sup>	31.5 ± 4.1 <sup>c</sup>	826.8 ± 47.7 <sup>b</sup>	561.1 ± 42.7 <sup>b</sup>	30.9 ± 7.4 <sup>b</sup>	23.3 ± 3.9 <sup>b</sup>
<i>S.excelsa</i> <sub>100</sub> +CCl <sub>4</sub>	6.9 ± 1.1 <sup>a</sup>	2.72 ± 0.2 <sup>b</sup>	66.1 ± 31.6	48.3 ± 9.3 <sup>d</sup>	1103.6 ± 88.1 <sup>f</sup>	877.4 ± 134.2 <sup>f</sup>	30.8 ± 12.9 <sup>b</sup>	3.2 ± 0.9 <sup>f</sup>
<i>S.excelsa</i> <sub>200</sub> +CCl <sub>4</sub>	6.5 ± 0.4 <sup>b,c</sup>	2.54 ± 0.1 <sup>b</sup>	82.3 ± 13.5 <sup>c</sup>	45.1 ± 10.9 <sup>d</sup>	1208.4 ± 100 <sup>f</sup>	545.4 ± 116.1 <sup>e</sup>	35.1 ± 9.8 <sup>b</sup>	3.6 ± 1.5 <sup>f</sup>
<i>S.excelsa</i> <sub>400</sub> +CCl <sub>4</sub>	8.8 ± 1.6	1.75 ± 0.4 <sup>c</sup>	84.4 ± 11.7 <sup>d</sup>	62.0 ± 10.9 <sup>f</sup>	1169.0 ± 109.7 <sup>f</sup>	757.5 ± 174.5 <sup>e</sup>	11.3 ± 4.8 <sup>b,f</sup>	2.2 ± 0.5 <sup>f</sup>
<i>P</i> <sub>ANOVA</sub>	0.001	0.0001	0.070	0.0001	0.0001	0.0001	0.0001	0.0001

**Key**

Sod = superoxide dismutase; cat = catalase; Gr = glutathione reductase; Gpx = glutathione peroxidase ;Gst = glutathione-s-transferase  
Pcc = protein carbonyl content

**Table 3:** Effect of pretreatment with *Smilax excelsa* L. Leaf aqueous extract after  $\text{CCl}_4$  intoxication, on the formation of tbars and glutathione system in the liver tissue in rats

Group	Mda	Gsh	Gssg	Gssg/gsh
	Nmol/g	$\mu\text{mol/g}$	$\mu\text{mol/g}$	-
Control	30.7 $\pm$ 4.5	3.9 $\pm$ 0.3	0.33 $\pm$ 0.04	0.091 $\pm$ 0.007
$\text{CCl}_4$	60.0 $\pm$ 10.3 <sup>a</sup>	2.2 $\pm$ 0.2 <sup>a</sup>	0.61 $\pm$ 0.16 <sup>g</sup>	0.156 $\pm$ 0.02 <sup>a</sup>
<i>S.excelsa</i> <sub>100+CCl4</sub>	40.2 $\pm$ 6.0 <sup>b,c</sup>	3.5 $\pm$ 0.9 <sup>f</sup>	0.39 $\pm$ 0.09 <sup>f</sup>	0.091 $\pm$ 0.02 <sup>c</sup>
<i>S.excelsa</i> <sub>200+CCl4</sub>	41 $\pm$ 6.3 <sup>d,c</sup>	2.3 $\pm$ 0.4 <sup>d</sup>	0.27 $\pm$ 0.04 <sup>b,c</sup>	0.136 $\pm$ 0.01 <sup>g</sup>
<i>S.excelsa</i> <sub>400+CCl4</sub>	35.1 $\pm$ 0.5 <sup>b,e</sup>	3.9 $\pm$ 0.7 <sup>c</sup>	0.25 $\pm$ 0.007 <sup>c</sup>	0.095 $\pm$ 0.02 <sup>c</sup>
<i>P</i> <sub>anova</sub>	0.0001	0.0001	0.0001	0.019

<sup>a</sup> $p < 0.0001$  vs control; <sup>b</sup> $p < 0.05$  vs control; <sup>c</sup> $p < 0.005$  vs  $\text{CCl}_4$ ; <sup>d</sup> $p < 0.01$  vs control;


<sup>e</sup> $p < 0.0001$  vs  $\text{CCl}_4$ ; <sup>f</sup> $p < 0.05$  vs  $\text{CCl}_4$ ; <sup>g</sup> $p < 0.005$  vs control; results are given as mean  $\pm$  sd.

**Key**

Gsh = reduced glutathione; Gssg = oxidized glutathione; Mda = malondialdehyde

**Histopathological Assessment**

The results of the light microscopic observations showed that the administration of  $\text{CCl}_4$  produces moderate degenerative changes such as necrosis, vacuolisation and hypertrophy in hepatocytes, hyperemia, mononuclear cell infiltration, necrotic areas, increase in the dark eosinophilic cells, the rupturing in the central vein of liver tissues in the rats. The degeneration was observed around central veins in the liver tissues of the rats given  $\text{CCl}_4$  (fig. 1 b and c) in comparison to normal control rats (fig. 1 a). In addition, the foamy cells around central veins were noticed in some animals of the experimental group. Similar degenerative changes were also detected in rats given *s. Excelsa* extract at 100 mg/kg (fig. 1 d), 200 mg/kg (fig. 1 e) or 400 mg/kg (fig. 1 f) dose together with  $\text{CCl}_4$ . The degeneration was observed to portal area from central vein in the groups given *s. Excelsa* extracts. The foamy cells around central veins were also observed still in some animals of experimental groups given *s. Excelsa* extracts (fig. 1 d, e, f). Pretreatment with *Smilax excelsa* extracts in the different doses did not show a protective effect in the liver tissue of  $\text{CCl}_4$ -treated rats, morphologically.



**Figure 1:** histological appearance of liver tissue of a control rat (a). Light micrographs of  $\text{CCl}_4$  group (b and c), *s. Excelsa* (100 mg/kg)-treated  $\text{CCl}_4$  group (d), *s. Excelsa* (200 mg/kg)-treated  $\text{CCl}_4$  group (e), and *s. Excelsa* (400 mg/kg)-treated  $\text{CCl}_4$  group (f). Hyperemia (\*) in central vein (cv), hypertrophy (↔) and vacuolisation (→) in hepatocytes, necrotic area (▶), mononuclear cell infiltration (mci), dark eosinophilic cells (▶), the rupturing of the central vein (→). H&e, original magnification x200.

**Discussion**

Detoxification of xenobiotic and toxic substances is an important hepatic function, in the course of which large amounts of reactive oxygen intermediates are produced. The excretion of many drugs depends on the integrity of the liver and cytochrome p-450 system (balistreri and rej, 1994).

In this study the experimental intoxication induced by  $\text{ccl}_4$  (mukai et al., 2002) was used for modeling liver injury in rats. Hepatotoxicity is connected with severe impairment of cell protection mechanisms. All acute injuries and/or necrotic lesions in the liver primarily cause a marked rise in the levels of the aminotransferases, ast and alt. Pretreatment of rats with an aqueous extract of *s. Excelsa* attenuated the  $\text{ccl}_4$ -induced rise in serum level of transaminases and total serum bilirubin, confirming the protective effect of the extract. Hepatic damage was also supported by the elevation of lipid peroxidation (expressed as mda) levels. Treatment with *s. Excelsa* produced a significant attenuation in  $\text{ccl}_4$ -induced increase in lipid peroxidation.

Sod is the first antioxidant enzyme to deal with oxidative free radicals by accelerating the dismutation of superoxide to hydrogen peroxide, while cat is a peroxisomal heme protein that catalyses the removal of hydrogen peroxide formed during the reaction catalyzed by sod. Thus, sod and cat act as mutually supportive antioxidative enzymes, and provide protective defense against ros. The increase in the activity of sod in the liver of  $\text{ccl}_4$ -treated rats may be due to the increased accumulation of superoxide radicals, which could further stimulate lipid peroxidation. Superoxide is inactivated by sod, the only enzyme known to use a free radical as a substrate. Any increase in sod activity is beneficial in the event of increased free radical generation. However, a rise in sod activity, without a concomitant rise in the activity of cat and/or gpx might be detrimental since sod generates hydrogen peroxide as a metabolite, which must be scavenged by cat or gpx. Thus, a simultaneous increase in catalase and/or gpx activity, as observed in *s. Excelsa* treated groups, is essential for an overall beneficial effect of an increase in sod activity (banerjee et al., 2002).

Glutathione antioxidant system plays a fundamental role in cellular defense against reactive free radicals and other oxidant species. This system consisted of reduced glutathione and an array of functionally-related enzymes, one of which is glutathione reductase, responsible for the regeneration of gsh. On the other hand, gpx and gst work together with gsh in the decomposition of hydrogen peroxide or other organic hydroperoxides. If the injury is associated with a generalized increase in tissue oxidative stress, it might well be reflected in the changes in tissue glutathione antioxidant system (mak et al., 1996). In this study, data consistent with the glutathione system also suggested high cell damage induced by  $\text{ccl}_4$  in the  $\text{ccl}_4$ -treated groups. Twenty four hours after  $\text{ccl}_4$  administration, hepatic gsh levels decreased significantly in  $\text{ccl}_4$  group, while gssg levels were elevated. The decrease in the gssg/gsh ratio, a potent indicator of tissue oxidative stress, in groups pretreated with 100, 200 and 400 mg/kg *s. Excelsa* aqueous extract, is also in accordance with the suggestion that free radical damage is decreased in the tissues of *s. Excelsa* treated groups.

Gst is a phase II enzyme that plays a key role in cellular detoxification of xenobiotics, electrophiles and ros through their conjugation to gsh. The decrease of hepatic gst might well be an adaptive response to the increased production of oxidized glutathione in tissues of  $\text{ccl}_4$ -treated rats (akerboom and sies, 1990).

Given that gpx works together with gsh in the decomposition of hydroperoxides generated from free radical-mediated reactions, the induction of gpx activity and the accompanied increase in liver glutathione level may merely be a manifest of antioxidant response to the increased oxidative stress (mak et al., 1996). Our findings of the concomitant increase in hepatic gpx and gst activities and gsh levels in rats treated with *s. Excelsa* aqueous extract (in the group treated with 400 mg/kg aqueous extract this increase was almost close to control group) are consistent with these observations.

Kupffer cell mpo may be an important source of oxidative damage during acute and chronic liver (brown et al., 2001). The hepatic levels of mpo in the  $\text{ccl}_4$ -treated groups were higher than in the normal control group, and there was a significant difference between  $\text{ccl}_4$ -treated and control groups. Treatment with 400 mg/kg *s. Excelsa* extract significantly reduced the levels of mpo in tissue. Thus, the results indicate that treatment with *s. Excelsa* extract causes significant suppression of neutrophil infiltration.

Measurement of protein carbonyls has been used as a sensitive assay for oxidative damage to proteins, partly because it measures several different consequences of oxidative damage (halliwell and whiteman, 2004). The results showed increased protein damage by ros in  $\text{ccl}_4$  groups. The treatment with 400 mg/kg *s. Excelsa* aqueous extract might have modified the oxidative status by increasing antioxidant defense.

As no has a range of effects on a variety of biological processes, it is unclear if the rise of no serum levels is beneficial or detrimental in the liver injury induced by hepatotoxins (kim et al., 2009). In this study the hepatotoxin has made a significant increase in serum no levels which has been significantly decreased at 200 and 400 mg/kg extract doses.

In view of these observations, we hypothesized that *s. Excelsa* extract would have protective effects in an *in vivo* induced free radical-mediated hepatotoxicity by attenuating the  $\text{ccl}_4$ -mediated oxidative stress through decreased production of free radical derivatives, as evidenced by the decreased mda, increased activity of phase II detoxifying enzyme (gst) and antioxidant enzymes (sod, cat, gpx and gr), and protection against lpo and gsh depletion. Accordingly, in our other report *s. Excelsa* shoots and leaves were proved as nephroprotective under *in vivo* conditions (ozsoy et al., 2013). Our results were consistent with earlier studies which have also shown that antioxidants could comprehensively ameliorate the oxidative damage by increasing antioxidative potential (yuan et al., 2014; gu et al., 2014).

Despite this conclusion, the improvement in antioxidant enzymes system seem to have been incapable of neutralizing increased  $\text{ccl}_4$  toxicity seen in the histopathology of liver cells. In the present study, the results of histological observations showed that  $\text{ccl}_4$  leads to moderate degenerative changes mainly necrosis, hyperemia, and vacuolar degeneration in the liver of rats. Our histopathological findings were in agreement with the degenerative structural changes reported to occur in liver tissues after the application of  $\text{ccl}_4$  (jaramillo-juárez et al., 2008; el denshary et al., 2011). In this study, pretreatment of the aqueous extracts of *s. Excelsa* shoots and leaves did not diminished  $\text{ccl}_4$ -induced degenerative injury in liver tissue, morphologically. It was stated that sometimes there could not be strict correlation between histological findings and serum transaminase values (kallai et al. 1964) and that entire histologic spectrum of liver disease can be seen in individuals with normal alt values (mofrad et al., 2003). It was also reported that alt values do not sometimes correlate with parameters of apoptosis and oxidative stress and that the severity of the disease could be only determined by liver biopsy (canbakan et al., 2010). Accordingly, the results of the present study showed that, at early stages of liver injury, biochemical findings could not always correlate with histological examination and that an antioxidant effect could not always be along with a restored tissue structure.

In conclusion, our study demonstrated that the aqueous extract of the shoots and leaves of *s. Excelsa* could increase antioxidative defense system against ccl<sub>4</sub>-induced oxidative stress.

## Acknowledgement

This study was supported by Istanbul university research projects (bap). Project numbers: 1402;udp-8368 and udp-7724

## References

1. Aebi, H. (1984). Catalase *in vitro*. Methods enzymol., **105**: 121-126.
2. Akerboom, T. and Sies, H. (1990). Glutathione transport and its significance in oxidative stress. In, viña j (ed). Glutathione: metabolism and physiological functions. 45-55, crc press, florida.
3. Anderson, M.E. (1985). Determination of glutathione and glutathione disulfide in biological samples. Methods. Enzymol., **113**: 548-555.
4. Aruoma, O.I., Halliwell, B., Hoey, B.M. and Butler, J. (1989). The antioxidant action of n-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide, and hypochlorous acid. Free radic. Biol. Med., **6**: 593-597.
5. Balistreri, W.F. and Rej, R. (1994). Liver function. Determination of total and conjugated bilirubin in serum (jeddrassik and gróf technique). In, burtis ca, ashwood er (eds): tietz textbook of clinical chemistry. 1466-1468, w.b. Saunders company, philadelphia.
6. Banerjee, S.K., Maulik, M., Mancahanda, S.C., Dinda, A.K., Gupta, S.K. and Maulik, S.K. (2002). Dose-dependent induction of endogenous antioxidants in rat heart by chronic administration of garlic. Life sci., **70**: 1509-1518.
7. Bergmeyer, H.U., Horder, M. and Rej, R. (1986). International federation of clinical chemistry (ifcc). Scientific committee, analytical section: approved recommendation (1985) on IFCC methods for the measurement of catalytic concentration of enzymes. Part 2. IFCC method for aspartate aminotransferase (l-aspartate: 2-oxoglutarate aminotransferase, ec 2.6.1.1). J. Clin. Chem. Clin. Biochem., **24**: 497-510.
8. Bergmeyer, H.U. (1980). Ifcc methods for measurement of catalytic concentrations of enzymes. Part 3, ifcc. Method for alanine aminotransferase (l-alanine 2-oxoglutarate aminotransferase, ec 2.6.1.2). Clin. Chim. Acta, **105**: 145f-172f.
9. Brown, K.E., Brunt, E.M. And Heinecke, J.W. (2001). Immunohistochemical detection of myeloperoxidase and its oxidation products in kupffer cells of human liver. Am. J. Pathol. (ajp), **159**: 2081-2088.
10. Buege, J.A. and Aust, S.D. (2010). Microsomal lipid peroxidation. Methods enzymol., **52**: 302-310.
11. Canbakan, B., Senturk, H., Canbakan, M., Toptas, T., Tabak, O., Balci, H., Olgac, V., and Ozbay, G. (2010). Is alanine aminotransferase level a surrogate biomarker of hepatic apoptosis in nonalcoholic fatty liver disease? Biomark. Med., **4**: 205-214.
12. Carlberg, I. And Mannervik, B. (1985). Glutathione reductase. Methods enzymol., **113**: 484-490.
13. El Denshary, E.S., Al-Gahazali, M.A., Mannaa, F.A., Salem, H.A., Hassan, N.S. and Abdel-Wahhab, M.A. (2011). Dietary honey and ginseng protect against carbon tetrachloride-induced hepatonephrotoxicity in rats. Exp. Toxicol. Pathol., **64**: 753-760.
14. Fiddler, R.N. (1977). Collaborative study of modified aoac method of analysis of nitrite in meat and meat products. J. Assoc. Off anal. Chem. (j. Aoac), **60**: 594-599.
15. Gu, H., Gu, X., Xu, Q. And Kang, W.-Y. (2014). Antioxidant activity *in vitro* and hepatoprotective effect of *phlomis maximowiczii* *in vivo*. Afr. J. Trad. Cam, **11**: 46-52.
16. Habig, W.H. And Jacoby, W.B. (1981). Assays for differentiation of glutathione-s-transferases. Methods enzymol., **77**: 398-405.
17. Halliwell, B. And Whiteman, M. (2004). Measuring reactive species and oxidative damage *in vivo* and in cell culture: how should you do it and what do the results mean? Brit. J. Pharmacol., **142**: 231-255.
18. Jaramillo-Juárez, F., Rodríguez-Vázquez, M.L., Rincón-Sánchez, A.R., Martínez, M.C., Ortiz, G.G., Llamas, J., Posadas, F.A. and Reyes, J.L. (2008). Acute renal failure induced by carbon tetrachloride in rats with hepatic cirrhosis. Ann. Hepatol., **7**: 331-338.
19. Kallai, L., Hahn, A., Röder, V. And Zupanec, V. (1964). Correlation between histological findings and serum transaminase values in chronic diseases of the liver. Acta med. Scand., **175**: 49-56.
20. Kim, S.H., Cheon, H.J., Yun, N., Oh, S.T., Shin, E., Shim, K.S. And Lee, S.M. (2009). Protective effect of a mixture of *aloe vera* and *silybum marianum* against carbon tetrachloride-induced acute hepatotoxicity and liver fibrosis. J. Pharmacol. Sci., **109**: 119-127.
21. Lawrence, R.A. and Burk, R.F. (1976). Glutathione peroxidase activity in selenium-deficient rat liver. Biochem. Biophys. Res. Commun., **71**: 952-958.
22. Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with the folin phenol reagent. J. Biol. Chem., **193**: 265-275.
23. Mak, D.H., Ip, S.P., Li, P.C., Poon, M.K. and Ko, K.M. (1996). Alterations in tissue glutathione antioxidant system in streptozotocin-induced diabetic rats. Mol. Cell biochem., **162**: 153-158.
24. Mofrad, P., Contos, M.J., Haque, M., Sargeant, C., Fisher, R.A., Luketic, V.A., Sterling, R.K., Shiffman, M.I., Stravitz, R.T. and Sanyal, A.J. (2003). Clinical and histologic spectrum of nonalcoholic fatty liver disease associated with normal alt values. Hepatology, **37**: 1286-1292.
25. Mukai, T., Mera, K., Nishida, K., Nakashima, M., Sasaki, H., Sakaeda, T. and Nakamura, J. (2002). A novel method for preparation of animal models of liver damage: liver targeting of carbon tetrachloride in rats. Biol. Pharm. Bull., **25**: 1494-1497.
26. Ozsoy, N., Can, A., Yanardag, R. and Akev, N. (2008). Antioxidant activity of *smilax excelsa* l. Leaf extracts. Food chem., **110**: 571-583.
27. Özsoy, N., Okyar, A., Arda-Pirinççi, P., Can, A., Bolkent, Ş and Akev, N. (2013). Evaluation of *Smilax excelsa* L. Use in experimentally induced nephrotoxicity. Kafkas univ. Vet. Fak. Derg., **19**: 807-814.
28. Reznick, A.Z. and Packer, L. (1994). Oxidative damage to proteins: spectrophotometric method for carbonyl assay. Methods enzymol., **233**: 357-363.
29. Singbartl, K., Green, S.A and Ley, K. (2000). Blocking p-selectin protects from ischemia/reperfusion induced acute renal failure. Faseb j., **14**: 48-54.
30. Yuan, L., Gu, X., Yin, Z. and Kang, W. (2014). Antioxidant activities *in vitro* and hepatoprotective effects of *nelumbo nucifera* leaves *in vivo*. Afr. J. Trad. Cam, **11**: 85-91.
31. Walter, K and Schütt, C. (1974). Acid and alkaline phosphatase in serum. In, bergmeyer hu (ed): methods of enzymatic analysis. 2nd ed. Vol 2, 856-864, academic press inc., newyork.