



Research Paper

Afr. J. Traditional,  
Complementary and  
Alternative Medicines  
www.africanethnomedicines.net

ISSN 0189-6016©2007

ANTIMALARIAL ACTIVITY IN CRUDE EXTRACTS OF SOME CAMEROONIAN MEDICINAL PLANTS.

Jean Bickii<sup>a</sup>, Guy Raymond Feuya Tchouya<sup>b</sup>, Jean Claude Tchouankeu<sup>b</sup> and Etienne Tsamo<sup>b</sup>.

<sup>a</sup>Institute for Medical Research and the studies of Medicinal Plants (IMPM), Laboratoire de Recherches sur le Paludisme, Yaoundé, Cameroon., <sup>b</sup>Faculty of Science, University of Yaoundé 1, P. O. Box 812 Yaoundé, Cameroon

\*E-mail: [jbickii@yahoo.fr](mailto:jbickii@yahoo.fr), Tel: +237 989 20 30

### Abstract

Fifteen crude extracts from the stem bark and seeds of four medicinal plants, viz: *Entandrophragma angolense*, *Picralima nitida*, *Schumanniohyton magnificum* and *Thomandersia hensii* were tested *in vitro* for their antimalarial activity against the chloroquine-resistant *Plasmodium falciparum* W2 strain. The results showed that the extracts of these plants possessed some antimalarial activity, the methanol extract of *Picralima nitida* demonstrating the highest activity *in vitro*. Further isolation and identification of some active compounds from these plants will justify their common use in traditional medicine for the treatment of malaria or fever in Cameroon.

**Key words:** *Entandrophragma angolense*; *Picralima nitida*; *Schumanniohyton magnificum*; *Thomandersia hensii*; *Plasmodium falciparum*; Antimalarial activity.

### Introduction

The spread of chloroquine-resistant *Plasmodium falciparum* and the alarming emergence of multi-drug resistant strains have raised an urgent need to search for new, safe and effective antimalarial drugs (O'Neill et al., 1986; Weenen et al., 1990). So, screening of herbal potions used in traditional medicine was indicated. Medicinal plant extracts as well as their occurring products have been used in the treatment of fever or other symptoms associated with malaria throughout the tropics and subtropics, with a folklore encompassing centuries of practical experience (Weenen et al., 1990). The continuing clinical use of quinine isolated from *Cinchona* species (Rubiaceae) and the recent discovery of a highly effective anti-malarial substance, artemisinin, from a Chinese herb, *Artemisia annua* L. (Compositae) have stimulated the interest in medicinal plants as sources of new potential antimalarial drugs.

Since 1980, large-scale *in vitro* and *in vivo* screenings of plants all over the world have been intensified and some of these have shown high antimalarial activities (Fandeur et al., 1985; Noster and Kraus, 1990; Koumaglo et al., 1991; Nkunya, 1992; Gessler et al., 1993).

In Cameroon, many plants are traditionally used as antimalarials or antipyretics (Tantchou et al., 1986; Kouamouo, 1988). A recent study using a crude extract and purified compounds of *Khaya grandifoliola* (Meliaceae) (Bickii et al., 2000) has yielded interesting results and stimulated us to investigate more plants from other families, traditionally used for the treatment of malaria or fever. *Entandrophragma angolense* (Welw) CDC (Meliaceae), *Picralima nitida* Stapf, 1910 (Apocynaceae), *Thomandersia hensii* De Wild and Th. Dur (Acanthaceae), and *Schumanniohyton* (Rubiaceae) (Letouzey, 1985) are some of these plants usually used in Cameroonian traditional medicine as antimalarials or antipyretics. Although, their therapeutic properties are claimed or evidenced in traditional medicine, few studies have been realised in our country to assess the activities of these medicinal plants.

In the present study, the *in vitro* antimalarial activity of these traditional medicinal plants, used in Cameroon, was assessed against *P. falciparum* W2 (Indochina I/CDC) chloroquine-resistant strain.

## Materials and methods

### Plant materials

Plants were selected on the basis of their ethnomedical studies and endemicity. Stem bark of *Entandrophragma angolense* Welwitsch C. D. C. (Meliaceae) was harvested in April 1999 in Yokadouma in the eastern Cameroon. Stem bark of *Thomandersia hensii* De Wild and Th. Dur (Acanthaceae) was harvested in march 2003 in Ngomedzap, southern Cameroon. Stem bark of *Schumanniophyton magnificum* Harms (Rubiaceae) and the seeds of *Picralima nitida* Stapf (Apocynaceae) were collected in April 2003 in Eseka, southern Cameroon. Plants were authenticated in the Cameroon National Herbarium, Yaoundé for *E. angolense* (N° 29932) and in the Herbarium of the Research Centre for Medicinal Plants and Traditional Medicine, Laboratory of Botany, Yaoundé, for *P. nitida* (LB Pn 0301), *T. hensii* (LB Th 0301) and *S. magnificum* (LB Sm 0301), where different Voucher specimens were conserved.

### Extraction procedures

Plant materials were dried and pulverised. The following extracts were made: hexane, 50% dichloromethane-methanol, methanol and water-ethanol (1: 1 v/v) of the stem bark of *E. angolense*; hexane, 50% dichloromethane-methanol and methanol extracts of the seeds powder of *P. nitida*; hexane, ethyl acetate, 50% dichloromethane-methanol and methanol extracts of the stem bark powder of *T. hensii* and hexane, 50% dichloromethane-methanol, methanol and water-ethanol (1: 1) extracts of the stem bark powder of *S. magnificum*.

Dry and powdered stem bark of *E. angolense*, *S. magnificum*, *T. hensii* and the seeds of *P. nitida*, were extracted separately with a mixture of water-ethanol (1: 1) (*E. angolense* and *S. magnificum*) and hexane. The defatted material obtained after hexane extraction was then extracted successively with ethyl acetate, dichloromethane – methanol (1:1), and methanol.

Hence, dried and ground plant material (3.34 kg *E. angolense*, 2.5 kg *S. magnificum*) were macerated with ethanol-water (1:1) mixture for 48h and 96h. The extracts were filtered and the filtrate were freeze-dried to yield (2.62 % w/w *E. angolense*, 3.28 % w/w *S. magnificum*) respectively.

For the hexane, ethyl acetate, dichloromethane-methanol (1:1) and methanol extracts, the ground materials (8 kg of each plant) were dried and successively extracted with hexane, ethyl acetate, dichloromethane-methanol (1:1) and methanol at room temperature for 48h and 96h. The extracts were filtered and the filtrates were submitted to concentration by a rotavapor. The solvents were evaporated *in vacuo* to yield hexane extracts (2.85 % w/w *E. angolense*, 1.14 % w/w *P. nitida*, 0.30 % w/w *S. magnificum*, 0.67 % w/w *T. hensii*), ethyl acetate extract (6.7 % w/w *T. hensii*), dichloromethane – methanol extracts (4.28 % w/w *E. angolense*, 5 % w/w *P. nitida*, 1.97 % w/w *S. magnificum*, 6.2 % w/w *T. hensii*) and methanol extracts (3.57 % w/w *E. angolense*, 3.57 % w/w *P. nitida*, 1 % w/w *S. magnificum*, 5.83 % w/w *T. hensii*).

### *In vitro* evaluation assay

#### Parasites cultivation

The *P. falciparum* chloroquine-resistant strain W2 was maintained in continuous culture in sealed flasks according to the technique described by Trager and Jensen (1976). The clone W2 was cultivated in the culture medium (RMPI 1640, 25 nM Hepes and 25 nM NaHCO<sub>3</sub>, supplemented with heat inactivated 10 % human serum) and the parasites were incubated at 37 °C in a 3 % O<sub>2</sub>, 5 % CO<sub>2</sub> and 91 % N<sub>2</sub> atmosphere. These were synchronised in the ring stage by serial treatment with 5 % sorbitol (Sigma) (Lambros and Vanderberg, 1979) and studied at 1 % parasitemia.

#### Preparation of tested extracts

Extracts were prepared as 20 mg/ml stock solutions in DMSO (dimethyl sulfoxide, Sigma Chemicals Co, St Louis, MO, USA) filtered, diluted as needed for individual experiments and tested in triplicate. The stock solutions were diluted in supplemented RPMI 1640 medium so as to have at most 0.2 % DMSO in the final reaction medium. Seven concentrations were tested: 100, 50, 25, 12.5, 6.25, 3.125 and 1.56 µg/ml.

#### Flow cytometric analysis

An equal volume of 1 % parasitemia, 4 % hematocrit culture was thereafter added and gently mixed thoroughly. Negative controls contained equal concentrations of DMSO. Positive controls contained 1 µM

chloroquine diphosphate (Sigma chemicals Co, S<sup>l</sup> Louis, MO, USA). Culture were incubated at 37 °C for 48 hrs (1 parasite erythrocytic life circle). Parasite at ring stage were thereafter fixed by replacing the serum medium by an equal volume of 1 % formaldehyde in PBS (Sigma chemicals Co, S<sup>l</sup> Louis, MO, USA). Aliquots (50 µl) of each culture were added to 5 ml round- bottomed polystyrene tubes containing 0.5 ml 0.1 % Triton X- 100 and 1 nM YOYO nuclear dye (Molecular Probs) in PBS.

Parasitemias of treated and negative cultures were compared using Beckton Dickinson FACS<sup>ort</sup> Flow cytometer to count parasitized erythrocytes. Data acquisition was performed using Cell Quest software. These data were normalized to percent control activity and 50% inhibitory concentrations (IC<sub>50</sub>s), calculated using Prism 3.0 software (Graph Pad) with data fitted by non linear regression to the variable slope sigmoidal dose-response formula  $Y = 100 / [1 + 10^{-(\log IC_{50} - x)H}]$ , where H is the Hill coefficient or slope factor (Singh and Rosenthal, 2001). This activity was analysed in accordance with the norms of plants antimalarial activity of Rasoanaivo et al. (1998). According to these norms, an extract is very active if IC<sub>50</sub> < 5 µg/ml, active 5µg/ml < IC<sub>50</sub> < 50 µg/ml, weakly active 50 µg/ml < IC<sub>50</sub> < 100 µg/ml and inactive IC<sub>50</sub> > 100 µg/ml.

## Results

The IC<sub>50</sub> value of chloroquine for the clone W2 was 133 nM. The IC<sub>50</sub> values of the plants extracts are shown in Table 1. These extracts demonstrated that they possess antimalarial activity. No significant differences were observed between the activity of methanol and water-ethanol extracts of *E. angolense* and *S. magnificentum*. The methanol extract of *P. nitida* was more active than those of the other plants. However, nearly all the extracts showed antimalarial activity *in vitro*.

**Table 1:** IC<sub>50</sub> values of plants extracts against *Plasmodium falciparum* chloroquine-resistant strain W2.

Plants	Extract	Extract Yield (% w/w)	IC <sub>50</sub> (µg/ml)* (Mean ± SD)
<i>Entandrophragma angolense</i>	Hexanic extract	2.85	33.4 ± 0.5
	50% dichloromethane/MeOH	4.28	18.4 ± 1.8
	Methanolic extract	3.57	26.2 ± 3.2
	50 % Water-EtOH extract	2.62	26.4 ± 2.8
<i>Picralima nitida</i>	Hexanic extract	1.14	129.6 ± 1.7
	50% dichloromethane/MeOH	5.00	20.1 ± 2.4
	Methanolic extract	3.57	10.9 ± 1.1
<i>Thomandersia hensii</i>	Hexanic extract	0.67	53.9 ± 3.2
	Ethyl acetate extract	6.70	24.7 ± 0.8
	50% dichloromethane/MeOH	6.20	77.2 ± 1.5
	Methanolic extract	5.83	68.2 ± 1.4
<i>Schumanniphyton magnificentum</i>	Hexanic extract	0.30	78.1 ± 1.5
	50% dichloromethane/MeOH	1.97	47.8 ± 1.3
	Methanolic extract	1.00	28.7 ± 1.9
	50 % Water-EtOH extract	3.28	25.5 ± 2.0

(\*) The results are expressed as geometric means ± SD from triplicate experiments.

## Discussion

*Entandrophragma angolense* is locally used in the treatment of malaria and some other diseases (Irvine, 1961; Obih et al., 1986; Bray et al., 1990). The stem bark, in aqueous decoctions, is used for the treatment of fever or malaria in Cameroon and Ivory Coast. The IC<sub>50</sub> of the 50% dichloromethane-methanol extract of *E. angolense* was more active than the methanol and the water-ethanol extracts. This higher activity could be explained by the fact that some extracting compounds show relative affinity with some solvents. Although there was no significant difference between the IC<sub>50</sub> values of the methanol and the water-ethanol extract of *E. angolense*, our result justifies the use of palm wine, instead of water in the preparation of traditional remedies.

The stem bark, the roots, the seeds and the fruits of *Picralima nitida* are always used in our traditional medicine to cure malaria or fever. The stem bark, in aqueous decoctions is also used for the treatment of

diarrhoea, intestinal worms, and gonorrhoea in many Central and West African countries, and for the antimalarial and their anti-inflammatory activities (Ezeamuzie et al., 1994; Fakeye et al., 2000). The methanol extract of the seeds of *P. nitida* showed the highest antimalarial activity ( $IC_{50} = 10 \mu\text{g/ml}$ ), although this  $IC_{50}$  value is very low compared to those reported in other studies (François et al., 1996).

*Thomandersia hensii* is extensively used in traditional medicine (Burkill, 1985). A leaf decoction is used in Cameroon and Zaïre as a remedy of diarrhoea and colitis. The irritant sap is the treatment of choice for furuncles, abscesses and syphilis ulcers. A decoction of roots and leafy twigs is used for uro-genital disorders and intestinal parasites and as a tonic in case of debility and tiredness. Pulped roots are used for the treatment of oedema, rheumatism and sometimes ear and eye inflammations (Ngadjui et al., 1994). Although the water-ethanol extract was not tested, the antimalarial testing of the ethyl acetate extract showed some activity ( $IC_{50} = 24.7 \mu\text{g/ml}$ ) which is similar to the results obtained from the methanol and the water-ethanol extracts of *E. angolense* in our study.

*Shumanniophyton magnificum* is a plant widely used in Africa ethno-medicine for the treatment of various diseases, particularly malaria and fever. A decoction of the bark is in great repute among some tribes of Cameroon as a remedy for dysentery and used as an enema (Tane et al., 1990), while other tribes use it as a lotion after circumcision (Irvine, 1961). The juice is used as a snake bite remedy in Nigeria and the protective effects of the extract against snake venom have been demonstrated (Akunyili and Akubue, 1986; Houghton et al., 1992). This plant is also well-known as a repellent for snakes (Okogun et al., 1983). The methanol and the water-ethanol extracts were active against *P. falciparum* W2 strain. Their activities were similar to those of the same extracts of *E. angolense*.

### Conclusion

The results of our study show that the extracts of *E. angolense*, *P. nitida*, *T. hensii* and *S. magnificum* possess some antimalarial activity. The methanol and the water-ethanol extracts showed the highest activities, thus justifying the use of water and ethanol (palm-wine, spirits) as common solvents in traditional medicine. Further pharmacological screening with chemical fractions of methanol and water-ethanol extracts of *E. angolense*, *P. nitida* and *S. magnificum* and the ethyl acetate extract of *T. hensii* would permit the isolation and the identification of active compounds of these plants.

### Acknowledgements

This study was financed by the World Health Organisation/UNDP/WORLD BANK/WHO Special Programme for Research and Training in Tropical Diseases (TDR) through Research grant N° 990899 (Dr J. BICKII). The authors thank Dr Fabrice Fekam Boyom and his collaborators at the University of California, San Francisco.

### References

1. Akunyili, D. N., Akubue, P. I. (1986). Schumannioside, the anti snake venom principle from the stem bark of *Schumanniophyton magnificum* Harms. J. Ethnopharmacol., 18: 67-72.
2. Bickii, j., Njifutie, N., Ayafor, J. F., Basco, L. K. and Ringwald, P. (2000). *In vitro* antimalarial activity of limonoids from *khaya grandifoliola* C.D.C. (Meliaceae). J. Ethnopharmacol., 69: 27-33.
3. Bray, D. H., Warhurst, D. C., Connolly, J. D., O'Neill, M. J. and Phillipson, J. D. (1990). Plants as sources of antimalarial drugs. Part 7. Activity of some species of Meliaceae and their constituent limonoids. Phytother. Res., 4: 29-35.
4. Burkill, H. M. (1985). The useful plants of West Tropical Africa. Royal Botanic Garden, Kew, UK. Vol. 1, 2<sup>nd</sup> Edn., p. 28.
5. Ezeamuzie, I. C., Ojinnake, M. C., Uzygna, E. O., OJI, S. E. (1994). Antiinflammatory, antipyretic and antimalarial activity of a West African medicinal plant-*Picralima nitida*. Afr. J. Med. Sci., 23 (1): 85-
6. Fandeur, T., Moretti, C., Polonsky, J. (1985). *In vitro* and *in vivo* assessment of the antimamalrial activity of sergeolide. Planta med., 61: 20-23.
7. Fakeye, T. O., Itiola, O. A., Odelila, H. A. (2000). Evaluation of the antimicrobial property of the stem bark of *Picralima nitida* (Apocynaceae). Phytother. Res., 14: 368-370.
8. François, G., Ake Assi, L., Holenz, J., Brungmann, G. (1996). Constituents of *Picralima nitida* display inhibitory activities against asexual erythrocytes forms of *Plasmodium falciparum* *in vitro*. J. Ethnopharmacol., 54 (2-3): 113-117.

9. Gessler, M. C., Nkunya, M. H. H., Mwasumbi, L. B., Henrich, M., Tanner, M. (1993). Screening Tanzanian plants for antimalarial activity. *Acta Tropica*, 56: 65-77.
10. Houghton, P. J., Osibogun, I. M., Sukhi Bansal. (1992). A peptide from *Schumanniphyton magnificum* with anti-cobra venom activity. *Planta med.*, 58: 263-265.
11. Irvine, F. R. (1961). *Woody plants of Ghana*. Oxford University Press, Oxford, p 517.
12. Kouamouo, J. (1988). Utilisation des métabolites cellulaires comme marqueurs de l'activité pharmacologique : application aux drogues anti-cancéreuses et anti-malariques. Thèse Doctorat d'Etat. Université de Reims, France.
13. Koumaglo, K., Gbeassor, M., Nikabu, O., De Souza, C., Wernsdorfer, W. (1991). Effects of three compounds from *Morinda lucida* on *Plasmodium falciparum*. *Planta med.*, 58: 533-534.
14. Lambros, C., Vanderberg, J. P. (1979). Synchronisation of *Plasmodium falciparum* erythrocytic stages in culture. *J. Parasitol.*, 65: 418-420.
15. Letouzey, R. (1985). Carte phytogéographique du Cameroun au 1/500000°. Institut de la Carte Internationale de la Végétation, Toulouse, France, p. 5.
16. Ngadjui, B. T., Dongo, E., Ayafor, J. F., Connolly, J. D. (1994). Thomandertiol, a tetraterpenoid from the twigs of *Thomandersia laurifolia*. *J. Nat. Prod.*, 57 (1): 161-163.
17. Nkunya, M. H. H. (1992). Progress in the search of antimalarials. Napreca Monograph Series, N° 4. Addis-Ababa. Ethiopia.
18. Noster, S., Kraus, L. J. (1990). *In vitro* antimalarial activity of *Contera latiflora* and *Exoxteme caribaeum* extracts on *Plasmodium falciparum*. *Planta med.*, 56: 63-65.
19. Obih, P. O., Makinde, J. M. and Laoje, J. (1986). Investigation of various extracts of *Morinda lucida* for antimalarial actions on *Plasmodium berghei* in mice. *Afr. J. Med. Sci.*, 14: 45-49.
20. Okogun, J. I., Adeboye, J. O., Okorie, D. A. (1983). Novel structure of two chromone alkaloids from the root-bark of *Schumanniphyton magnificum*. *Planta med.*, 49: 95-99.
21. O'Neill, M. J., Bray, D. H., Boardman, P., Phillipson, D. J., Warhurst, D. C., Peters, W., Suffness, M. (1986). Plants as sources of antimalarial drugs: *in vitro* antimalarial activities of some quassinoids. *Antimicrob. Agents Chemother.*, 30: 101-104.
22. Rasoanaivo, P., Oketch-Rabah, H. (1998). Preclinical considerations on antimalarial phytomedicines. Part II. Efficacy evaluation. *Inst. Malgache de Recherches Appliquées*, p 1-13.
23. Singh A., Rosenthal P. J. (2001). Comparison of efficacies of cysteine protease inhibitors against five strains of *Plasmodium falciparum*. *Antimicrob. Agents Chemother.*, 45: 949-951.
24. Tane, P., Ayafor, J. F., Sodengam B. L., Connolly, J. D. (1990). Chromone glycosides from *Schumanniphyton magnificum*. *Phytochemistry*, 29 (3): 1004-1007.
25. Tanchou, J., Titanji, V. P. K., Divo, A., Jensen, B. (1986). Studies of Cameroonian medicinal plants. II. Antimalarial activities of the extract of *Alstonia bonei* and *Guiboursia tessamanii* on Viet-Nam strain of *Plasmodium falciparum*. *Rev. Sci. Tech.*, 3: 69-73.
26. Trager, W., Jensen, J. B. (1976). Human malaria parasites in continuous culture. *Science*, 193: 673-675.
27. Weenen, H., Nkunya, M. H. H., Bray, D. H., Mwasumbi, L. B., Kinabo, L. S. and Kilimali, V. A. E. B. (1990). Antimalarial activity of Tanzanian medicinal plants. *Planta Med.*, 56: 368-370.