



**Research Paper**

*Afr. J. Traditional,  
Complementary and  
Alternative Medicines*  
[www.africanethnomedicines.net](http://www.africanethnomedicines.net)

ISSN 0189-6016©2009

**NEW ANTITRYPANOSOMAL TETRANOTRITERPENOIDS FROM *AZADIRACHTA INDICA***

**\*Mercy Githua<sup>1</sup>, Ahmed Hassanali<sup>2</sup>, Joseph Keriko<sup>1</sup>, Grace Murilla<sup>3</sup>, Mary Ndungu<sup>1</sup> and Gathu Nyagah<sup>1</sup>**

<sup>1</sup>Department of Chemistry, Jomo Kenyatta University of Agriculture and Technology (JKUAT), P.O Box 62000-00200 Nairobi, Kenya. <sup>2</sup>International Center of Insect Physiology and Ecology (ICIPE), P.O Box 30772, Nairobi, Kenya, <sup>3</sup>Trypanosomiasis Research Center (TRC), P.O Box 362, Kikuyu, Kenya

\*E-mail: [githuamercy@yahoo.co.uk](mailto:githuamercy@yahoo.co.uk)

**Abstract**

Organic extracts of the leaves of *Azadirachta indica* A. Juss. yielded ten antitrypanosomal terpenoids. Three of these (**1 – 3**), are novel and are derivatives of nimbolide and nimbin. They were extracted from chloroform fraction of methanol extract. These compounds were found to exhibit strong antitrypanosomal activities against *Trypanosoma brucei rhodesiense* with MIC values ranging of 6.9, 15.6 and 7.8 µg/ml respectively and were more active than Cymerlarsan ( a standard drug), which had an MIC value of 187.5 µg/ml when tested against *T. b. rhodesiense*. The structures were elucidated by spectroscopic methods including; NMR, MS, UV and IR.

**Key words:** Meliaceae, limonoids, *Trypanosoma brucei rhodesiense*, *Azadirachta indica*, antitrypanosomal activity.

**Introduction**

*Azadirachta indica* (Meliaceae) is a large tree growing in tropical/subtropical regions. The leaves, bark, wood, roots and fruits are intensely bitter. Extracts from different parts of *A. indica* have shown activity against several diseases and their vectors that affect human and animals (Nagui, 1987; Mackinnon et al., 1997). According to Ayurveda (Hindu System of Medicine) the different parts of this tree possess different medicinal properties. A tea prepared from the leaves and bark is used to treat fever (Kokwaro, 1993); a decoction of fresh leaves is used as a favorite wash to cure old ulcers of long standing and the aqueous extract of the leaves in particular is used as remedy for malaria, similar to the practice in Nigeria. The methanol extract of *A. indica* exerts a pronounced anti-inflammatory (rat paw oedema) and a fairly good antipyretic effect (pyrogen induced hyperpyrexia) in rabbits, and thus may justify its use in the treatment of fever resulting from malaria (Okpanyi and Ezeukwu, 1981). The biological activities of *A. indica* have been widely attributed to the presence of limonoids; modified triterpenes with or derived from a precursor with a 4, 4,8-trimethyl-17-furanylsteroid skeleton. Crude extracts of some of the limonoids constituents of the *A. indica* have been shown to exhibit antimalarial activities against chloroquine resistant K1 strain of *Plasmodium falciparum* (Bray et al., 1990). We now report the structures of three new limonoids that were isolated from the leaves of *A. indica* and their activities against *T. b. rhodesiense*.

## Materials and Methods

### Plant materials

The leaves of *A. indica* were collected from Shimba Hills National Park, Kwale, South coast of Kenya and a voucher specimen No MG/1/2002 has been deposited in the herbarium of the Botany department of University of Nairobi (Kenya).

### Extraction and isolation of compounds from *A. indica*

Air dried and grounded powder of *A. Indica* leaves (6 Kg), was soaked in methanol (10 liters) for 48 hours and then filtered. The residue was soaked for another 48 hours in methanol (10liters) and then filtered. The combined filtrates were concentrated to dryness under reduced pressure at 45°C to yield a viscous oil (636 g). The extract was partitioned between water (100 ml) and chloroform (250 ml) x 3. The combined organic layer was evaporated to dryness under reduced pressure. Thirty grams of the chloroform layer was column chromatographed on silica gel (column size, 81 x 4.5 cm; 230 - 400 mesh) using hexane/ethyl acetate gradient (90 - 10% hexane). For each solvent system, 1 liter of solvent was used to elute and 50ml fractions collected. Separation was monitored using thin layer chromatography (TLC). The plates were developed in hexane: ethylacetate (2:1), then sprayed with Ehrlich's reagent (2% 4- dimethylaminobenzaldehyde) and developed in hydrogen chloride gas chamber. Typical purplish spots were obtained for limonoids rich fractions, which eluted at 55 - 100% ethylacetate (Maier and Edward, 1970).

A limonoid-bearing fraction (12 mg) was analyzed by semi-preparative HPLC on Beckmann Ultrasphere ODS reverse phase column (250 x 10 mm i.d) on Varian 5000 liquid chromatograph using 1:1 water/acetonitrile isocratic system at a flow rate of 2.5 ml per min and detected at 215 nm. Three new limonoids **1** (6.9 mg), **2** (15.6 mg) and **3** (7.4 mg), were obtained and tested for antitrypanosomal activity. NMR experiments on the compounds were recorded on Bruker Avance (500 MHz) Spectrophotometer. The <sup>1</sup>H and <sup>13</sup>C spectra were recorded in CDCl<sub>3</sub> solvent. Chemical shifts (δ) were reported in parts per million (ppm) relative to tetramethylsilane (TMS) as the internal standard and coupling constants (J), in hertz (Hz). Peaks on <sup>1</sup>H NMR are reported as singlets (s), doublets (d), doublet of doublet (dd), triplets (t) and multiplets (m). The <sup>13</sup>C multiplicity was determined by DEPT experiments, which gave chemical shifts values in ppm for assignments.

The IR spectra of the compounds were recorded on a Fourier Transform infrared spectrometer (FT-IR-8101, DR- 8001). Electron Impact Mass Spectra (EIMS) were obtained on a fission VG Platform II GC/LC-MS spectrometer at 70 eV by solid probe. Melting points of the compounds were determined on Sanyo Gallenkamp electronic melting point apparatus (Cal No. MPD350. BM 3.5 Serial No. SG97/12/425 UK and are uncorrected.

### Propagation of trypanosomes cultures

*Trypanosomiasis brucei rhodesiense* (KETRI 3438) procyclic forms were obtained from Kenya Trypanosomiasis Research Institute (KETRI) trypanosomiasis bank and stored in liquid nitrogen. They were resuscitated and cultured until correct parasitaemia for *in vitro* screening was obtained. The strain was isolated on 7<sup>th</sup> July 1997 from a human patient and passaged through mice. The pre-patent period was six days while the pre-freezing period was twenty three days.

### Antitrypanosomal bioassay

The procedure developed by Burri et al. (1993) to determine the levels of melarsoprol in serum samples of animals after drug treatment was employed. The bioassay was carried out in 96 flat-bottomed well plates with a lid (Corning Glass Works, Corning, New York) arranged in a matrix of 8 rows A - H and 12 columns 1 - 12 (Figure 1). The tests were carried out in columns 11 - 2 and rows B - G. The wells in columns 1 and 12 and rows A and H were not used due to high rate of evaporation. The stock solutions were prepared in sterilized distilled water (1 mg/ml) and diluted to appropriate concentrations. For every extract, three different concentrations were prepared, that is, 500, 250 and 125 µg/ml. Trypanosomes in logarithmic growth phase below 10<sup>6</sup>/ml one day after sub-passaging were diluted to 2 x 10<sup>3</sup> cells/ml in medium. Hundred µl of this trypanosome suspension was placed into each well. 100 µl extract solution of each concentration was added to two adjacent wells of column 11. Serial dilutions were carried out by transferring 100 µl from wells of column 11 to wells of column 10 by multi-channel pipettes, then from wells of column 10 to 9, from 9 to 8 and so on up to wells of column 4. For every test, the highest concentration was in column 11 and in twofold dilution to the lowest in column 4 in 128 fold range, so that, for every extract, the highest

concentration was 250  $\mu\text{g/ml}$  and the lowest was 0.488  $\mu\text{g/ml}$ . Columns 3 and 2 were the control, that is, no extract was added. Plates were incubated for 72 hrs at 27°C in 5%  $\text{CO}_2$ .

### Determination of minimum inhibition concentration

The test was observed under inverted microscope at 100-fold magnification. In every row the highest dilution of either standard or samples with less than 5 motile trypanosomes of normal shape was determined. The concentration in this well is defined as the minimum inhibitory concentration (MIC).

$$\text{MIC} \cdot \text{D} \cdot 2^n = \text{C}_x$$

MIC = Minimum Inhibitory Concentration

D = Dilution of extract by medium in wells of column 11 (maximum concentration)

n = Steps of dilution to the first well with no living trypanosomes

C<sub>x</sub> = Concentration of extract solution

\* = Multiplication

### Results and Discussion

The HREIMS of compound **1** gave a molecular ion at  $m/z$  554.2254 corresponding to  $\text{C}_{31}\text{H}_{38}\text{O}_9$ .  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Tables 1 and 2) showed resonances for four distinctive tertiary methyl groups  $\delta_{\text{H}}$  1.23 (H-19), 1.68 (H-18), 1.58 (H-29), 1.28 (H-30) on carbons at  $\delta_{\text{C}}$  16.4 (q),  $\delta_{\text{C}}$  12.8 (q),  $\delta_{\text{C}}$  17.1 (q),  $\delta_{\text{C}}$  17.5 (q), respectively. The positions of these methyl groups were assigned by analysis of HMBC spectrum. Correlations of H-18 with the carbon resonance at  $\delta_{\text{C}}$  134.9 (C-13) and  $\delta_{\text{C}}$  146.3 (C-14); H-19 with the carbon resonance at  $\delta_{\text{C}}$  202.6 (C-1),  $\delta_{\text{C}}$  43.6 (C-5),  $\delta_{\text{C}}$  41.4 (C-9) and  $\delta_{\text{C}}$  47.3 (C-10); H-29 with the carbon resonance at  $\delta_{\text{C}}$  148.5 (C-3),  $\delta_{\text{C}}$  47.7 (C-4),  $\delta_{\text{C}}$  43.6 (C-5) and  $\delta_{\text{C}}$  175.9 (C-28) and H-30 with the carbon resonance at  $\delta_{\text{C}}$  86.9 (C-7),  $\delta_{\text{C}}$  49.6 (C-8) and  $\delta_{\text{C}}$  146.3 (C-14) (Figure 2).  $^1\text{H}$  NMR spectrum indicated presence of a  $\beta$ -substituted furan moiety at  $\delta_{\text{H}}$  7.23, 6.29 and 7.28. Examination of COSY spectra revealed that methine protons at  $\delta_{\text{H}}$  6.29 coupled with methine protons at  $\delta_{\text{H}}$  7.28, which allowed the assignment of the hydrogen atoms at  $\delta_{\text{H}}$  6.29 to C-22, 7.28 to C-23 and 7.23 to C-21. The carbonyl group on C-1 was deduced from the absence of H-1 resonance and was confirmed from HMBC correlation of H-2  $\delta_{\text{H}}$  7.08 (d,  $J = 9.7\text{Hz}$ ), H-3  $\delta_{\text{H}}$  6.44 (d,  $J = 10.1\text{Hz}$ ), H-5  $\delta_{\text{H}}$  3.43 (d,  $J = 11.6\text{Hz}$ ), H-9  $\delta_{\text{H}}$  2.67 (t,  $J = 11.0, 5.5\text{Hz}$ ) and methyl protons at C-19  $\delta_{\text{H}}$  1.23 (s) with the carbon resonance at  $\delta_{\text{C}}$  202.6. Cross peaks between methyl protons at  $\delta_{\text{H}}$  1.58 (H-29), to carbons at  $\delta_{\text{C}}$  148.5 (C-3),  $\delta_{\text{C}}$  47.7 (C-4),  $\delta_{\text{C}}$  43.6 (C-5) and  $\delta_{\text{C}}$  175.9 (C-28) confirmed the position of H-29 and also confirmed C-28 as a ketone. The position of methoxy groups was deduced from HMBC correlations of the methoxy protons on C-12 with the carbon resonance at  $\delta_{\text{C}}$  51.6 (OCH<sub>3</sub>-12) and of the methoxy protons on C-15 with the carbon resonance at  $\delta_{\text{C}}$  50.6 (OCH<sub>3</sub>-15). The structure is proposed to be 7 $\alpha$ -acetyl-15 $\beta$ -methoxy-29 methylene 7,15-deoxo nimbolide.


HREIMS of compound **2** gave a molecular ion at  $m/z$  556.1342 corresponding to  $\text{C}_{31}\text{H}_{40}\text{O}_9$ . IR spectrum indicated the presence of a hydroxyl ( $3446\text{ cm}^{-1}$ ), CH aliphatic stretch ( $2997\text{ cm}^{-1}$ ), C=O ( $1729\text{ cm}^{-1}$ ), olefinic double bond ( $1652\text{ cm}^{-1}$ ) and C-O ( $1232\text{-}1150\text{ cm}^{-1}$ ). NMR spectral data showed a basic structure similar to that of nimbolide (**4**) a known compound isolated from ethanoic extract of *A. indica* (Sudaratana et al., 1985).  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data (Tables 1 and 2), showed presence of six tertiary methyl groups  $\delta_{\text{H}}$  0.925 (s, H-19), 1.64 (s, H-18), 1.12 (s, H-29), 1.28 (s, H-30), 2.17 (s, H-1a) and 1.95 (s, H-1b). The positions of these methyl groups were assigned by analysis of HMQC and HMBC correlations (Figure 2). Examination of COSY spectrum revealed that methine protons at  $\delta_{\text{H}}$  2.58 (dd,  $J = 9.5, 7.1, 2.4\text{ Hz}$ , 1H) coupled with methylene protons at  $\delta_{\text{H}}$  2.03 (d,  $J = 16.1\text{Hz}$ , 2H). HMBC correlation of the proton at  $\delta_{\text{H}}$  2.03 with carbons at resonance  $\delta_{\text{C}}$  172.5 (carbonyl carbon) and 39.3 allowed the assignment of protons at  $\delta_{\text{H}}$  2.03 to C-11 and 2.58 to C-9. From HMBC correlations, cross peaks between methyl protons at  $\delta_{\text{H}}$  1.95 (s),  $\delta_{\text{H}}$  2.17 (s) with carbons at  $\delta_{\text{C}}$  158.3 and  $\delta_{\text{C}}$  115.4 and cross peaks between protons at  $\delta_{\text{H}}$  5.74 (s) with carbons at  $\delta_{\text{C}}$  164.7, 20.4 and 27.5 suggests the side chain to be tigloyl. HMBC correlation of methine proton at  $\delta_{\text{H}}$  3.61 (t,  $7.2\text{Hz}$ , 1H) with furan carbons allowed the assignment of that proton to C-17 ( $\delta_{\text{C}}$  49.3). The HMBC cross peaks between methyl protons at  $\delta_{\text{H}}$  (1.64, s, 3H) with carbons at  $\delta_{\text{C}}$  134.7, 146.7 and 49.3 permitted the assignment of quaternary carbons C-13 and C-14. There is a double bond between them as they have no HMQC correlation and no COSY coupling. From  $^1\text{H}$  and  $^{13}\text{C}$  NMR there is one methoxy group whose position was revealed by the HMBC correlation. The assignment of relative stereochemistry was based on the NOESY experiments. The H-9 $\alpha$  resonance showed a NOESY correlation with OH-3, H-5, H-15 and H-16 $\alpha$  confirming the  $\alpha$  orientation of H's on C-5, C-15 and hydroxyl group on C-3. H-16 $\beta$  resonance showed a NOESY correlation with 3H-30 $\beta$ , H-7, 3H-19 $\beta$  and 3H-29 $\beta$  whereas H-17 resonance showed a NOESY correlation with 3H-

19 $\beta$ , 3H-29 $\beta$  and H-16 $\beta$ . NOESY correlation of H-6 with H-7, H-3, 3H-29 $\beta$  and 3H-30 $\beta$  confirmed the  $\beta$ -orientation of H's on C-3, C-6, C-7 and C-17 (Figure 3). The compound is proposed to be 2-oxo-3-deacetyl salannin.


Compound **3** has a molecular formula C<sub>30</sub>H<sub>38</sub>O<sub>10</sub> (m/z 558). The IR spectrum indicated presence of a hydroxyl group (3427 cm<sup>-1</sup>), CH aliphatic stretch (2924 cm<sup>-1</sup>), C=O (1732 cm<sup>-1</sup>), olefinic double bonds (1680 cm<sup>-1</sup>) and C-O (1234 cm<sup>-1</sup>). <sup>1</sup>H and <sup>13</sup>C NMR spectral data (Tables 1 and 2 indicated the presence of a  $\beta$ -substituted furan moiety at  $\delta_H$  7.2 (H-21, s), 6.32 (H-22, s) and 7.31 (H-23, d, 9.9 Hz) corresponding to carbons at  $\delta_C$  138.9 (d), 110.3 (d) and 143.0 (d), respectively. The signals at  $\delta_C$  148.0 (d), 126.7 (d) and 202.1 (s) corresponding to protons at  $\delta_H$  6.41 (1H, d, J = 4.1 Hz) and 5.8 (1H, dd, J = 16.6, 9.7, 6.9 Hz) are typical of  $\alpha$ ,  $\beta$ -unsaturated ketone moiety in the A-ring. HMBC correlation of protons at  $\delta_H$  (5.86, dd, J = 16.6, 9.7, 6.9 Hz), 6.41 (d, J = 4.1 Hz), 3.39 (d, J = 12.6 Hz), 2.74, (t, J = 9.3, 3.8 Hz) and methyl protons at 1.20(s) with the carbon resonance at  $\delta_C$  202.1(s) allowed the assignment of  $\delta_H$  6.41 to C-3 and proton at  $\delta_H$  5.86 to C-2 and carbonyl to C-1. This was confirmed by the examination of COSY spectra which revealed that methine protons at  $\delta_H$  6.41 (1H, d, J = 4.1 Hz) coupled with methine protons at  $\delta_H$  5.86 (1H, dd, J = 16.6, 9.7, 6.9 Hz). <sup>1</sup>H and <sup>13</sup>C NMR indicate the presence of 2 methoxy groups attached to keto carbons. The compound is proposed to be 7 $\alpha$ -hydroxy-15 $\beta$ -hydroxy-7,15-deoxo nimbin

**Table 1:** <sup>1</sup>H NMR spectral data for compounds 1-3 [ $\delta_H$  ppm, (multi, JHz)]


H	1	2	3
1		4.99 (s)	
2	7.08(d, J = 9.7)		5.86 (dd, J = 16.6, 9.7, 6.9)
3	6.44(d, J=10.1)	3.83 (d, J = 9.4)	6.41 (d, J = 4.1)
5	3.43 (d, J = 11.6)	2.64 (d, J = 11.7)	3.39 (d, J = 12.6)
6	3.79 (d, J = 7.1)	3.97 (dd, J = 11.4, 9.1, 3.1)	4.01 (d, J = 3.2)
7	4.01 (d, J = 3.2)	4.15 (d, J = 3.1)	3.88 (m)
9	2.67 (t, J = 11.0, 5.5)	2.58 (dd, J = 9.5, 7.1, 2.4)	2.74 (t, J = 9.3, 3.8)
11	2.33 (dd, J = 16.2	2.03 (d, J = 16.1)	2.22 (d, J = 3.6)
11	3.27 (d, J = 16.1, 5.4, 0.7)	2.26 (m, J = 12.7, 6.4)	2.86 (dd, J = 13.5, 6.3, 5.5)
15	5.57 (t, J = 7.8, 6.6)	5.41(t, J = 14.2, 7.2))	5.54 (t, J = 14.4, 6.3)
16	2.22 (m)	2.13 (m)	2.03 (m)
16	2.89 (dd, J = 16.1, 5.4, 0.7)	2.23 (m)	2.18 (m)
17	2.07 (m)	3.61 (t, J = 7.2)	2.81 (d, J = 2.7)
18	1.68 (s)	1.64 (s)	1.67 (s)
19	1.23 (s)	0.92 (s)	1.20 (s)
21	7.23 (s)	7.30 (s)	7.2 (s)
22	6.29 (d, J = 12.2)	6.29 (s)	6.32 (s)
23	7.28(d, J = 9.9)	7.24 (s)	7.31 (d, J = 9.9)
28		4.09 (d, J = 8.3)	
29	1.58 (s)	1.12 (s)	1.58 (s)
30	1.28 (s)	1.28 (s)	1.28 (s)
3OH		5.82 (s)	
6CH <sub>3</sub>			1.32 (s)
7CH <sub>3</sub>	1.36 (s)		
12OCH <sub>3</sub>	3.56 (s)		3.64 (s)
15OCH <sub>3</sub>	3.68 (s)		
28a	4.23(s)		
1b		5.74 (s)	
12OCH <sub>3</sub>		3.17 (s)	
1a CH <sub>3</sub>		2.17 (s)	
1b CH <sub>3</sub>		1.95 (s)	
15OH			3.79 (s)



**Figure 1:** Chemical structures of compounds 1-4



**Figure 2:** Long range HMBC correlation elucidating methyl groups



**Figure 3:** NOESY correlation of compound 2

**Table 2:**  $^{13}\text{C}$  NMR spectral data for compounds 1-3 ( $\delta_{\text{C}}$  (ppm))

carbon	1	2	3
1	202.6 (s)	71.8 (d)	202.1 (s)
2	126.4 (d)		126.7(d)
3	148.5 (d)	71.0 (d)	148.0 (d)
4	47.7 (s)	44.2 (s)	47.7 (s)
5	43.6 (d)	38.8 (d)	43.6 (d)
6	66.2 (d)	72.4 (d)	87.3 (d)
7	86.9 (d)	85.7 (d)	66.1 (d)
8	49.6 (s)	48.9 (s)	47.4 (s)
9	41.4 (d)	39.3 (d)	39.0 (d)
10	47.3 (s)	40.4 (s)	47.4 (s)
11	32.5 (t)	39.4 (t)	34.3 (t)
11	32.5 (t)	30.4 (t)	34.3 (t)
12	173.6 (s)	172.5 (s)	173.6 (s)
13	134.9 (s)	134.7 (s)	134.8 (s)
14	146.3 (s)	146.7 (s)	146.7 (s)
15	87.7 (d)	87.7 (d)	86.8 (d)
16	34.2 (t)	41.0 (t)	41.4 (t)
16	34.2 (t)	41.0 (t)	41.4 (t)
17	41.9 (d)	49.3 (d)	51.1(d)
18	12.8 (q)	13.1 (q)	12.5 (s)
19	16.4 (q)	15.3 (q)	16.3 (s)
20	126.8 (s)	127.1 (s)	126.3 (s)
21	138.9 (d)	142.7 (d)	138.9 (d)
22	110.4 (d)	110.7 (d)	110.3 (d)
23	148.1 (d)	138.7 (d)	143.0 (d)
28	175.9 (s)	78.3 (t)	173.6(s)
29	17.1 (q)	19.7 (q)	17.1 (s)
30	17.5 (q)	16.9 (q)	17.5 (s)
7C=O	174.1 (s)		152.9 (s)
7CH <sub>3</sub>	20.5 (q)		
12C=O	173 (s)		
12OCH <sub>3</sub>	51.6 (q)	51.3 (q)	51.6 (q)
15OCH <sub>3</sub>	50.6 (q)		
28a	85.3 (d)		
1C=O		164.7 (s)	
1a		158.3 (s)	
1b		115.4 (d)	
1aCH <sub>3</sub>		20.4 (q)	
1bCH <sub>3</sub>		27.5 (q)	
6C=O			152.9 (s)
6Me			20.0 (s)

### Antitrypanosomal Activity

Crude methanolic extracts and chloroform fractions of *A. indica* leaves exhibited significant antitrypanosomal *in vitro* activities against *T. b. rhodesiense* procyclic forms (KETRI 3438) whose MIC values are  $51.2 \pm 0$  and  $4.4 \pm 0$   $\mu\text{g/ml}$ , respectively and exhibited higher activities than cymerlarsan (drug in the market) which had an MIC value of  $187.5$   $\mu\text{g/ml}$ . Chloroform extracts were more potent than the methanol extracts, indicating that the active compounds are of medium polarity.

Repeated chromatographic separations of the chloroform fraction yielded three new tetranotriterpenoids (**1 - 3**) that were tested against *T. b. rhodesiense* procyclic forms (KETRI 3438). The MIC values were  $6.9 \pm 0$ ,  $15.6 \pm 0$  and  $7.8 \pm 0$   $\mu\text{g/ml}$ , respectively. All the three compounds had lower activities than that of the chloroform fraction suggesting that there are other antitrypanosomal compounds present or possible synergistic blend effects. Based on

their spectroscopic data and comparison with published data, the structures of the compounds were readily elucidated and were found to have the nimbolide (**4**) basic structure (C-seco limonoids). Compound **2** which has an epoxy group between C<sub>7</sub> and C<sub>15</sub> has the lowest activity whereas compounds **1** and **3** with no epoxy group and with more methoxy groups exhibited higher activities. This suggests that antitrypanosomal activity can be enhanced by chemical modifications of the basic structure (nimbolide) and its derivatives to generate new effective antitrypanocidal drugs with novel modes of action.

### Acknowledgments

We are grateful to Mr. S. G. Mathenge and Mr. Mutiso both of University of Nairobi, Botany Department and Mr. D. Gitau for their assistance in collection and identification of plant materials. We acknowledge Dr. Tony Hopper of IACR- Rothamsted for performing the Nuclear Magnetic Resonance (NMR) experiments and his valuable advice and guidance on the interpretation of the data. The staff of tissue culture section of Trypanosomiasis Research Center (TRC) are acknowledged for their assistance in bioassays.

### References

1. Bray, A. D.; Warhust, D.; Connolly, J. D.; Neil, M. J. O.; Phillipson, J. D. (1990). Plants as sources of Antimalarial Drugs. Part 7. Activity of some species of Meliaceae plants and their constituent Limonoids, *Phytotherapy Research* **4**, 29-35.
2. Burri, C.; Baltz, T.; Giroud, C.; Doua, F.; Welker, H. A.; Brun, R. (1993). Pharmacokinetic properties of trypanocidal drug melarsoprol. *Chemother.* **39**, 225-234.
3. Kokwaro, J. O. (1993). Medicinal plants of East Africa 2<sup>nd</sup> Edn. Kenya Literature Bureau Nairobi, Kenya. p157-158.
4. Maier, V. P. and Edward, R. G. (1970). Specific thin-layer chromatography assay of limonin, a citrus bitter J. *Agric Food Chem.*, **18**, 250
5. Okpanyi, S. N. and Ezeukwu, G. C., (1981). Anti-inflammatory and antipyretic activities of *Azadirachta indica*. *Planta. Med.*, **41**: 34
6. Mackinon, S.; Durst, T.; Arnason, J. T. (1997). Antimalaria activity of tropical Meliaceae extracts and Gedunin derivatives. *J. Nat. Prod.* **60**, 336-341
7. Nagui, S. N. H. (1987). [Eds.] In: Natural pesticides from the Neem tree and other Tropical plants. Proceedings of the 3<sup>rd</sup> Neem conference (Nairobi, 1996) GTZ, Eschborn Germany, p315.
8. Sudaratana, R. ; Yodhathai, T .; Chavi, Y .; Yongyuth, Y. (1985). Nimbolide, a constituent of *Azadirachta indica* inhibits *Plasmodium Falciparum* in culture. *Southeast Asian J.Trop. Med. Pub. Hlth.*, **16** 1