




## African Journal of Urology

Official journal of the Pan African Urological Surgeon's Association  
web page of the journal

[www.ees.elsevier.com/afju](http://www.ees.elsevier.com/afju)  
[www.sciencedirect.com](http://www.sciencedirect.com)



### Endourology

Original article

# Ureterscopy for treatment of obstructing ureteral calculi in pregnant women: Single center experience



T.K. Fathelbab\*, A.M.A. Hamid, E.M. Galal

Department of Urology, Faculty of Medicine, Minia University, Egypt

Received 9 October 2014; received in revised form 6 December 2014; accepted 19 December 2014  
Available online 25 November 2015

#### KEYWORDS

Stones;  
Ureterscopy;  
Pregnancy

#### Abstract

**Objectives:** To evaluate our experiences with ureteroscopic treatment of ureteral calculi in pregnancy.  
**Patients and methods:** Between April 2006 and October 2013, 41 pregnant women with persistent renal colics and/or hematuria refractory to conservative measures were treated with ureteroscopy. The patients' mean age was 23 (range 19–37) years. Most of the patients (56.1%) presented in the 2nd trimester. Loin pain and colic were the most common presenting symptoms (90.2%). Twenty-seven patients (65.9%) had an obstruction on the right side. All patients underwent ureteroscopy under epidural anesthesia.  
**Results:** Ureterscopy revealed the presence of ureteric stones in 36 of the 41 cases. The stone size ranged from 5 to 16 (mean 8.9) mm. Distal ureteric stones were found in 29 patients. The pneumatic lithoclast was used for stone fragmentation in 22 of them (75.9%), while the stone was directly extracted in 4 patients. In 3 patients the stone migrated proximally and was not accessible any more. Proximal ureteric stones were detected in 7 of the remaining 12 cases. When attempting to manipulate these stones, they migrated more proximally and became unreachable. In the last 5 patients the entire ureter was free of stones; they only had edema at the ureteric orifice. A long lasting JJ stent was left until the end of the pregnancy in all cases with migrated inaccessible stones. In all patients successfully treated, a JJ stent with dangle extraction strings was left for two weeks. Minor urologic complications were encountered in the form of mild dysuria in 12 cases (29.2%) and mild hematuria in 5 cases (12.2%). All patients completed their pregnancy until full term without any serious obstetric complications requiring intervention.  
**Conclusion:** Ureterscopy is a safe and effective therapeutic option for the treatment of obstructing ureteral stones in pregnancy with stone-free and complication rates comparable to the non-pregnant population.

© 2015 Pan African Urological Surgeons' Association. Production and hosting by Elsevier B.V. All rights reserved.

\* Corresponding author.

E-mail addresses: [tarekhalaf86@yahoo.com](mailto:tarekhalaf86@yahoo.com) (T.K. Fathelbab), [amrabelhamid2002@yahoo.com](mailto:amrabelhamid2002@yahoo.com) (A.M.A. Hamid), [Galaluro71@yahoo.com](mailto:Galaluro71@yahoo.com) (E.M. Galal).  
Peer review under responsibility of Pan African Urological Surgeons' Association.

<http://dx.doi.org/10.1016/j.afju.2014.12.005>

1110-5704/© 2015 Pan African Urological Surgeons' Association. Production and hosting by Elsevier B.V. All rights reserved.

## Introduction

Urolithiasis is the most common cause (after urinary tract infection) of urinary tract-related abdominal pain in pregnant women with a reported prevalence of 1/200–1/2000 women [1–3]. It is assumed that physiological alterations associated with pregnancy, such as urine stasis and hypercalciuria [4] which may be attributed to an increased glomerular filtration rate and increased urinary citrate, magnesium and glycoprotein levels, have an inhibitory and protective effect against stone formation [5]. Nevertheless, there is no difference in the incidence of urolithiasis between pregnant and non-pregnant women.

Ureteral stones are more common than renal pelvic or calyceal stones. Both sides are equally affected and most of the stones are composed of calcium phosphate or calcium oxalate [6]. The majority of pregnant patients with urolithiasis present in the second or third trimester [7,8].

Symptomatic urolithiasis during pregnancy has been reported to be the most common cause of abdominal pain and hospital admission for non-obstetric reasons [9].

Treatment of symptomatic ureteral calculi during pregnancy is mainly conservative in the form of hydration, analgesics, antispasmodics, and antibiotics in cases with associated infection [2]. Failed conservative treatment necessitates definitive stone management, either by insertion of a double-J stent or a nephrostomy tube, or by ureteroscopic manipulation of the stone [10–15]. We herein report on our experiences with ureteroscopic treatment of ureteral calculi in pregnant patients.

## Patients and methods

This prospective study conducted at the Urology Department, Minia University, Egypt, between April 2006 and October 2013 included 78 pregnant patients presenting with renal colics with or without hematuria and/or lower urinary tract symptoms (LUTS).

At first, conservative management in the form of hydration, analgesics, antispasmodics and, if needed, antibiotics was attempted for a duration of two weeks. Thirty-seven out of 78 patients (47.5%) responded well to conservative treatment and were thus excluded from the study. The remaining 41 patients who showed no improvement after conservative therapy were subjected to ureteroscopic manipulation.

The patients' mean age was 23 (range 19–37) years. The demographic characteristics of the patients are summarized in Table 1. The symptoms were renal colics in 37 (90.2%) and hematuria in 23 (56.1%) patients.

Laboratory work-up included urine analysis and urine culture, when needed, blood urea, serum creatinine and a complete blood picture in all patients. On admission, the patients were subjected to sonographic examination for the diagnosis of urinary tract obstruction. Further examinations included resistive index (RI) measurements and Doppler ultrasound to differentiate between definitive ureteral obstruction due to calculi and physiologic hydronephrosis. Right-side obstruction was found in 27 (65.9%) and left-side obstruction in 14 patients (34.1%). Neither KUB nor computed tomography (CT) was used to avoid hazards of radiation exposure to the fetus.

**Table 1** Patient's demographic characters.

Characteristic	Value
NO	41
Age (YRS)	19–37
<i>Gestational age</i>	
1st trimester	4 (9.8%)
2nd trimester	23 (56.1%)
3rd trimester	14 (34.1%)
<i>Presentation</i>	
Renal colic	37 (90.2%)
hematuria	23 (56.1%)
<i>Stone side</i>	
RT. Ureter	27 (65.9%)
LT. Ureter	14 (34.1%)

All patients were subjected to a complete obstetric examination pre- and postoperatively. The study was approved by the hospital ethics committee, and the patients were informed about the nature of the procedure. All patients gave a written consent to the intervention.

They received 2nd generation cephalosporin preoperatively. Epidural anesthesia was used in all cases to avoid risks of general anesthesia.

Ureteroscopy (7.5–8 F) was performed without dilatation using only semirigid ureteroscopes (Karl Storz, Tuttlingen, Germany). Obstetric monitoring was carried out during the whole procedure. Cystoscopy was done to identify the ureteric orifice. The guidewire was advanced through the ureteroscope into the ureteric orifice. The ureteroscope followed the wire up to the site of the stone, then the wire was advanced under vision, past the stone to the kidney. The ureteroscope was then removed and reintroduced beside the wire until it reached the stone. The pneumatic lithoclast was used for stone disintegration, while a basket or grasper was used for stone extraction. After ureteroscopic manipulation, a ureteral stent was left in situ, and its proximal coil was checked by immediate renal ultrasound.

## Results

Ureteric stones were found in 36 of the 41 patients. The stone size ranged from 5 to 16 mm (mean 8.9 mm). Distal ureteric stones were found in 29 cases. Pneumatic lithotripsy was used for stone disintegration in 22 cases (75.9%), while the stones were extracted without fragmentation, either by grasper or basket, in 4 cases (13.8%). In 3 cases (10.3%) the stone migrated proximally and became inaccessible.

Proximal ureteric stones were detected in the 7 of the remaining 12 cases. When attempting to manipulate these stones, they migrated more proximally and became inaccessible. In the last 5 patients the entire ureter was free of stones; they only had edema at the ureteric orifice.

As at the time of the study we did not have flexible ureteroscopes or a laser lithotripter, a long lasting JJ stent (Percuflex™, Boston Scientific Corp., USA) was left until the end of pregnancy in all cases with inaccessible proximally migrated stones (10 cases). After successful ureteroscopic stone extraction and/or disintegration, a JJ stent was left for two weeks in all patients. The stent was then

**Table 2** Procedure details and outcome.

Variable	Number
<i>Anesthesia</i>	
Epidural	41 (100%)
Stone size	5–16 mm (mean 8.9 mm)
<i>Stone manipulation</i>	
Lithotripsy	22 (75.9%)
Direct extraction	4 (13.8%)
<i>Stent</i>	
Double J	41 (100%)
Success rate	89.7% (26 out of 29 pts)
<i>Complications</i>	
Dysuria	12 (29.2%)
Hematuria	5 (12.2%)

removed in the outpatient clinic by pulling its dangle extraction string.

All patients had a smooth postoperative course apart from mainly stent-related mild dysuria and urgency in 12 (29.2%) and hematuria in 5 patients (12.2%). The details and the outcome of the procedure are shown in [Table 2](#).

All patients completed their pregnancy until full term without any serious obstetric complications requiring intervention.

## Discussion

Urolithiasis is more common in multiparous women, especially during the mid and late pregnancy, with no difference in incidence compared to the non-pregnant population [16]. The majority of our cases presented in the second and third trimesters (56.1 and 34.1%, respectively). This is consistent with the results of Song et al. [17] and Rana et al. [14] who observed that most of their cases presented in mid and late pregnancy.

Treatment of symptomatic ureteral stones is controversial [11], but it is widely accepted that conservative measures should be tried first. The reported incidence of spontaneous passage of the stone following conservative measures ranges from 22.8 to 66.6% [11,18]. In our study, successful conservative treatment was observed in 37 out of 78 patients (47.5%).

Failed conservative measures mandate definitive treatment of ureteral stones aiming to prevent renal function deterioration, urosepsis secondary to obstruction and obstetric complications, as well as to minimize maternal discomfort [13].

Definitive treatment of symptomatic ureteral calculi traditionally consists of either placement of a percutaneous nephrostomy tube or ureteral stent insertion. Disadvantages of these methods include the fact that they are more or less temporary measures with an increased risk of infection or incrustation requiring regular exchange [19] and the reported risk of stent migration, especially when inserted during early pregnancy [20]. Considering these disadvantages on one hand and the recent technological advancement in ureteroscopic design (small and flexible), the constant development of stone fragmentation devices and increased experience and skills in ureteroscopic techniques on the other, ureteroscopy has gained significant importance as a definitive treatment of symptomatic calculi, also during pregnancy [14]. While ureteroscopy during pregnancy, especially

in late pregnancy, used to be considered difficult because of the anatomic distortion, various studies have proven that it is a technique which can be safely and effectively carried out throughout the whole duration of pregnancy [12,21–24]. Stone fragmentation and subsequent stone extraction can be done with lithotripter devices, graspers or dormia baskets [25,26]. However, when using a lithotripter, the device should have the ability to deliver its energy to a very localized area, causing no or only minimal collateral damage, and it must enable the surgeon to approach the stone using a semirigid or flexible ureteroscope [14].

In our study, pneumatic lithotripsy was used for stone fragmentation in 22 of 29 cases (75.9%), while direct retrieval of the stones was possible in 4 patients, without any major complications. This result is in accordance with Ishii et al. [27] who used pneumatic lithotripsy in most of their cases (62%) and accomplished stone retrieval either by basket or forceps (26%). The stone-free rate in our study was 89.7% (26/29 patients) which is similar to the stone-free rates reported by Bozkurt et al. (92.6%) [13] and by Johnson et al. (88%) [28]. On the other hand, Travassos et al. [29] reported a 100% stone-free rate which may be explained by the small number of patients included in his study (9 patients).

Complications of ureteroscopy during pregnancy are uncommon and consist of either urologic or obstetric complications. In our study, we encountered only minor urologic complications which were mainly stent-related and were comparable to those in the non-pregnant population. The complications seen in our study, like those reported by Semins et al. [12], could be classified as complications grade 1 according to the Clavien classification [30]. In contrast, in a recently published study, Ishii et al. [26] reported a relatively higher incidence of complications (16.9%) which they attributed to an increased use of pneumatic lithotripsy. Obstetric complications during ureteroscopy are rare [11]. In a meta-analysis of 14 studies including 108 patients, the authors reported only one case of premature uterine contractions [12]. Similarly, Hoscan et al. [11] in their study of 34 cases reported one case of pre-term uterine contractions (2.9%). In the current study we did not encounter any obstetric complications.

## Conclusion

Diagnosis and management of ureteral calculi in pregnant females is a challenge. Conservative measures should be tried first, but failed conservative measures necessitate definitive treatment. Ureteroscopy and intracorporeal lithotripsy are safe and reasonable options with success and complications rates comparable with the non-pregnant population.

## Consent

Consent from the patient/parents. A written consent was obtained from the patient in their files.

## Ethical committee approval

This study has gained approval from the local research ethics committee of faculty of Medicine Minia University.

## Conflict of interests

No conflict of interests.

**Source of funding**

None.

**Authors' contributions**

Tarek Khalaf Fathelbab: Writing the manuscript, data collection, publication.

Amr Mohamed Abdel Hamid: Idea of the study, revision of manuscript.

Ehab Mohamed Galal: Writing the manuscript.

**References**

- [1] Gorton E, Whitfield HN. Renal calculi in pregnancy. *Br J Urol* 1997;80(Suppl. 1):4–9.
- [2] Srirangam SJ, Hickerton B, Van Cleynenbreugel B. Management of urinary calculi in pregnancy: a review. *J Endourol* 2008;22(5):867–75.
- [3] Waymont R, Schwartz BF. Pregnancy and urolithiasis. *Emedicine* 2009. <http://emedicine.medscape.com/article/455830-overview>.
- [4] Coe FL, Parks JH, Lindheimer MD. Nephrolithiasis during pregnancy. *N Engl J Med* 1978;298:324–6.
- [5] Maikranz P, Lindheimer M, Coe FL. Nephrolithiasis in pregnancy. *Bailliere's Clin Obstet Gynaecol* 1994;8(2):375–86.
- [6] Ross AE, Handa S, Lingeman JE, Matlaga BR. Kidney stones during pregnancy: an investigation into stone composition. *Urol Res* 2008;36:99–102.
- [7] Stothers L, Lee LM. Renal colic in pregnancy. *J Urol* 1992;148(5):1383–7.
- [8] Pais Jr VM, Payton AL, LaGrange CA. Urolithiasis in pregnancy. *Urol Clin N Am* 2007;34(1):43–52.
- [9] Laing KA, Lam TB, McClinton S, Cohen NP, Traxer O, Somani BK. Outcomes of ureteroscopy for stone disease in pregnancy: results from a systematic review of the literature. *Urol Int* 2012;89(4):380–438.
- [10] Deters LA, Belanger G, Shah O, Pais VM. Ultrasound guided ureteroscopy in pregnancy. *Clin Nephrol* 2013;79(2):118–23.
- [11] Hoscan MB, Ekinci M, Tunçkıran A, Oksay T, Özorak A, Özkardeş H. Management of symptomatic ureteral calculi complicating pregnancy. *Urology* 2012;80(5):1011–4.
- [12] Semins MJ, Trock BJ, Matlaga BR. The safety of ureteroscopy during pregnancy: a systematic review and meta-analysis. *J Urol* 2009;181(1):139–43.
- [13] Bozkurt Y, Penbegül N, Soylemez H, Atar M, Sancaktutar AA, Yıldırım K, et al. The efficacy and safety of ureteroscopy for ureteral calculi in pregnancy: our experience in 32 patients. *Urol Res* 2012;40(5):531–5.
- [14] Rana AM, Aquil S, Khawaja AM. Semirigid ureteroscopy and pneumatic lithotripsy as definitive management of obstructive ureteral calculi during pregnancy. *Urology* 2009;73:964–7.
- [15] Shokeir AA, Mutabagani H. Rigid ureteroscopy in pregnant women. *Br J Urol* 1998;81(5):678–81.
- [16] Maikranz P, Coe FL, Parks J. Nephrolithiasis in pregnancy. *Am J Kidney Dis* 1987;9:354–8.
- [17] Song Y, Fei X, Song Y. Diagnosis and operative intervention for problematic ureteral calculi during pregnancy. *Int J Gynecol Obstet* 2013;121:115–8.
- [18] Isen K, Hatipoglu NK, Dedeoglu S, Atilgan I, Çaça FN, Hatipoglu N. Experience with the diagnosis and management of symptomatic ureteric stones during pregnancy. *Urology* 2012;79(3):508–12.
- [19] Tóth C, Tóth G, Varga A, Flaskó T, Salah MA. Percutaneous-nephrolithotomy in early pregnancy. *Int Urol Nephrol* 2005;37(1):1–3.
- [20] Kavoussi LR, Jackman SV, Bishoff JT. Re: Renal colic during pregnancy: a case for conservative treatment. *J Urol* 1998;160:837–8.
- [21] Ulvik NM, Bakke A, Høisaeter PA. Ureteroscopy in pregnancy. *J Urol* 1995;154(5)(7):1660–3.
- [22] Scarpa RM, De Lisa A, Usai E. Diagnosis and treatment of ureteral calculi during pregnancy with rigid ureteroscopes. *J Urol* 1996;155:875–7.
- [23] Watterson JD, Girvan AR, Beiko DT, Nott L, Wollin TA, Razvi H, et al. Ureteroscopy and holmium:YAG laser lithotripsy: an emerging definitive management strategy for symptomatic ureteral calculi in pregnancy. *Urology* 2002;60:383–7.
- [24] Tawfik ER. Ureteroscopy during pregnancy using follow-the-wire technique. *Urotoday Int J* 2009;2(6).
- [25] Lifshitz DA, Lingeman JE. Ureteroscopy as a first-line intervention for ureteral calculi in pregnancy. *J Endourol* 2002;16(1):19–22.
- [26] Akpinar H, Tufek I, Alici B, Kural AR. Ureteroscopy and holmium laser lithotripsy in pregnancy: stents must be used postoperatively. *J Endourol* 2006;20:107–10.
- [27] Ishii H, Aboumarzouk OM, Somani K. Current status of ureteroscopy for stone disease in pregnancy. *Urolithiasis* 2014;42:1–7.
- [28] Johnson EB, Krambeck AE, White WM, Hyams E, Beddies J, Marien T, et al. Obstetric complications of ureteroscopy during pregnancy. *J Urol* 2012;188(1):151–4.
- [29] Travassos M, Amselem I, Filho NS, Miguel M, Sakai A, Consolmagno H, et al. Ureteroscopy in pregnant women for ureteral stone. *J Endourol* 2009;23(3):405–7.
- [30] Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–13.