



Post-tonsillectomy hemorrhage after bipolar diathermy vs. cold dissection surgical techniques in Alahsa region, Saudi Arabia[☆]



Ibrahim K. Aljabr, Fathelrahman M. Hassan^{*}, Khalid A. Alyahya

College of Medicine, King Faisal University, Alahsa, Saudi Arabia

Received 12 May 2015; accepted 21 June 2015

Available online 17 August 2015

KEYWORDS

Post-tonsillectomy hemorrhage;
Bipolar diathermy;
Cold dissection;
Risk factors;
Saudi Arabia

Abstract *Introduction:* Tonsillectomy is a common surgical procedure done by otolaryngologists. Tonsillectomy is a relatively simple procedure. The concept of implementing it as a day case operation has become increasingly popular.

Material and Methods: This is a cross sectional study done in Alahsa city, eastern province, Saudi Arabia during the period from January 2014 to March 2015. This study reported the postoperative hemorrhage after Bipolar diathermy and Cold dissection surgical techniques to evaluate the incidence of the hemorrhage and to identify the possible risk factors associated with its occurrence.

Results: Postoperative bleeding occurred in 45 (3.6%) out of 1232 patients. Post-tonsillectomy hemorrhage according to operation technique was significantly higher among patients who underwent bipolar diathermy than cold dissection technique ($p < 0.05$).

Conclusion: Bleeding after operation by bipolar diathermy technique was occurring more frequently within the first five days. Hemoglobin level was significantly decreased in post-tonsillectomy hemorrhage.

© 2015 The Authors. Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Tonsillectomy dates back over 2000 years¹ and is now the most commonly performed surgical procedure in Pediatrics.² It is a common surgical procedure done by otolaryngologists.

^{*} Corresponding author. Tel.: +966 53029004.

E-mail addresses: fathmaga@yahoo.com, fgameel@kfu.edu.sa (F.M. Hassan).

[☆] **Financial support used for the study:** No financial support was used. Peer review under responsibility of Alexandria University Faculty of Medicine.

Tonsillectomy is a relatively simple procedure. The concept of implementing it as a day case operation has become increasingly popular.³ However, a number of postoperative complications have been well-documented in the literature; the most common and potentially life-threatening of which is post-tonsillectomy hemorrhage (PTH).⁴ Considering that it is an elective procedure, it is often connected with a comparatively high bleeding rate. Furthermore, bleeding in this location (the upper airways) always represents a significant risk.⁵ PTH occurs at a rate between 0.28% and 20%. This wide range may reflect the diversity in the otolaryngological community on how to properly define significant PTH.⁶ This study

reported the postoperative hemorrhage after Bipolar diathermy and Cold dissection surgical techniques at the Alahsa city, Saudi Arabia, to evaluate the incidence of this hemorrhage and to identify the possible risk factors associated with its occurrence.

2. Materials and methods

This is a cross sectional study done in Alahsa city, Saudi Arabia, during the period from January 2014 to March 2015, and the outcome of Pediatric tonsillectomy performed as day case procedures was studied. A total number of patients who underwent tonsillectomy were one thousand two hundred and thirty-two patients (928 of them by bipolar diathermy technique, 304 by cold dissection technique). This study included only forty-five patients coming back after tonsillectomy with bleeding. Indications for tonsillectomy among most of patients were recurrent tonsillitis (Throat infections), snoring or problems with sleep. Surgical procedures were performed using the techniques of bipolar diathermy or cold dissection. Postoperatively, they received the same medications (antibiotics and analgesics). Postoperative tonsillectomy hemorrhage incidents were identified. Data collected included patient's age, day of postoperative bleeding, operation duration, site of bleeding and the vital signs. To determine statistical significance, cross tabulation and chi-squared analysis were performed. Statistical significance was set at $P < 0.05$.

3. Results

Postoperative bleeding occurred in 45 (3.6%) out of 1232 patients. Comparisons for age, medication used, medication allergy, other illness, development and growth were done. The age of ten patients (22.2%) was on the range of 0–6 years, and the age of the rest 35 patients (77.8%) was > 6 years.

Table 1 Patients characteristics.

Characteristics	Frequency, <i>n</i>	%
<i>Ages (mean 8.3 years)</i>		
0–6 Years	10	22.2
> 6 Years	35	77.8
<i>Medication used</i>		
1 – Ventolin	9	20.0
2 – Flixonase	2	4.50
3 – Nasonex	2	4.50
4 – Others	32	70.1
<i>Medication allergy</i>		
1 – Penicillin	4	8.80
2 – Others	1	2.20
3 – None	40	89.0
<i>Other illness</i>		
1 – Asthma	9	20.0
2 – DM	2	4.40
3 – Allergic rhinitis	3	6.60
4 – Epilepsy	1	2.20
5 – None	30	66.8
<i>Development and growth delay</i>		
1 – Normal	43	95.5
2 – Abnormal	2	4.50

About 95.5% of patients with postoperative bleeding were having normal growth and development as shown in Table 1. Thirty-eight patients (84.4%) were suffering from recurrent tonsillitis (Throat infection), twenty-eight (62.2%) of them had snoring and 25 (55.6%) were suffering from sleeping difficulties (e.g. Apnea) (see Fig. 1). Post-tonsillectomy hemorrhage according to operation technique was significantly higher among patients who underwent the bipolar diathermy than cold dissection technique ($p < 0.05$). On the other hand, there were no statistically significant differences in age groups, other illness, vital sign and sides of bleeding according to patients' operation technique as shown in Table 2. The mean hemoglobin levels were significantly decreased among postoperation patients compared to preoperation ($p < 0.05$), while the hemoglobin variation between the two tonsillectomy techniques revealed a non-significant difference ($P > 0.05$), Table 2. The levels of prothrombin time (PT), Activated partial thromboplastin time (APTT) and international normalize ratio (INR) were statistically not significant ($(p = 0.166)$, ($p = 0.183$), ($p = 0.415$), respectively) in postoperation compared to preoperation, Table 3.

4. Discussion

Tonsillectomy-related morbidity and mortality are sources of potential malpractice claims in the field of otolaryngology. Tonsillectomy operation with hemorrhage is the major complication which, can be potentially life threatening. The previously reported clinical risk factors for post-tonsillectomy hemorrhage included age, sex, surgical technique and device, surgeon's skill level, and tonsillectomy indication.^{7–11} Our results agree that age, tonsillectomy indication, and operation technique are predictive of post-tonsillectomy hemorrhage. Several studies have described the patient's age as a significant risk factor.¹² Tomkinson et al.⁹ reported that patients older than 12 years had a 3-fold higher likelihood of severe post-tonsillectomy hemorrhage, and this finding was in agreement with our present study results which indicated that most of the patients with post-tonsillectomy hemorrhage were in older age group. This is probably related to the fact that older patients, who were the majority of the bleeder group in our study, have more time to get infected which leads to more fibrosis and more aggressive operation. Several studies have found an association between the indication for tonsillectomy and the post-tonsillectomy hemorrhage. In our study, the indication for tonsillectomy was mainly related to recurrent throat infection, snoring and sleep difficulties, as reported previously.^{8,9} No statistical association was found between operative time and post-tonsillectomy hemorrhage. In the present study, bleeding after operation was observed among bipolar diathermy patients (22 out of 45) within less than 5 days after operation compared to (2 out of 45) patients in cold dissection group. Our result was in agreement with the study done by Weimert et al. who performed a double blinded study to compare bipolar diathermy tonsillectomy and cold dissection. They found no difference in the incidence of post-tonsillectomy hemorrhage.¹³ Previous authors have noted that bipolar diathermy may cause increased severity of pain and may increase the risk of delayed hemorrhage.¹⁴ However, proponents of the technique state that it is a much faster procedure with minimal intraoperative blood loss and negligible incidence of immediate post-tonsillectomy




Figure 1 Indication for tonsillectomy.

Table 2 Post-tonsillectomy hemorrhage according to operation techniques.

	Bipolar diathermy	Cold dissection	P value
Frequency, n			
<i>Age groups/years</i>			
0-6	8	2	0.338
>6	29	6	
<i>Other illness</i>			
Asthma	7	2	0.641
D.M.	1	1	
Allergy	3	0	
Epilepsy	1	0	
Others	20	5	
<i>Sides of bleeding</i>			
Left side	19	5	0.810
Right side	16	5	
<i>Bleeding after operation</i>			
< 5 days	22	2	0.032
5-10 days	8	5	
> 10 days	8	0	
<i>Vital sign</i>			
With tachycardia	7	2	0.116
Without tachycardia	34	2	
<i>Hemoglobin variation</i>			
< 1.0 gm/dl	13	2	1.000
> 1.0 gm/dl	26	4	

P value < 0.05 is revealed significant, D.M.: diabetes mellitus.

hemorrhage.¹⁵ Most have found no difference between the two methods, but prefer diathermy because of the decreased operating time needed and drier field that was come to a similar conclusion.¹⁶ Our results were similar to previous studies that have not shown the chronic infection is a risk factor for

Table 3 The mean level of laboratory analysis in pre and postoperation patients.

Laboratory analysis	Mean ± SD		P value
	Pre-operation	Post-operation	
Hemoglobin (g/dl)	13.4 ± 1.3	12.3 ± 1.1	0.000
PT (s)	12.1 ± 1.5	12.5 ± 1.2	0.166
APTT (s)	28.8 ± 1.2	29.2 ± 1.6	0.183
INR	0.9 ± 1.0	1.1 ± 1.3	0.415

P value < 0.05 is revealed significant; PT, prothrombin time; APTT, activated partial thromboplastin time; INR, international normalize ratio.

post-tonsillectomy hemorrhage.¹⁷ The preoperative hematologic evaluation was also considered. The postoperative mean hemoglobin was significantly decreased ($P < 0.05$) compared to preoperative values, and most likely had an association with bleeding. There were no patients with abnormally elevated PT/PTT and INR. These patients were medically cleared pre-operatively. It is suggested that screening PT/PTT should be reserved for patients with known or suspected coagulopathies.¹⁸ For the vital signs, our data found no significant association between tachycardia after both operation techniques was in agreement with the data that show an increased incidence of bleeding in patients with normal heart rates in the study done by Cantor and Rogers.¹⁹ The greatest effect on hemoglobin level drops up to more than 1.0 gm/dl was seen in 26 patients with bipolar diathermy compared to cold dissection (4 patients). Our results found no statistical significance in hemoglobin concentration variation in both techniques ($P > 0.05$).

5. Conclusions

The post-tonsillectomy hemorrhage occurred in 45 patients (3.6%) out of 1232 patients underwent tonsillectomy in the

study period. Hemoglobin level was significantly decreased in post-tonsillectomy hemorrhage. Bleeding after operation by bipolar diathermy technique was common among patients within less than five days. Other illness, site of bleeding, method of tonsillectomy, PT, APTT, INR, total operative time and postoperative pulse rate were not significantly associated with post-tonsillectomy hemorrhage. The selection of the suitable operation technique and known patient's coagulopathies may help the otolaryngologist in identifying patients at risk for post-tonsillectomy hemorrhage.

Conflict of interest

We have no conflict of interest to declare.

References

1. Elsobky S, Ahmad N, Qureshi M, Izzath W, Sadiq H. Paediatric day case tonsillectomy: a safe, feasible and an economical way to treat patients – Yorkhill experience. *Scottish Med J* 2014;**59**(1):5–8.
2. Hern J, Waddell T. Day case paediatric tonsillectomy: a review of three years experience in a dedicated day case unit. *Clin Otolaryngol Allied Sci* 1999;**24**:208212.
3. Ibekwe TS, Obasikene G, Offiong E. Tonsillectomy: vasoconstrictive hydrolytic cold dissection method. *Afr J Paediatr Surg* 2013;**10**(2):150–3.
4. Opatowsky MJ, Browne JD, McGuirt Jr WF, et al. Endovascular treatment of hemorrhage after tonsillectomy in children. *AJNR Am J Neuroradiol* 2001;**22**(4):713–6.
5. Senska G, Schröder H, Pütter C, et al. Significantly reducing post-tonsillectomy haemorrhage requiring surgery by suturing the faucial pillars: a retrospective analysis. *PLoS One* 2012;**7**(10):e47874.
6. Liu JH, Anderson KE, Willging JP, et al. Post tonsillectomy hemorrhage. What is it and what should be recorded? *Arch Otolaryngol Head Neck Surg* 2001;**127**(10):1271–5.
7. Perkins JN, Liang C, Gao D, Shults L, Friedman NR. Risk of post-tonsillectomy hemorrhage by clinical diagnosis. *Laryngoscope* 2012;**122**:2311–5.
8. Kim MK, Lee JW, Kim MG, Ha SY, Lee JS, Yeo SG. Analysis of prognostic factors for postoperative bleeding after tonsillectomy. *Eur Arch Otorhinolaryngol* 2012;**269**:977–81.
9. Tomkinson A, Harrison W, Owens D, Harris S, McClure V, Temple M. Risk factors for postoperative hemorrhage following tonsillectomy. *Laryngoscope* 2011;**121**:279–88.
10. Ali RB, Smyth D, Kane R, Donnerly M. Post-tonsillectomy bleeding: a regional hospital experience. *Ir J Med Sci* 2008;**177**:297–301.
11. Tomkinson A, De Martin S, Gilchrist CR, Temple M. Instrumentation and patient characteristics that influence postoperative hemorrhage rates following tonsil and adenoid surgery. *Clin Otolaryngol* 2005;**30**:338–46.
12. Windfuhr JP, Chen YS, Remmert S. Hemorrhage following tonsillectomy and adenoidectomy in 15,218 patients. *Otolaryngol Head Neck Surg* 2005;**132**:281–6.
13. Weimert TA, Babyak JW, Richter HJ. Electrodissection tonsillectomy. *Arch Otolaryngol Head Neck Surg* 1990;**116**:1866188.
14. Roy A, Dela Rosa C, Vecchio YA. Bleeding following tonsillectomy – a study of electrocoagulation and ligation techniques. *Arch Otolaryngol Head Neck Surg* 1976;**102**:9–10.
15. Kennedy KS, Strom CG. A comparison of postoperative bleeding incidence between general and local anesthesia tonsillectomies. *Otolaryngol Head Neck Surg* 1990;**102**:654–7.
16. Papangelou L. Hemostasis in tonsillectomy – a comparison of electrocoagulation and ligation. *Arch Otolaryngol Head Neck Surg* 1972;**96**:3588360.
17. Kennedy KS, Strom CG. A comparison of postoperative bleeding incidence between general and local anesthesia tonsillectomies. *Otolaryngol Head Neck Surg* 1990;**102**:654–7.
18. Manning SC, Beste D, McBride T, Goldberg A. An assessment of preoperative coagulation screening for tonsillectomy and adenoidectomy. *Int J Pediatr Otorhinolaryngol* 1987;**13**:2377244.
19. Cantor RJ, Rogers J. Post tonsillectomy haemorrhage in children: the value of routine monitoring of the pulse. *J Laryngol Otol* 1984;**98**:993–5.