



Bayero Journal of Pure and Applied Sciences, 8(2): 117 – 122

Received: November, 2014

Accepted: October, 2015

ISSN 2006 - 6996

EVALUATION OF THE AFFINITY OF THE COMPONENTS OF BLOOD AND FUR OF ALBINO RAT FOR BIOACCUMULATION OF CHROMIUM (VI) ION FOLLOWING MILD EXPOSURE

Omoniyi K. I. * and Ekwumemgbo A. P.

Chemistry Department, Ahmadu Bello University, Zaria, Nigeria.

*Correspondence e-mail: israeliflourish@yahoo.com

*Telephone Number: +234-08036257789

Abstract

The study investigates the affinity of chromium (VI) ion for the components of blood and fur of female albino rats after spaced dosing with 5 mL and 1 mL of 1.0 µg/mL K₂Cr₂O₇ solution intraperitoneally in May, 2012; for application in forensic analyses following industrial and environmental exposures. The level of total-Cr ions was determined in the serum, red blood cell, plasma and fur of the experimental female albino rats after dosing with 5 mL of 1.0 µg/mL K₂Cr₂O₇ solution intraperitoneally in day 1, and 1 mL of the same solution on day 8 of the experiment, using atomic absorption spectrophotometry. After two weeks of K₂Cr₂O₇ administration, 12.127 ± 0.726 µg/mL, 2.956 ± 0.003, 2.210 ± 0.002 and 4.090 ± 0.0078 µg/mL were the mean levels of ions in the serum, red blood cell (RBC), plasma and fur respectively. There was no significant weight increase in the experimental group compared to the control group during the dosing period; however, the experimental group showed passiveness, consumed less feed, but had more water intake compared to the control group. The ranking for the mean level of total-Cr ions in the experimental rat followed the ranking, serum > fur > red blood cell (RBC) > plasma; while the control had the ranking serum > fur > plasma > RBC. There was high statistical increase in the ions in the serum of the experimental rat compared to the control; likewise the level in the experimental fur was highly elevated compared to the control (P < 0.05). The total-Cr ions in the fur and serum were doubled due to the administration of K₂Cr₂O₇ solution to the rats. The findings of the study signaled that the fur and serum can equally serve as specimen for detection of mild exposure to Cr (VI) during forensic investigation.

Key words: albino rat; blood component; chromium (VI) ion; forensic; mild exposure

1. INTRODUCTION

Kortenkamp (1997) surveyed the most important chromium-containing materials or objects as: Cr ore, lubricating oils, artificial dentures, wood preservation salts, cement, and leather tanned with Cr, dust particles beside busy roadways and use of tobacco products. People who work with materials containing mere traces of Cr salts are more at risk than workers who come into contact with high concentration of Cr salts (Adya *et al.*, 2005).

Chromium (VI) is the primarily form of Cr that has been implicated to be harmful to the biosphere (Amoikon *et al.*, 1995). The major emission source of Cr (VI) is the chrome plating industry. Chromic acid, used to electroplate metal parts, is the most common Cr (VI) compounds produced; chromic acid is also registered as a fungicide and pesticide in California for use in wood and lumber protection treatments (Kortenkamp, 1997; Becker *et al.*, 1991). Emissions of Cr (VI) may also result from lead chromate in paint, chromate production, coatings on stainless steel, fuel combustion and auto refinishing. If not all, most of this emitted

chromium is in the Cr (VI) state (Amoikon *et al.*, 1995; Becker *et al.*, 1991).

Cr (VI) exists as a component of a complex anion that varies with pH and may take the form of chromate (CrO₄²⁻), hydrochromate (HCrO₄⁻), or dichromate (Cr₂O₇²⁻). These ionic Cr⁺⁶ forms are highly soluble in water and thus mobile in the aquatic environment or biofluids (Ecological Analysts, 1981).

Accumulations of Cr in tissues and organism depend heavily on its chemical form, route of entry and amount administered. On exposure of the body to Cr (VI), the body changes Cr (VI) to Cr (III) (Yamaguchi and SanoK, 1983). Cr (III) in biological materials functions in mammals by maintaining efficient glucose, lipid, and protein metabolism (Steven *et al.*, 1976). In general, the toxicity of Cr (III) to mammals is low because it tends to form stable complexes with negatively charged inorganic or organic compounds in aqueous or particulate form; so its membrane permeability is poor and it is noncorrosive; further, there is little tendency for Cr⁺³ to biomagnify in food chains in the inorganic form.

However, the oxidizing potential of Cr (VI) is high, so it easily penetrates biological membranes and so is more toxic than the +3 form (Steven *et al.*, 1976; Taylor and Parr, 2002; Yasuko *et al.*, 2006). All stable Cr⁺⁶ anionic compounds strongly oxidize organic matter on contact and yield oxidized organic matter and Cr⁺³ (Yasuko *et al.*, 2006).

Toxicity of Cr (VI) among warm-blooded organisms has been reported, for instance it was fatal to dogs in 3 months at 100 mg/kg in their food and killed most mammalian experimental animals at injected doses of 1 to 5 mg Cr/kg body weight (Ronald, 1986). Cr (VI) compounds may cause skin ulceration, irritative dermatitis, ulcerations in mucous membranes, and perforations of the nasal septum. That inhalation of Cr⁺⁶ compounds may cause bronchial carcinomas has been well documented in humans (Langard and Norseth, 1979). Chromate producing industrial workers developed respiratory cancer after been exposed to 30 to 1,100 µg/m³ Cr (VI) in air for periods of 4 to 24 years, and workers producing chromate pigment reportedly developed respiratory cancer after been subjected to an estimated Cr⁺⁶ exposure of 500 to 1,500 µg/m³ for 6 to 9 years (Post and Campbell, 1980).

At least three distinct Cr⁺⁶ excretion patterns exist in rats: blood has a single component, with a biological half-life of 13.9 days; testes, brain, kidney and lung have two components; and liver has three components with half-lives of 2.4h, 52.8h and 15.7 day (Yamaguchi and SanoK, 1983). Cr ion accumulation in organs of albino rats 72h after each of the four-weekly intraperitoneal administration of 1.0 mL of 1.0 µg/mL K₂Cr₂O₇ solution and another four weeks of recovery using atomic absorption spectrophotometry indicated that the total Cr ion in the organs of both sexes of the rat followed the ranking blood > heart > liver; though the male accumulated more (Omoniyi *et al.*, 2010).

Wiegand *et al.* (1985) described the *in vitro* uptake kinetics of hexavalent chromium in erythrocytes of rats and humans. Blood sample is the commonest for evaluating systemic accumulation of heavy metal pollutants compared to hair, saliva and urine. The body's excretory mechanisms may eliminate particles and particle breakdown products as substances are transported in the blood or the lymph. So the levels of released toxins such as metal ions may be detected and determined in body fluids such as serum or plasma, urine and sweat (Howie *et al.*, 1996; Merrit and Brown, 1996). Compared to urine and sweat, hair has the advantage of long term memory; as a three-inch strand of human hair will give a six month history of the body chemistry (Wein-Schwartz and Oderda, 2000).

The concentrations of metals such as As, Cd, Hg and Pb in hair samples has been reported to tend to correlate with the levels of these metals in internal organs (Maugh, 1978; Sharma and Kumar, 2004), this is one of the reasons why hair is used as a screening tool for forensic purposes. The ease of collecting hair samples compared to the blood of an animal also places hair ahead of blood for screening purposes. For instance, the determination of the level of Cr in the hair samples of diabetic patients is carried during the screening procedure, since low levels of chromium in the hair

have been demonstrated in victims of juvenile-onset diabetes (Maugh, 1978).

This study reports the affinity of the components of blood of female albino rat (*Rattus nervigicus*) for bioaccumulation of Cr ions after 7 days of recovery from two weekly intraperitoneal administrations of 5.0 mL of 1.0 µg/mL K₂Cr₂O₇ solution on day 1 and 1.0 mL of the solution on day 8; and to compare these to the bioaccumulation level in the fur of the animal model, in order to furnish additional scientific data for forensic analyses of industrial and environmental exposures to Cr.

2. MATERIALS AND METHODS

Intraperitoneal administration of K₂Cr₂O₇ solution to albino rat

Twelve female albino rats (about 5-7 weeks old) of body weight ranging from 150 - 178 g were collected and bred in the animal house unit of Faculty of Pharmaceutical Science, Ahmadu Bello University, Zaria, Nigeria in May 2012. Six rats (E1 - E6) served as the experimental and six (C1 - C6) as control group. The two sets of rats were fed adequately with commercial rat pellets (Bendel Feeds and Flour Mill, Ewu, Nigeria) and tap water *ad lib*. The feed and water were analyzed for the level of chromium to serve as a baseline.

Five millilitre of 1.0 µg/mL K₂Cr₂O₇ solution - prepared by dissolution in Hank's solution (0.9% NaCl solution) (Lori and Hanawa, 2001) - was injected intraperitoneally into the experimental rats in the morning of day 1 and 1 mL of the same solution on day 8 of the experiment. The responses to the exposures were observed from day 8-15; after which both groups were sacrificed by lethal dose of chloroform. The experiment conducted in this study was in accordance with the Ahmadu Bello University committee on animal use and care. And it was ensured that the animal received humane care according to the European Convention on Animal Care.

The concentration of the Cr solution was chosen based on the result of Omoniyi *et al.* (2010) in which the administration of 1.0 mL of 0.1, 1.0 or 5.0 µg/mL K₂Cr₂O₇ solution - prepared by dissolution in Hank's solution - to female and male rats done intraperitoneally was used to determine the dose response 72 h after the administrations. An administration of 1.0 mL of 1.0 µg/mL f K₂Cr₂O₇ solution gave the optimum dose for an obscured physical effect in the animal model (Omoniyi *et al.*, 2010).

The fur of the animal from both groups was collected using a cleaned pair of scissors. Blood (3.5 mL) was collected from both study groups using a 5 mL syringe after dissection of the animal. The blood was separated into erythrocytes, serum and blood plasma by centrifugation, thus:

Separation of serum from whole blood

A 1.0 mL aliquot of the blood sample in a micro-centrifuge tube was positioned slantingly and undisturbed at room temperature for 2 h to allow the blood to clot. The blood was then centrifuged at 2000 rpm (1650 x g) for 20 minutes. The resulting supernatant which is the serum was measured and transferred into a beaker using a Pasteur pipette (<http://www.nottingham.ac.uk/methods/serumseparationfro.html>).

Separation of blood plasma and erythrocytes from whole blood

To an aliquot of 2.5 mL fresh whole blood collected from each of the experimental and control rat in a glass tube, 0.1 mL of 2% potassium oxalate solution was added; the mixture was kept on ice to maintain a temperature of 4°C. The blood was then centrifuged at 700 x g/5 minutes to separate the components. The resulting component was a clear solution of blood plasma in the upper phase of the centrifuge tube which was then separated by using a Pasteur pipette. The buffy coat, which is a thin layer of leukocytes (white blood cells), mixed with platelets in the middle. Erythrocytes (red blood cells) at the bottom of the centrifuge tube were carefully collected by using a Pasteur pipette to get rid of the leukocytes (Christain, 1978; David, 2011).

Quality assurance and acid digestion

Quality assurance of the atomic absorption machine was carried out by using four kidneys of the experimental rats; these were dried in an oven at a temperature of 60°C for about 24 h. The dried kidneys were pulverized using an agate mortar and pestle. A 5.0 mL aliquot of 1.0 µg/mL K₂Cr₂O₇ solution was added to one half of the pulverized kidney in a beaker, while the other half had no addition of the solution.

Wet digestion of each of the 1.0 mL blood components, the fur obtained from the two groups of the animal and the kidney for spiking experiment was carried out using concentrated HNO₃ and HCl in the ratio of 1:3 on a hotplate at 90°C for 10 minutes, while adding warm distilled water at the point of dryness. To the filtrate of each digest, distilled water was added in order to make it up to 10 mL in a volumetric flask. The level of Cr in each solution was then determined by atomic absorption spectrophotometry (Buck Scientific) at wavelength 357.9 nm.

RESULTS AND DISCUSSION

Behavioural pattern of the rats after injection

On day 1 of the experiment, when 5 mL of 1 µg/mL K₂Cr₂O₇ solution was administered, few hours later the experimental group looked dull and inactive compared to the control group. On the day 2 water intake of the experimental group was 25 mL which was relatively low compared to the 50 mL of the control group. This trend of inactivity of the experimental group continued for day 3 and 4 of the experiment with volume of water intake being 35 mL and 50 mL for the experimental and control group respectively on day 3, and 48 mL and 50 mL for the experimental and control group respectively on day 4 of the experiment. The same trend was observed for the experimental group in terms of the amount of feed intake which was less than the control group. The recovery period started from the late hours

of day 4 and continued up to the early hours of day 8, during which the experimental group became active and showed increase in the amount of water and feed intake, with the volume of water intake ranging from 50 mL – 75 mL and 50 mL – 73 mL for the experimental and control group respectively.

In the second week of the experiment i.e. on day 8 to day 15 during which 1.0 mL of 1 µg/mL K₂Cr₂O₇ solution was administered to the experimental group; there was a slightly different behavioural pattern for the experimental group. Few hours after the administration, the experimental group looked dull and inactive compared to the control group. Water and feed intake of the experimental group were less than the control group, with volume of water intake being 60 mL and 75 mL for the experimental and control group respectively. The same trend was observed on day 9 of the experiment. The recovery period for the second administration started during the late hours of day 9 and continued up to the end of the experiment at day 15 when the rats were sacrificed.

The finding led to the deduction that an intraperitoneal dose of 5 mL of 1.0 µg/mL K₂Cr₂O₇ solution to albino rat is a mild exposure dose with noticeable behavioural recovery after the fourth day of the intraperitoneal administration. However, intraperitoneal administration of 1 mL 1.0 µg/mL K₂Cr₂O₇ solution led to noticeable behavioural recovery after the second day of the administration.

Mean chromium concentration in components of blood and fur of experimental and control rats

The mean % recovery for the spiking experiment carried out using kidney in order for quality assurance ranged from 78.9 ± 0.23 to 88.7.2 ± 0.52.

The mean level of total-Cr ions in the drinking water and feed given to the rats during the study were 0.0023 and 0.0036 µg/mL respectively; since these were below the allowable limits, the levels were insignificant to affect the accuracy of the results for the blood and fur.

The mean chromium concentration (n = 6) in the serum, red blood cells (RBC), plasma and fur of the experimental and control groups depicted in Figure 1 indicated that for all the samples studied, the levels of Cr in the control group were lower than the experimental counterparts. The ranking for the mean level of total-Cr ions in the experimental followed the ranking, serum > fur > RBC > plasma; while the control had the ranking serum > fur > plasma > RBC. Following from Table 1, the highest variability in the level of total-Cr ions between the experimental and control group was found in serum having coefficient of variance (CV) being 68.32%, this was closely followed by fur (63.98%) while the lowest was in plasma (4.02%).

Table 1: Minimum and maximum levels of total-Cr ions in the components of blood and fur experimental and control rats following intraperitoneal administration of K₂Cr₂O₇ solution

Sample	Minimum Cr (µg/mL)	Maximum Cr (µg/mL)	SD	CV (%)
Serum experiment	9.633	15.650	0.726	68.32
Serum control	5.700	8.175	0.181	
RBC experiment	1.854	3.100	0.080	12.10
RBC control	1.345	3.460	0.072	
Plasma experiment	2.013	3.944	0.004	4.02
Plasma control	1.480	3.238	0.048	
Fur experiment	4.01	6.432	0.0078	63.98
Fur control	1.987	2.87	0.021	

The information from CV indexes that the systemic Cr ions in rat has highest affinity for serum followed by the fur and least in plasma. This conforms to the assertion of Katharine and Stanley (1985) in the study undertaken *in vivo* and *in vitro* on the binding of metal ions as metal salts or as corrosion products (Co⁺², Ni⁺², Cr⁺³, Cr⁺⁶) to tissues and cells; the findings reveal that

most of the metal was in serum with some binding to cells. The most extensive binding to cells was recorded for Cr⁺⁶, which was the most strongly bound to red blood cells compared to white blood cells and tissue (Katharine and Stanley, 1985; Merrit and Brown, 1996). The presence of serum does not affect binding to the RBC and white blood cells.




Figure 1: Mean concentrations of total-chromium ions in the components of blood and fur of experiment and control rats following intraperitoneal administration of K₂Cr₂O₇ solution

The highest level of total-Cr ions recorded for serum in this study (12.127 ± 0.726 µg/mL) is explained by the fact that the albumin and gamma globulin in serum has a very high propensity for binding metal ions. The blood pH of about 6.7 is basic to the isoelectric point of albumin (a protein with negative charges) so affinity for bioaccumulation of Cr⁺⁶ was highest in this component of blood; while the blood pH is acidic to the isoelectric point gamma globulin (a protein with negative charges; the amount of albumin in serum is about five times that of gamma globulin (Katharine and Stanley, 1985).

The mean level of Cr ions bioaccumulated in the experimental fur was 4.090 ± 0.0078 µg/mL; this was almost two-fold relative to the 2.23 ± 0.021 µg/mL obtained in the control rats. The affinity can be traced to the absorption by the keratinous component of fur where the large number of disulphide bonds of keratin can form electrostatic bond with Cr⁺⁶ in the system; thus

Cr likely has a higher degree of protein binding (Maletin, 1988; Pautard, 1963).

Following from Figure 1, the serum level of total-Cr ion in the experimental animal was 3-fold that in fur, 4-fold that in RBC, and 6-fold that in plasma. For the control rat, the serum level of Cr ion was 3-fold than those recorded in fur, RBC and plasma.

Though, the animals were sacrificed after the intraperitoneal administration of 5.0 mL and 1.0 mL of the 1.0 µg/mL K₂Cr₂O₇ solution. It is anticipated that after the administration of the higher amount of K₂Cr₂O₇ solution on day 1, the plasma to erythrocyte (RBC) ratio of total chromium ions accumulated would decrease with increasing hexavalent chromium concentration administered intraperitoneally. In accordance to the finding of Corbett *et al.* (1998), the rate of excretion of Cr ion in the plasma is much more than the rate of excretion in the serum and red blood cells; because Cr binds to both transferrin and to a lesser degree albumin in plasma (Borguet *et al.*, 1990; Vincent, 2000;

Borguet *et al.*, 1995). The plasma to erythrocyte ratio of total chromium ions accumulated decreased with increasing hexavalent chromium concentration administered intraperitoneally. This infers that the components of blood (serum, red blood cells) in the female animal are more prone to chromium toxicity than the plasma.

The Cr levels in the fur and serum were doubled due to the administration of $K_2Cr_2O_7$ solution to the rats; this signals that the fur and serum can equally serve as specimen for detection of mild exposure to Cr (VI) during forensic investigation.

Duncan multiple range test (two-way) at $P < 0.05$ was used to investigate the statistical relationship that existed between the groups and the blood components/fur. There was high statistical increase in the total-Cr ions in the serum of the experimental rat compared to the control; likewise the level in the experimental fur was highly elevated compared to the control ($P < 0.05$). A slight significant increase was recorded in the level of Cr ion of the experimental RBC compared to the control. There was no statistical difference between the bioaccumulated total-Cr ions in plasma of the experimental to the control. There was significant difference in the Cr ion level in the serum compared to the other specimens studied for both groups ($P < 0.05$).

Weight of rat to component bioaccumulation of Cr ions

The mean body weight (154.0 g) of the control rats increased by 19% at day 15; this indicated a significant difference in weight during the 15-day experiment. The mean body weight of the experimental rats was 163 g, this increased by 2% at day 15. There was no significant difference in weight of the experimental rats during the 15-day experiment.

The total amount of chromium ions accumulated in the whole blood of the experimental rats, decreased as the weight increases. These was highest in rat E1 of Cr ion concentration 24.79 $\mu\text{g/mL}$ followed by E5 of Cr ion concentration 14.756 $\mu\text{g/mL}$ of and least in E4 with concentration 12.33 $\mu\text{g/mL}$; recording the initial weights 153 g, 178 g and 158 g and final weights 155 g, 162 g and 182 g respectively. The same trend was observed for the control group. These was highest in C4 of chromium ion concentration 16.873 $\mu\text{g/mL}$ followed by C3 of chromium ion concentration 6.766 $\mu\text{g/mL}$ and least in C1 having Cr concentration 6.587 $\mu\text{g/mL}$; these had initial weights 146 g, 150 g, 167 g and final weights 156 g, 165 g and 172 g respectively.

The trend of the decrease in total-Cr ion concentration as weights increases might have resulted from the rats of smaller weights not been able to excrete effectively the intraperitoneally injected chromium solution. This

implies that the female rats with the larger weights did more effectively. This trend indicates that body weight has a negative correlation to Cr^{+6} bioaccumulation by whole blood of the animal.

4. Conclusion

High dose of chromium (VI) resulted to inactivity and less consumption of feed and water intake. While there was insignificant increase in the weights of the experimental group at day 15, the control group showed a significant weight gain of 19% at day 15.

The ranking for the mean level of Cr ion in the experimental followed the ranking, serum > fur > RBC > plasma; while the control had the ranking serum > fur > plasma > RBC. The highest variability in the level of total-Cr ions between the experimental and control group was found in serum CV 68.32%, this was closely followed by fur (63.98%) and the lowest was in plasma (4.02%). Therefore, this signals that serum and fur can serve as bio-indicator of the system level of Cr (VI). The serum level of Cr ion in the experimental animal was 3-fold that in fur, 4-fold that in RBC, and 6-fold that in plasma. For the control rat the serum level of Cr ion was 3-fold than those recorded in fur, RBC and plasma.

There was high statistical increase in the Cr ions in the serum of the experimental rat compared to the control; likewise the Cr level in the experimental fur was highly elevated compared to the control ($P < 0.05$). A slight significant increase was recorded in the level of Cr ion of the experimental RBC compared to the control. There was no statistical difference between the bioaccumulated Cr ions in plasma of the experimental to the control.

There is negative correlation between the body weight of the rat and Cr^{+6} bioaccumulated in whole blood samples.

The standard level of chromium in the serum, plasma and red blood cells by the National institute of health is in the range of 0.05 to 0.5 $\mu\text{g/g}$ and the WHO international standards for drinking water recommend a maximum allowable concentration of 0.05 mg/L for chromium (VI). Although few adverse effect have been linked to high intakes of chromium, mean concentration of Cr (VI) found in the serum, red blood cells and plasma were 12.128 $\mu\text{g/g}$, 2.956 $\mu\text{g/g}$ and 2.21 $\mu\text{g/g}$ respectively, showing that concentration found in the animal model could still be detrimental to the animals.

The study indicates that mild exposure of the physiological system to Cr ions could lead to mean accumulated levels of Cr ions that might pose health risk in the future. Therefore, it is recommended that there should be caution in the exposure of the body to Cr ions through ingestion, inhalation or dermal contact during occupational and domestic activities, even at mild doses.

REFERENCES

- Adya, N., Alam, M., Ravindranath, T., Mubeen, A. and Saluja, B. (2005). Corrosion of titanium dental implants: Literature review. *Journal of Indian Prosthodontic Society*,**5**: 126-131.
- Amoikon, E.K., Fernandez, J.M, Southern, L.L, Thompson, D.L., Ward, T.L., Olcott, B.M. (1995). Effect of chromium tripicolinate on growth hormones. *Journal of Animal Science*,**58**: 102-104.
- Becker, N., Chang-Claude, J., Frentzel-Beyme, R. (1991). Risk of cancer for arc welders in the Federal Republic of Germany: Results of a second follow-up (1983-1988). *Britanian Journal of Industrial Medicines*,**48**: 675-683.

- Borguet, F., Cornelis R, Lameire N. (1990). Speciation of chromium in plasma and liver tissue of end stage renal failure patients on continuous ambulatory peritoneal dialysis. *Biological Trace Elements Research*, **26**: 449–460.
- Borguet, F., Cornelis, R., Delanghe, J. (1995). Study of the chromium binding in plasma of patients on continuous ambulatory peritoneal dialysis. *Clinical Chemimica Acta*, **238**: 71–84.
- Christain, G.D. (1978). Analytical Chemistry. Vol. II, Pp. 553-557.
- Corbett, G.E., Finley, B.L., Paustenbach, D.J. (1998). Systemic uptake of chromium in human volunteers following dermal contact with hexavalent chromium (22 mg/L). *Exposure Analytical and Environmental Epidemiology*, **7**(2): 179–189.
- David, R.C. (2011). Rice University experimental biosciences. Blood and Erythrocyte Fractionation.
- Ecological Analysts, Inc. (1981). The sources, chemistry, fate, and effects of chromium in aquatic environments. American Petroleum Institute. 2101 L St., N.W., Washington, DC 20037. Pp. 207.
- Howie, D.W., Rogers, S.D., McGee, M.A., Haynes, D.R. (1996). Biologic effects of cobalt chrome in cell and animal models. *Clinical Orthopaedic Related Research*, **329S**: S217-S232.
- Katharine, M., Stanley, A.B. (1985). Biological effects of corrosion products from metals: Corrosion and degradation of implant materials. second symposium, American Society for Testing and Materials, Philadelphia, pp. 195-207.
- Kempson, I.M., Skinner, W.M., Kirkbride, K.P. (2007). The occurrence and incorporation of copper and zinc in hair and their potential as bioindicators: a review. *Journal of Toxicology and Environmental Health*, **B 10**: 611–622.
- Kortenkamp, A. (1997): Problems in the biological monitoring of chromium (VI) exposed individuals. *Biomarkers*, **2**:73-80.
- Langard, S., Norseth, T. (1979). Chromium. In: Friberg, L., Nordberg, G.F., Vouk, V.B. (eds.). Handbook on the toxicology of metals. Elsevier/North Holland Biomedical Press. Pp. 383-397.
- Lori, J.A. and Hanawa, T. (2001). Adsorption characteristics of albumin on gold and titanium in Hank's solution using EQCM. *Corrosion Science*, **43**: 2111.
- Maletin, Y.A. (1988). Institute of general and inorganic chemistry. Academy of Sciences of the Ukrainian SSR, Kiev. Translated from Teoreticheskaya I Eksperimental'naya Khimiya, **24** (4): 450–455.
- Maugh, T.N. (1978). Hair: a diagnostic tool to complement blood serum and urine. *Science*, **202**:1271–1273.
- Merrit, K., Brown, S.A. (1996). Distribution of cobalt-chromium wear corrosion products and biologic reactions. *Clinical Orthopaedics*, **329S**: S233-S243.
- Methods of serum separation. Accessed 27 March 2012. Available: <http://www.nottingham.ac.uk/methods/serums/eparationfro.html>.
- Omoniyi, K.I., Ekwumemgbo, P.A., Aligwo, C., Balogun, M., Adewusi, A.M. (2010). Effect of gender on the dose accumulation of chromium ions in organs of *Rattusnervigicus*. *Electronic Journal of Environment, Agriculture and Food Chemistry*, **9**(4): 735-741.
- Pautard, F.G.E. (1963). Mineralization of keratin and its comparison with the enamel matrix. *Nature*, **199**: 531–535.
- Post, M.A., Campbell, P.G. (1980). Lead chromate pigments—a literature survey on environmental and toxic effects. U.S. Dep. Comm. National Bureau of Standard Report, NBSIR 80-1974. Pp. 38.
- Ronald, E. (1986). Chromium hazards to fish, wildlife, and invertebrates: a synoptic review. *Contaminant Hazard Review of Biological Report*, **85**(1.6) January(6).
- Sharma, R. and Kumar, A. (2004). Trace element contents in human head hair of residents from Agra City (India). *Bulletin of Environmental Contamination and Toxicology*, **72**: 530-534.
- Steven, J.D., Davies, L.J, Stanley, E.K, Abbott, R.A., Ihnat, M., Bidstrup, L., Jaworski, J.F. (1976). Effects of chromium in the Canadian environment. Natural Resources Council Canada, NRCC No. 15017. NRCC/CNRC, Ottawa, Canada, K1A OR6. Pp.168.
- Taylor, F.G. (Jr.), Parr, P.D. (2002). Distribution of chromium in vegetation and small mammals adjacent to cooling towers. *Journal of Tenn Academy of Science*, **53**:87-91.
- Vincent, J.B. (2000). The biochemistry of chromium. *Journal of Nutrition*, **130**: 715–718.
- Wein-Schwaitz, W., Oderda, G.M. (2000). Clinical toxicology. In: Therapeutics, drug and disease management. (7th Ed.). Baltimore Press, Wicliams and Wilkins. Pp. 51.
- Wiegand, H.J., Ottenwalder, H., Bolt, H.M. (1985). Fast uptake kinetics *in vitro* of ⁵¹Cr (VI) by red blood cells of man and rat. *Archives of Toxicology*, **57**(1): 31–34.
- Yamaguchi, S., SanoK, S.N. (1983). On the biological half-time of hexavalent chromium in rats. *Industrial Health*, **21**:25-34.
- Yasuko, T., Takao, H., Daisuke, K., Masahiro, E. (2006). Corrosion of stainless steel sternal wire after long-term implantation. *Journal of Artificial Organs*, **10**(2):145-148.