



# Biokemistri

An International Journal of the Nigerian Society for Experimental Biology

## Original Article

# Antipyretic and analgesic activities of aqueous extract of *Acacia nilotica* root

Lukman A. Alli<sup>1\*</sup>, Mikhail O. Nafiu<sup>2</sup>, Abdulfatai A. Adesokan<sup>3</sup>, Musbau A. Akanji<sup>2</sup>, Adeniyi Y. Tijani<sup>4</sup>, Oluwakanyinsola A. Salawu<sup>4</sup>.

<sup>1</sup>Department of Medical Biochemistry, College of Health Sciences, University of Abuja, Nigeria.

<sup>2</sup>Department of Biochemistry, Faculty of Life Sciences, University of Ilorin, Nigeria.

<sup>3</sup>Department of Biochemistry, College of Health Sciences, Ladoke Akintola University, Ogbomosho, Nigeria.

<sup>4</sup>Department of Pharmacology and Toxicology, National Institute for Pharmaceutical Research and Development (NIPRD), Abuja, Nigeria.

\*Corresponding author: Lukman Alli. Email: adewalealli@gmail.com; Tel.: +2348033664037.

Received: 21 May 2014; Revised 10 June 2014; Accepted: 10 June 2014.

**ABSTRACT:** This study was designed to investigate the scientific basis for the use of *Acacia nilotica* root extract for treatment of fever and pain in traditional medical practice. Anti-Pyretic study was carried out using Brewer's yeast suspension to induce pyrexia. The hot plate, tail immersion and acetic acid-induced writhing tests were the nociceptive models used for analgesic study. Anti-pyretic and analgesic activity of the extract was compared with acetaminophen that was used as control drug. Five groups comprising five animals per group were used for each study. Group 1 was administered 10 ml/kg body weight of distilled water, Group 2 was administered 150 mg/kg body weight of acetaminophen while groups 3, 4 and 5 were administered 100, 200 and 400 mg/kg body weight of extract respectively as single oral dose. The extract produced significant dose-dependent reduction in rectal temperature of rats at 200 and 400 mg/kg body weight. Significant analgesic activities were also observed in the hot plate, tail immersion and acetic acid induced writhing, after administration of 200 and 400 mg/kg b.w of extract which is comparable to the control drug, acetaminophen. The results from this study showed that aqueous extract of *Acacia nilotica* root at 200 and 400 mg/kg body weight possess significant antipyretic and analgesic activities. This provides scientific support for its traditional medical use in the treatment of fever and pain.

**KEYWORDS:** Antipyrexia, Analgesia, *Acacia nilotica*, Brewer's yeast.

BKM.2014.015 © 2014 Nigerian Society for Experimental Biology; All rights reserved. Printed in Nigeria  
This article is downloadable online in PDF format at <http://www.bioline.org.br/bk>

## INTRODUCTION

*Acacia nilotica* (Linn.) Willd. Ex Del. (also called Gum Arabic tree; 'Gabaruwa' in Hausa and 'Igi Kasia/Booni' in Yoruba) is an important plant used in traditional medical practice in Northern Nigeria and many African countries (Bargal and Bargali, 2009), for the treatment of different ailments such as diabetes, diarrhea (van Wyk, 2000), tuberculosis (Oladosu *et al.*, 2007) and malaria (Alli *et al.*, 2011). It is rich in phytochemicals such as alkaloids, flavonoid, tannins, terpenes and phenolics [Banso, 2009; Alli *et al.*, 2011] that are responsible for some of its pharmacologic activities. Although studies have been carried out to validate some of

the local uses of the stem and bark extract of this plant, there is need to investigate the anti-pyretic and analgesic activity of the extract of *A. nilotica* root in order to validate the use of the root extract in traditional treatment of fever and pain. As pain and fever are the main symptoms of malaria, this study is therefore investigating the anti-pyretic and analgesic activity of aqueous extract of *Acacia nilotica* root, after its anti-plasmodial activity has been established from previous studies (Alli *et al.*, 2011).

Pain and Pyrexia are common physiologic response to infection, inflammation and tissue damage (Spacer and Breder, 1994). Pain may occur as a result of nociceptive,

inflammatory or neuropathic conditions that could involve biochemical mediators like prostaglandins, substance and glutamate (Raffa, 2006). Pyrexia could occur as body's natural defense mechanism to prevent survival of infectious agents (Cooper, 1995; Woolf, 2010). It is initiated when infectious agents or damaged tissues triggers increase production of pro-inflammatory mediators (pyrogens such as, Tumour Necrosis Factor- $\alpha$  {TNF- $\alpha$ }, interleukin 1b and interleukin-6) which increase the synthesis of prostaglandin E-2 (PGE-2) near the pre-optic area of anterior hypothalamus. The hypothalamus is consequently triggered to elevate the body temperature (Akpan *et al.*, 2010). The mechanism of action of commonly used anti-pyrexia involves inhibition of cyclooxygenase enzyme (COX), specifically COX-2, thereby interrupting the synthesis of prostaglandin E-2, and thus reducing body temperature (Cheng *et al.*, 2005). Some of the commonly available synthetic antipyretic agents, with analgesic properties, (such as acetaminophen, acetyl salicylic acid and other non-steroidal anti-inflammatory drugs) could be toxic to hepatic cells and glomeruli (Cheng *et al.*, 2005). Therefore, search for safe and effective plant-based antipyretic and analgesic agents is highly essential and desirable.

## MATERIALS AND METHODS

### Experimental animals

Healthy Wistar albino rats (190  $\pm$  10 g) and albino mice (25  $\pm$  5 g) used for this study, were obtained from the animal facility center of National Institute for Pharmaceutical Research and Development (NIPRD), Abuja, Nigeria. They were housed in well ventilated cages, fed with rat pellets (Pfizer feeds Plc, Lagos, Nigeria) and water *ad libitum*. Standard laboratory conditions were maintained in accordance with the guideline for the care and use of Laboratory animals by National Academy of Sciences. The ethical committee on animal use and care of NIPRD approved the use of laboratory animals for this study.

### Plant materials

Root samples of *Acacia nilotica* were collected by Mallam Muazzam, from Bamburu-Chaza village, Suleja, Niger State, Nigeria. The samples were identified and authenticated at the herbarium of NIPRD, by a taxonomist, Mrs Grace Ugbabe. Voucher specimen (NIPRD/H/6401) was prepared and deposited at the herbarium for referencing.

### Preparation of Aqueous Extract of *Acacia nilotica* root

The root sample of *A. nilotica* was air dried under shade for two weeks and pulverized using grinding machine. The powder was stored in an airtight container and kept in a cool dry place. Aqueous extraction was carried out following the cold maceration method described by Adzu *et al.*, (2003). Two hundred grams of the powdered root of *A. nilotica* was

soaked in 1 litre of distilled water and kept for 24 hours. The suspension was then filtered with muslin cloth followed by filtration with Whatman filter paper (No.1). The filtrate was freeze-dried using AMSCO/FINN-AQUA GT2 Freeze dryer (Germany).

### Reagents

Acetaminophen (Emzor Pharmaceutical brand, NAFDAC: 04-0411) was obtained from Novac Pharmacy, Kubwa, Abuja, Nigeria. Brewer's yeast was obtained from Nigerian Breweries Plc, Lagos, Nigeria. Solutions of acetic acid (1% v/v) and 150 mg/kg body weight acetaminophen were prepared with distilled water. All other reagents were of analytical grade and prepared in glass bottles using distilled water.

### Acute toxicity study

Acute toxicity study of the aqueous extract of *A. nilotica* root was carried out following the Organization for Economic Cooperation and Development (OECD) Guideline 423 (OECD, 2001). Nine healthy, nulliparous and non-pregnant female albino mice weighing 20  $\pm$  2 g were used for this study. They were weighed and randomized into three groups of three mice per group. The animals were fasted for 4 hours prior to dosing. After 4 hours of fasting, the control (Group 1) was administered 10 ml/kg body weight of distilled water, while Groups 2 and 3 were administered oral doses of the extract at 300 and 2000 mg/kg body weight respectively. The mice were observed individually for signs of toxicity during the first 30 minutes, periodically during the first 24 hours and daily for 14 days. Weight changes were recorded twice weekly and mortality recorded daily. LD<sub>50</sub> of the root extract was determined by comparing number of mortality with fixed LD<sub>50</sub> cut-off values in the guideline.

### Phytochemical screening

Phytochemical screening of the crude extract was carried out following the methods described by Sofowora (1993) to determine the presence of secondary metabolites such as alkaloids, phenolics, flavonoids, saponins, terpenes, tannins, anthraquinones

### Induction of Pyrexia and administration of extract

The method described by Singh *et al.*, (2010), using Brewer's yeast suspension, was followed to induce pyrexia in the rats. Subcutaneous injection of 15% brewer's yeast suspension into rats will produce pyrexia in the animals. The basal rectal temperatures of these rats were measured by inserting clinical thermometer (OMRON digital thermometer, Eco Temp Basic) up to 1 cm into the anus and recording the temperature after 1 minute. The baseline temperature was recorded at 0 hour, after which the rats were administered subcutaneous injection of 10 ml/kg body weight of 15% Brewer's yeast suspension in the abdominal region. Nineteen hours after administration of yeast, the rectal temperature of

each rat was observed and rats that showed an increase in rectal temperature of at least 0.5 °C were selected and randomized into five groups containing five rats per group.

Group 1 served as the control, and received 10 ml/kg body weight of distilled water, Group 2 received 150 mg/kg body weight of acetaminophen, while Groups 3, 4 and 5 received 100, 200 and 400 mg/kg body weight of extract respectively. All the animals received a single oral dose of treatment for each group. The rectal temperatures were taken again, at the 20th, 21st, 22nd and 23rd hour after brewer's yeast injection.

**Table 1: Phytochemical constituents of aqueous extract of *A. nilotica* root.**

Phytochemicals	Quantity (mg/kg)
Phenolics	34.50 ± 1.55
Tannins	27.00 ± 1.75
Alkaloids	23.30 ± 1.86
Saponins	9.80 ± 0.89
Anthraquinones	4.70 ± 0.97
Flavonoids	0.50 ± 0.06
Terpenes	0.10 ± 0.05
Sterols	0.10 ± 0.05

Values are expressed as mean ± S.E.M of 3 observations.

#### Analgesic Activity (Hot Plate method)

The hot-plate latency method described by Hosseinzadeh *et al.* (2000) was used. Healthy Wistar Albino rats (190 ± 10 g) were screened for thermal sensitivity by placing them one-by-one on a hot plate maintained at 55.0 ± 0.5 °C. Rats that stayed on the hot plate for ≥ 2 sec without jumping or licking paws (response to thermal stimulus) were selected for the study. The selected rats were fasted for 12 hours and then randomized into five groups of five rats each. Group 1 served as the control, and received 10 ml/kg body weight of distilled water, Group 2 received 150 mg/kg body weight of acetaminophen, while Groups 3, 4 and 5 received 100, 200 and 400 mg/kg body weight of extract orally, respectively.

The rats were each placed on a hot plate (maintained at 55.0 ± 0.5 °C) 30 minutes after administration of distilled water, acetaminophen and extract. The time taken for the rats to respond to the thermal stimulus was noted as the latency of response (in seconds) and the mean latency for each group was calculated. The effects of the administration of extracts, acetaminophen and distilled water on response of the rats to thermal stimulus (by licking paws or jumping) were also determined after 30, 60, 90 and 120 minutes. A latency time

of 20 seconds was regarded as complete analgesia and measurement was terminated if 20 sec is exceeded to avoid tissue injury to the animals.

#### Tail Immersion Method

The tail immersion method described by Jansen and Jagenav (1959) was followed. Healthy Wistar Albino rats (190 ± 10 g) were screened for thermal sensitivity by putting the tip of their tail (lower 5 cm portion) inside a beaker containing hot water maintained at 55.0 ± 0.5 °C. Rats that maintained their tail in the hot water for ≥ 2 sec were selected for the study. Twenty five albino rats selected for this study were randomized into 5 groups of 5 rats each, and placed into restraining cages leaving the tail to hang out freely. Group 1 served as the control, and received 10 ml/kg body weight of distilled water, Group 2 received 150 mg/kg body weight of acetaminophen, while Groups 3, 4 and 5 received 100, 200 and 400 mg/kg body weight of extract orally, respectively.

The lower 5 cm portion of the tail was immersed in a beaker filled with hot water maintained at 55.0 ± 0.5 °C. The time (seconds) taken for the withdrawal of the tail from hot water induced pain recorded with a stop watch as the reaction time, was recorded for each rat at intervals of 30, 60, 90 and 120 minutes after the extract and drug administration. The maximum cut-off time for immersion in hot water was 15 seconds to avoid injury to tissues of the tail.

#### Acetic acid induced writhing test

Analgesic activity of the aqueous extract of *A. nilotica* root to peripheral pain was assessed using the method described by Salawu *et al.*, (2008). Twenty-five mice were randomly divided into five groups of 5 mice each. Group 1 served as the control, and received 10 ml/kg body weight of distilled water, Group 2 received 150 mg/kg body weight of acetaminophen, while Groups 3, 4 and 5 received 100, 200 and 400 mg/kg body weight of extract orally, respectively. Thirty minutes after treatment, each mouse received intraperitoneal injection of 10 ml/kg body weight of 1% (v/v) acetic acid solution and subsequently placed in a glass beaker for observation. The number of writhes (contraction of abdomen, twisting of trunk and extension of the hind limbs) per animal was counted within 30 minutes observation period, beginning from 5 min after acetic acid injection. Inhibition (%) was calculated using the following expression.

$$\text{Percentage Inhibition (\%)} = \frac{\text{Average writhe (Control)} - \text{Average writhe (Test)}}{\text{Average writhe (Control)}}$$

**Table 2: Antipyretic activity of aqueous extract of *A. nilotica* (A.N) root**

Treatment/Dose (mg/kg body weight)	Rectal Temperature (°C) of rats					
	0 hour pre- injection of yeast	19 hour post- injection of yeast	Temperature after administration of extract			
			20 hour	21 hour	22 hour	23 hour
Control	37.60 ± 0.25	38.50 ± 0.20	38.60 ± 0.25	38.70 ± 0.22	38.90 ± 0.24	39.00 ± 0.25
Acetaminophen 150	37.60 ± 0.26	38.60 ± 0.25	38.40 ± 0.23	38.00 ± 0.22*	37.80 ± 0.18*	37.60 ± 0.15*
Extract 100	37.50 ± 0.31	38.50 ± 0.20	38.40 ± 0.22	38.40 ± 0.25	38.30 ± 0.25	38.20 ± 0.20
200	37.50 ± 0.29	38.60 ± 0.28	38.40 ± 0.25	38.30 ± 0.22	38.20 ± 0.24	37.90 ± 0.23*
400	37.60 ± 0.26	38.60 ± 0.17	38.40 ± 0.25	38.20 ± 0.21	37.90 ± 0.19*	37.70 ± 0.15*

Values are expressed as mean ± S.E.M of five observations; \* = significantly different at  $P < 0.05$  when compared to control.

**Table 3: Analgesic activity of aqueous extract of *A. nilotica* root using hot plate test**

Treatment/Dose (mg/kg b.w)	Mean Basal time (sec)	Mean reaction time (sec) at interval of 30 min			
		30	60	90	120
Control	2.54 ± 0.15	2.55 ± 0.20	2.54 ± 0.18	2.56 ± 0.22	2.55 ± 0.20
Acetaminophen (150)	2.55 ± 0.17	5.50 ± 0.15 <sup>*</sup>	6.10 ± 0.20 <sup>*</sup>	6.85 ± 0.22 <sup>*</sup>	7.65 ± 0.25 <sup>*</sup>
Extract 100	2.55 ± 0.20	2.55 ± 0.25	2.65 ± 0.22	2.65 ± 0.27	2.75 ± 0.28
200	2.50 ± 0.17	3.45 ± 0.24	4.80 ± 0.23	5.30 ± 0.20 <sup>*</sup>	5.75 ± 0.25 <sup>*</sup>
400	2.55 ± 0.20	4.85 ± 0.22	5.45 ± 0.15 <sup>*</sup>	5.95 ± 0.25 <sup>*</sup>	6.30 ± 0.30 <sup>*</sup>

Values are expressed as mean ± S.E.M of five observations; \* = significantly different at  $P < 0.05$  when compared to control

### Statistical analysis

Data were expressed as mean ± standard error of mean of five replicates. Statistical analysis was done using Graph pad prism version 5.0. The differences between the mean were compared using analysis of variance (ANOVA). Values with  $P < 0.05$  were considered statistically significant.

### RESULTS

There were no remarkable behavioural (such as reaction to food supply, and contact) changes in the treated mice, during the acute toxicity study period. No mortality was recorded during the observation period following administration of the

extract at the doses of 300 and 2000 mg/kg body weight. The oral median lethal dose ( $LD_{50}$ ) of the extract was estimated to be 5000 mg/kg in mice following the OECD Guideline 423 (2001) for oral acute toxicity study. Absence of mortality in the experimental animals after administration of 2000 mg/kg body weight of extract corresponds to a  $LD_{50}$  value of 5000 mg/kg body weight on the guideline chart.

Initial tests were carried out to detect the presence of common phytochemicals in the aqueous extracts of *A. nilotica* root. The aqueous extract were found to produce positive reactions to phenolics, tannins, alkaloids, anthraquinones, flavonoids, terpenes and sterols (Table 1).

**Table 4: Analgesic Activity of *A. nilotica* root extract using tail immersion test**

Treatment/Dose (mg/kg b.w.)	Mean basal time (sec)	Mean reaction time (sec) at interval of 30 min			
		30	60	90	120
Control	2.20 ± 0.15	2.50 ± 0.20	2.55 ± 0.15	2.45 ± 0.18	2.50 ± 0.15
Acetaminophen (150)	2.35 ± 0.25	4.70 ± 0.22*	5.40 ± 0.25*	6.50 ± 0.22*	7.60 ± 0.20*
Extract 100	2.20 ± 0.20	2.25 ± 0.23	2.25 ± 0.20	2.40 ± 0.25	2.45 ± 0.22
200	2.25 ± 0.23	3.50 ± 0.25	4.20 ± 0.22	5.25 ± 0.21*	5.85 ± 0.23*
400	2.45 ± 0.20	3.85 ± 0.20	4.95 ± 0.26*	5.90 ± 0.23*	6.70 ± 0.25*

Values are expressed as mean ± S.E.M of five observations; \* = significantly different at  $P < 0.05$  when compared to control

**Table 5: Analgesic activity of *A. nilotica* on acetic acid-induced writhes in rats**

Treatment	Dose (mg/kg b.w)	Number of writhes	Inhibition (%)
Distilled water	10	32.50 ± 2.50	-
Acetaminophen	150	8.50 ± 0.65*	74.6*
<i>A. nilotica</i>	100	22.50 ± 1.50	30.7
<i>A. nilotica</i>	200	14.00 ± 1.00*	56.9*
<i>A. nilotica</i>	400	9.50 ± 0.57*	70.8*

Values are expressed as mean ± S.E.M of five observations; \* = significantly different at  $P < 0.05$  when compared to control

There was a significant dose-dependent reduction in the rectal temperature in rats administered with 200 and 400mg extract /kg body weight at 22 and 23 hr after extract administration respectively (Table 2).

Significant dose-dependent increase in reaction time, in the hot plate and tail immersion test, was observed after oral administration of 200 and 400 mg/kg b.w dose of aqueous extract of *A. nilotica* (Tables 3 and 4). The extract at 200 and 400 mg/kg body weight significantly inhibited the acetic acid induced writhing in the mice when compared with the control (Table 5).

## DISCUSSION

The antipyretic and analgesic activity of aqueous extract of *A. nilotica* was studied in rats using standard models. Acetaminophen (paracetamol) is used in this study as control drug because it is one of the most widely used analgesic and antipyretic in both children and adults (Chou *et al.*, 2007; Zhang *et al.*, 2007; Loke, 2009). It has also been shown to have significant concentration in the central nervous system at analgesic dose (Courade *et al.*, 2001). The antipyretic activity of the extract at doses of 200 and 400 mg/kg b.w, is comparable to the antipyretic activity of acetaminophen that

was used as the control drug. This result is similar to the antipyretic activity of aqueous stem extract of *Enantia chlorantha* at 100 and 200 mg/kg dose (Adesokan et al., 2008). Several other plant extracts, such as methanol extract of *Bauhinia racemosa* stem bark (Gupta et al., 2005), aqueous leave extract of *Parquetina nigrescens* (Owoyele et al., 2009), methanolic extract of *Capparis zeylanica* plant (Padhan et al., 2010) and ethanol extract of *Strophanthus sarmentosus* (Agbaje and Ajidahun, 2011) have been reported to possess antipyretic activity in Wistar rats. The dose dependent antipyretic activity of *A. nilotica* may be due to the presence of phenolics and alkaloids which have been reported to possess antipyretic and analgesic properties (Kanagusuku et al., 2007; Owoyele et al., 2008). Flavonoids such as baicalin and alkaloids such as boldine have been reported to exert antipyretic effect by suppressing TNF- $\alpha$  (Chang et al., 2007) or inhibiting the synthesis of prostaglandin E<sub>2</sub> (Backhouse et al., 1994). Yeast-induced pyrexia is based on increase synthesis of prostaglandin (PGE<sub>2</sub>) (Akpan et al., 2012), therefore anti-pyretic activity of the extract may be mediated via inhibition of PGE<sub>2</sub> synthesis. It may also be due to reduction of prostaglandin concentration in the hypothalamus through inhibition of cyclooxygenase pathway or through vasodilatation of superficial blood vessels leading to increase heat loss (Rang et al., 2007).

The analgesic activity of aqueous extract of *A. nilotica* root (using the hot plate and tail immersion test) at 200 and 400 mg/kg body weight dose is significant. This is similar to the result obtained for analgesic activity of aqueous extract of *Curcum longa* rhizome (Neha et al., 2009), aqueous extract of *Parquetina nigrescens* leaves (Owoyele et al., 2009) and aqueous extract of *Mimosa albida* root (Rejón-Orantes et al., 2013). Substances that can produce inhibitory effects in hot plate and tail immersion test can inhibit centrally-induced pain and therefore act as analgesics (Prado et al., 1990). The reason for this is that analgesic action in hot plate and tail immersion tests involves supra-spinal and spinal components of the central nervous system (Mustaffa et al., 2010). The writhes induced by acetic acid were used in this study to assess peripherally acting analgesics (Chou et al., 2007). Intraperitoneal injection of acetic acid can liberate prostaglandins and histamine which will stimulate local peritoneal receptors involved in the writhing response (Kanagusuku et al., 2007). Secondary metabolites from medicinal plants such as alkaloids (*myristica fragrance* seed) (Hayfaa et al., 2013), tannins and flavonoids (such as ethylquercetin (Picq et al., 1991), rutin and bioflavonoids (Calixto et al., 2000; Bittar et al., 2000) have been reported to be responsible for analgesic and antipyretic activities. The significant reduction in acetic acid-induced writhes by aqueous extract of *A. nilotica* suggests that the extract also possesses peripheral analgesic activity.

**Conclusion:** The observed activities of *A. nilotica* extract in the above tests suggest that the aqueous root extract possesses significant antipyretic and analgesic activities which may be due to the alkaloids and flavonoids present in the extract. It may therefore be used as a potent antipyretic and analgesic herbal medication alongside its common use as an antimalarial.

### Author's Contributions

This work was carried out in collaboration between all authors. LAA contributed to conception and design, data acquisition, analysis and interpretation; and writing of the manuscript and its post-review revisions. All other authors contributed to data analysis and interpretation. They were involved in drafting and revising the manuscript for important intellectual content.

### REFERENCES

- Adesokan AA, Yakubu MT, Owoyele BV, Akanji MA, Soladoye AO, Lawal OK (2008) Effect of administration of aqueous and ethanolic extracts of *Enantia chlorantha* stem bark on brewer's yeast-induced pyrexia in rats. *African Journal of Biochemistry Research* 2: 165–69
- Adzu B, Abbah J, Vongtau H, Gamaniel K (2003) Studies on the use of *Cassia singueana* in malaria ethnopharmacy. *Journal of Ethnopharmacology* 88: 261–267.
- Agbaje EO, Ajidahun OA (2011) Analgesic, anti-inflammatory and antipyretic effects of dried root ethanolic extract of *Strophanthus sarmentosus* (*apocynaceae*). *International Research Journal of Pharmacy and Pharmacology* 1: 62–69.
- Akpan EJ, Okokon JE, Etuk IC (2012) Antiplasmodial and antipyretic studies on root extracts of *Anthocleista djalensis* against *Plasmodium berghei*. *Asian Pacific Journal of Tropical Diseases* 2: 36–42.
- Alli LA, Adesokan AA, Salawu OA, Akanji MA, Tijani AY (2011) Anti-plasmodial activity of aqueous root extract of *Acacia nilotica*. *African Journal of Biochemistry Research* 5: 214–219
- Backhouse N, Delporte C, Givernan M, Cassels BK, Valenzuela A, Speisky H (1994) Anti-inflammatory and antipyretic effects of boldine. *Inflammation Research* 42: 114–117.
- Banso A (2009) Phytochemical and antibacterial investigation of bark extracts of *Acacia nilotica*. *Journal of Medicinal Plants Research* 3: 082–085.
- Bargal K, Bargali SS (2009) *Acacia nilotica*: a multipurpose leguminous plant. *Nature and Science* 7:11–19.
- Bittar M, de Souza MM, Yunes RA, Lento R, Delle MF, Cechinel FV (2000) Anti nociceptive activity of 13,118-binarigenin, a biflavonoid present in plants of the *guttiferae*. *Planta Medica* 66: 84–86.

- Calixto JB, Beirith A, Ferreira J, Santos AR, Cechinel FV, Yunes RA (2000) Naturally Occurring antinociceptive substances from plant. *Phytotherapy Research* 37: 319–321.
- Chang CP, Huang WT, Cheng BC, Hsu CC, Lin MT (2007). The flavonoid baicalin protects against cerebrovascular dysfunction and brain inflammation in experimental heatstroke. *Neuropharmacology* 52: 1024–1033.
- Cheng L, Ming-Liang H, Lars B (2005) Is COX-2 a perpetrator or a protector? Selective COX-2 inhibitors remain controversial. *Acta Pharmacologica sinica*. 26: 926–933.
- Chou R, Huffman LH, American Pain Society, American College of Physicians (2007) Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians Clinical Practice guideline. *Annals of Internal Medicine* 147: 505–514.
- Cooper KE (1995) Fever and antipyresis: role of the nervous system. 1<sup>st</sup> ed. Cambridge. Cambridge University press, 100-126.
- Courade JP, Besse D, Delchambre C (2001) Acetaminophen distribution in the central nervous system. *Life sciences* 69: 1455–1464.
- Gupta M, Mazumder UK, Sambath-Kumar P, Gomathi P, Rajeshwar Y, Kakoti BB, Tamilselven V (2005). Anti-inflammatory, analgesic and antipyretic effects of methanolic extract from *Bauhinia racemosa* stem bark in animal models. *Journal of Ethnopharmacology* 98: 267–273.
- Hayfaa AA, Sahar AA, Awatif MA (2013) Evaluation of analgesic activity and toxicity of alkaloids in *myristica fragrans* seeds in mice. *Journal of Pain Research* 2013: 611–615.
- Hosseinzadeh H, Ramezani M, Salman G (2000) Anti-nociceptive, anti-inflammatory and acute toxicity effects of *Zataria multiflora* Boiss extracts in mice and rats. *Journal of Ethnopharmacology* 73: 379–385.
- Jansen PAJ, Jagenav A (1959) A new series of potent analgesics, Dextro-2,2-diphenyl-3-methyl-4-morpholinobutyl pyrolidine and related amides. *Journal of Pharmacy and Pharmacology* 9: 381–385.
- Kanagusuku M, Sbars D, Bastos ES, Desouza MM, Cechinol-Filho V, Yunes RA, Delle Monache F, Niero R (2007) Pytochemical and analgesic activity of extract fractions and a 19-hydroxyursane-type triterpenoid obtained from *Rubus rosaefolius* (Rosaceae). *Biological and Pharmaceutical Bulletin* 30: 999–1002.
- Loke YK (2009) Aspirin or paracetamol- what is good for you? *Journal of Clinical Pharmacy and Therapeutics* 34: 247–248.
- Mustafa F, Indurkar J, Ismail S, Mordi MN, Ramanathan S, Mansor SM (2010). Analgesic activity, toxicity study and phytochemical screening of standardized *Cinnomomum iners* leaves methanolic extract. *Pharmacognosy Research* 2: 76–81.
- National Academy of Sciences (NAS) guide for care and use of laboratory animals. National Academy Press, 4-6. Available online; <http://www.nap.edu/catalog/5140.html>.
- Neha S, Ranvir GD, Jangade CR (2009) Analgesic and antipyretic activities of *Curcuma longa* rhizome extract in wistar rats. *Veterinary World* 2: 304–306.
- Oladosu P, Samuel BB, Okhale SE, Ibrahim K, Okogun JI (2007) Antitubercular activity of the dried fruits of *Acacia nilotica*. *Journal of Phytomedicine and Therapeutics* 12: 76–79.
- Organization for Economic Cooperation and Development (OECD) guideline 423. (2001). Oral Toxicity Study in Rodents. OECD guideline for the testing of chemicals, 423: 1–8.
- Owoyele BV, Nafiu AB, Oyewole IA, Oyewole LA, Soladoye AO (2009) Studies on the analgesic, anti-inflammatory and anti-pyretic effects of *Parquetina nigrescens* leaf extract. *Journal of Ethnopharmacology* 122: 86–90.
- Owoyele BV, Oguntoye SO, Dare K, Ogunbiyi BA, Arubuola EA, Soladoye AO (2008) Analgesic, anti-inflammatory and anti-pyretic activities from flavonoid fractions of *Chromolaena odorata*. *Journal of Medicinal Plants Research*. 2: 219–225.
- Padhan AR, Aghraham AK, Meher A (2010) A study on anti-pyretic activity of *Capparis zeylanica* Linn. Plant methanolic extract. *International Journal of Pharmaceutical Sciences and Research* 1: 169–171.
- Picq M, Chear SL, Prigent AF (1991) Effect of two flavonoid compounds in central nervous system. Analgesic activity. *Life Science* 49: 1979–1988.
- Prado WA, Tonussi CR, Rego EM, Corrado AP (1990) Antinociception induced by intraperitoneal injection of gentamycin in rats and mice. *Pain* 41: 365–371.
- Raffa R (2006) Pharmacological aspects of successful long-term analgesia. *Clinical Rheumatology* 25 (Suppl. 1) S9–S15.
- Rang HP, Dale MM, Ritter JM, Moore PK (2007) *Pharmacology*, 6<sup>th</sup> ed. Edinburgh: Churchill Livingstone.
- Rejón-Orantes JC, Perdomo-Suaréz DP, Rejón-Rodríguez A, Hernández-Hernández S, García-Liévana OE, López-Rodríguez D, Pérez de la Mora M (2013) Aqueous root extracts from *Mimosa albida* Humb. and Bonpl. ex Willd display antinociceptive activity in mice. *Journal of Ethnopharmacology*. 149: 522–526.

Salawu OA, Chindo BA, Tijani AY, Adzu B (2008) Analgesic, anti-inflammatory, antipyretic and antiplasmodial effects of the methanol extract of *Crossopteryx febrifuga*. *Journal of Medicinal Plants Research* 2: 213–218.

Singh N, Khatri P, Samanthar KC, Damor R (2010) Antipyretic activity of methanolic extract of leaves of *Quisqualis indica* Linn. *International Journal of Pharmaceutical Research and Development* 2: 122–128.

Sofowora A (1993) Medicinal Plants and Traditional Medicine in Africa. Spectrum Books Limited, Ibadan, Nigeria. 151–153.

Spacer CB, Breder CD (1994) The neurologic basis of fever. *New England Journal of Medicine* 330:1880–1886.

Van Wyk P (2000) Field Guide to trees of Southern Africa. Struik (Cape Town) 2: 25–27.

Woolf CJ (2010) What is this thing called pain? *Journal of Clinical Investigation* 120: 3742–3744.

Zhang W, Doherty M, Leeb BF (2007) EULAR evidence-based recommendation for the management of hand osteoarthritis: report of a taskforce of the EULAR Standing Committee for International Clinical Studies Including Therapeutics (ESCISIT). *Annals of the Rheumatic diseases* 66: 377–388.