



# Biokemistri

An International Journal of the Nigerian Society for Experimental Biology

## Original Article

# Honey increases sperm count in male albino rats by enhancing testosterone production.

Toyin Mohammed Salman<sup>1\*</sup>, Isiaka Abdullateef Alagbonsi<sup>2</sup>, Lukman Aribidesi Olayaki<sup>1</sup>, Sikiru Ayobami Biliaminu<sup>3</sup>, Hussein Mofomosara Salahdeen<sup>4</sup>, Olumide Ayodeji Olowu<sup>1</sup>

<sup>1</sup>Department of Physiology and <sup>3</sup>Chemical Pathology, College of Health Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria.

<sup>2</sup> Department of Physiology, Faculty of Medicine, Kogi State University, Anyigba, Kogi State, Nigeria.

<sup>4</sup> Department of Physiology, Lagos State University College of Medicine, Ikeja, Lagos State, Nigeria.

\*Corresponding Author: Toyin M. Salman. E-mail: drsalman111@yahoo.com; Tel.: +234 8032337422

Received: 02 May 2013; Revised: 03 July 2013; Accepted: 04 July 2013

**ABSTRACT:** We investigated the effects of different doses of honey (H) and testosterone (T) on sperm count and reproductive hormones in male albino rats. Thirty-five male albino rats were randomly divided in a blinded fashion into 7 groups of 5 rats each. Group 1 (control) was given 0.2 ml of distilled water. Groups 2, 3 and 4 were given 100 mg/kg, 200 mg/kg and 400 mg/kg of H orally respectively. Groups 5, 6 and 7 were given 2.5 mg/kg, 5 mg/kg and 7.5 mg/kg of T intraperitoneally respectively. All doses of H significantly ( $P < 0.05$ ) increased sperm count in rats, while all the doses of T significantly reduced sperm count in rats. Plasma T was increased and FSH was reduced ( $P < 0.001$ ) by all the three doses of H. On the contrary, LH was significantly reduced ( $P < 0.05$ ) by 100 mg/kg and 200 mg/kg of H but not by 400 mg/kg of H. All the three doses of T reduced the plasma T and LH in rats. Lastly, 5 mg/kg and 7.5 mg/kg but not 2.5 mg/kg of T reduced FSH in rats. The results suggest that honey enhanced sperm count in rat by increasing testosterone production.

**KEYWORDS:** Gonadotropins, Honey, Sperm count, Testosterone.

BKM.2013.011 © 2013 Nigerian Society for Experimental Biology; All rights reserved. Printed in Nigeria  
This article is downloadable online in PDF format at <http://www.bioline.org.br/bk>

## INTRODUCTION

Honey is a natural hive product with an extensive history of traditional human medicinal use in a large number of societies. It is widely available in most communities, and its therapeutic potential is grossly underutilized. The mechanism of action of several of its properties remain obscure, thus there is a need for further investigation (Zumla and Lulat, 1989).

Studies have shown that honey minimizes cellular injuries of the skin and post-radiotherapies mucosal trauma. Moreover, it possesses some biological properties such as antioxidant, antimicrobial, anti-inflammatory and immunomodulatory effects (Mohamed *et al.*, 2010; Estevinho *et al.*, 2008; Prakash *et al.*, 2008; Timm *et al.*, 2008; Perez *et al.*, 2006). The natural antioxidants, especially flavonoids, exhibit a wide range of biological effects, including antibacterial, anti-inflammatory, anti-allergic, anti-thrombotic, and vasodilatory actions (Cook and Sammon, 1996). Other reported properties of honey include, anti-platelet and anti-nociceptive (Kamran

*et al.*, 2006), prophylaxis against biofilm formation (Irish *et al.*, 2006), an aid or remedy to manage diversity of wound aetiologies (Gethin and Cowman, 2005) and dressing (Rahal *et al.*, 1984; Chirife *et al.*, 1983; Knutson *et al.*, 1981), dyspepsia and peptic ulcers (Al Somai *et al.*, 1994) and as a preservative for herbal medicines (Molan, 1998).

Honey is a natural product with very complex chemical composition. The composition of a particular honey sample greatly depends on the composition of nectar, where it originates. It contains more than 180 substances (White, 1979) including moisture; sugars such as glucose and fructose; enzymes such as catalase and glutathione reductase; trace essential elements such as iron, copper, zinc and calcium; vitamins such as vitamin A, C and E as well as some flavonoids and phenolic acids (Michalkiewicz *et al.*, 2008; Yao *et al.*, 2004; Al-Waili, 2003; Gheldof *et al.*, 2002). It is composed primarily of fructose and glucose but also contain 4-5% of fructooligosaccharide which serves as prebiotic agent (Chow, 2002). Flavonoids of antibacterial activity including pinocembrin (Bogdanov, 1984; Villanueva *et*

*al.*, 1970), kaempferol and quercetin, as well as naringenin and pinocembrin were detected in sunflower honey (Sabatier *et al.*, 1989). The presence of galangin and chrysin in several Swiss honeys has also been reported (Bogdanov, 1989). Thus, the composition of honeys varies with different floral sources as well as climatic and environmental conditions (Perez *et al.*, 2007; Gheldof *et al.*, 2002).

Traditionally, the local Malaysian and other population consume honey as a nutrient as well as for enhancement of fertility. Honey is farmed and used all over Nigeria. Initially local farmers harvested the honey from the wild but today apiculture is a growing industry in many parts of the country. Some studies on the healing effects and antimicrobial activity of honey collected from Nigeria on burns and wounds have been reported (Adesunkanmi and Oyelami, 1994). However, there is a paucity of data on the effects of honey collected from Nigeria on reproductive functions. The decline in male reproductive health and fertility in the last 30 years has been linked to environmental toxicants or xenobiotics (Sikka, 2008). This has led to increased interest to investigate the possible beneficial effect of honey in enhancing fertility in males.

Studies on the effect of honey on reproductive parameters are few and inconclusive. In a study using honey as vehicle for tamoxifen, a nonsteroidal antiestrogenic drug, honey was reported to increase sperm count without affecting other seminal parameters and reproductive hormones in male bonnet monkey (Gill-Sharma *et al.*, 2003). A recent study showed that treatment with 5% solution of honey collected from Palestine for 20 days orally to adult male rats increased epididymal sperm count and testicular sorbitol dehydrogenase activity as well as reduced lactate dehydrogenase activity (Abdul-Ghani *et al.*, 2008). Moreover, honey collected from Malaysia has recently been shown to increase sperm count without affecting other semen parameters and reproductive hormones in male rat (Mohamed *et al.*, 2012). The mechanism underlying an increase in sperm count without any change in reproductive hormones is not clear and of interest to us. The observation of Mohamed *et al.* (2012) therefore raises a question on the possibility of an increase in sperm count without any change in reproductive hormones. Testosterone is needed for the growth and development of male reproductive organs (Mooradan *et al.*, 1987) and in association with follicular stimulating hormone, it acts on the seminiferous tubules to initiate and maintain spermatogenesis (Christensen, 1975).

The study reported here was designed to evaluate the effects of different doses of honey on sperm count, testosterone and gonadotropins in male albino rats. To investigate whether the spermatogenic effect of honey is testosterone-related, we also studied the effect of different doses of testosterone on sperm count, testosterone level and gonadotropins in rats.

## MATERIALS AND METHODS

### Honey sample

The honey used in this study was obtained from the Federal College of Forestry, Ibadan. It was freshly diluted daily to the doses required using distilled water (as a vehicle).

### Animals and treatment protocol

Thirty-five male albino rats (200-250 g) were used for the study. The animals were obtained from the animal house of the Department of Physiology, College of Health Sciences, University of Ilorin, Kwara State, Nigeria. They were housed at five per cage and provided with standard laboratory feed and water *ad libitum*. They were maintained in a well-ventilated room at  $25 \text{ }^{\circ}\text{C} \pm 2 \text{ }^{\circ}\text{C}$  on a 12-hour light/dark cycle. Study protocol and animal use were approved, prior to the beginning of the study, by our institutional research and ethical committee. All necessary protocols were followed to ensure the humane treatment of the animals.

The animals were randomly divided in a blinded fashion into 7 groups (5 rats per group). Group 1 (control) was given 0.2 ml of distilled water. Groups 2, 3 and 4 were given 100 mg/kg, 200 mg/kg and 400 mg/kg of honey respectively by oral gavage once daily for 4 weeks. Groups 5, 6 and 7 were given 2.5 mg/kg, 5 mg/kg and 7.5 mg/kg of Testosterone (T) (Laborate Pharmaceutical, India) intraperitoneally respectively once daily for 4 weeks. At the end of the study (24 h after the final treatment), laparotomy was performed under ether anaesthesia. Epididymis was removed and blood was collected into sample bottles and allowed to clot at room temperature and sera separated by centrifugation at 1000 rpm for 10 min and stored at  $-20 \text{ }^{\circ}\text{C}$  for hormone estimations.

### Estimation of sperm count

The testes from each rat were carefully exposed and one of them was removed together with its epididymis. For each separated epididymis, the caudal part was removed and placed in a beaker containing 1 ml of normal saline solution. It was macerated with a pair of sharp scissors and left for few minutes to liberate the sperm cells into the normal saline. Semen drops were placed on a clean grease-free glass slide and two drops of warm 2.9% sodium citrate were added. The improved Neubauer counting chamber was charged with the semen solution and the number of sperm cells, appearing as black dots were counted in 25 small squares within the central counting area of the counting chamber as earlier described (Cheesebrough, 2000).

### Estimation of reproductive hormones

Serum-free testosterone and gonadotrophins (luteinizing hormone and follicle-stimulating hormone) were measured by tube-based enzyme immunoassay (EIA) method (Raji *et al.*, 2005) using commercial kits (IBL-Hamburg GmbH, Germany). The EIA is a standardized method used by WHO and part of its program for human reproduction research. The procedures for the assay as contained in the manufacturer's manual were strictly followed. The within assay variation was 8.1% and the sensitivity was 0.3 ng/ml. The optical density was read using a spectrophotometer (Jenway, 6300 spectrophotometer, UK) that was sensitive at wavelengths between 492 nm and 550 nm.


### Statistical Analysis

Data were analysed using SPSS version 16.0 for windows. All values given were the mean  $\pm$  S.E.M of the variables measured. Significance was assessed by the one-way analysis of variance (ANOVA), followed by a post-hoc Turkey multiple range test for multiple comparisons. P-Values of 0.05 or less were taken as statistically significant.

## RESULTS

### Effects of different doses of honey on sperm count in male albino rats

Effects of different doses of honey (H) on sperm count in male albino rats are shown in Figure 1. Oral administration of the three different doses of H significantly ( $P < 0.05$ ) increased sperm count in rats. The increases in sperm count were not dose-dependent as there was no significant change when the values of sperm count for the different doses were compared.



**Figure 1: Effects of honey and testosterone administration on sperm count in male rats. H=Honey, T=Testosterone. Values are expressed as Mean  $\pm$  S.E.M (n=5).  $P < 0.05$ , \*\* $P < 0.01$  vs control.**

**Table 1: Effects of honey (H) and testosterone (T) administration on plasma T, Follicle Stimulating Hormone (FSH) and Luteinizing Hormone (LH) in male rats. Values are expressed as Mean  $\pm$  S.E.M (n=5).  $P < 0.05$ , \* $P < 0.01$ , \*\*\* $P < 0.001$  vs control.**

	T (ng/ml)	LH (ng/ml)	FSH (ng/ml)
Control	3.76 $\pm$ 0.05	2.75 $\pm$ 0.16	8.26 $\pm$ 0.52
100 mg/kg (H)	6.18 $\pm$ 1.50***	2.57 $\pm$ 0.05*	0.13 $\pm$ 0.04***
200 mg/kg (H)	5.73 $\pm$ 0.79***	2.27 $\pm$ 0.07*	0.41 $\pm$ 0.07***
400 mg/kg (H)	8.37 $\pm$ 0.89***	2.78 $\pm$ 1.38	0.46 $\pm$ 0.13***
2.5 mg/kg (T)	2.88 $\pm$ 0.1***	2.06 $\pm$ 0.21*	6.84 $\pm$ 0.66
5 mg/kg (T)	2.02 $\pm$ 0.1***	1.58 $\pm$ 0.2**	4.66 $\pm$ 0.35***
7.5 mg/kg (T)	1.46 $\pm$ 0.14***	1.14 $\pm$ 0.13***	3.92 $\pm$ 1.03**

### Effects of different doses of testosterone on sperm count in male albino rats.

Effects of different doses of testosterone (T) on sperm count in male albino rats are shown in Figure 1. Oral administration of the three different doses of T caused significant reduction in sperm count in rats. The reduction was more significant ( $P < 0.01$ ) at 2.5 mg/kg and 7.5 mg/kg than 5 mg/kg ( $P < 0.05$ ).

### Effects of honey on plasma testosterone and gonadotropins levels in male albino rats

The effects of different doses of H on T, Follicle Stimulating Hormone (FSH) and Luteinizing Hormone (LH) in male albino rats are shown in Table 1. Plasma T was significantly increased ( $P < 0.001$ ) by oral administration of all the three doses of H. On the contrary, plasma LH was significantly reduced ( $P < 0.05$ ) by oral administration of 100 mg/kg and 200 mg/kg of H but not by 400 mg/kg of H. Moreover, plasma FSH was significantly reduced ( $P < 0.001$ ) by all the three doses of H.

### Effects of testosterone administration on plasma testosterone and gonadotropins levels in male albino rats

The effects of different doses of T on plasma T, FSH and LH in male albino rats are shown in Table 1. All the three doses of T significantly ( $P < 0.001$ ) reduced the plasma testosterone in rats. In addition, plasma LH was also significantly reduced by 2.5 mg/kg ( $P < 0.05$ ), 5 mg/kg ( $P < 0.01$ ) and 7.5 mg/kg (0.001) of T. Lastly, 5 mg/kg and 7.5 mg/kg but not 2.5 mg/kg of T significantly reduced ( $P < 0.001$ ,  $P < 0.01$ ,  $P > 0.05$  respectively) the plasma FSH in rats.

## DISCUSSION

Previous studies have reported that honey collected from Palestine and Malaysia increases sperm count in rat and monkey (Mohamed *et al.*, 2012; Abdul-Ghani *et al.*, 2008; Gill-Sharma *et al.*, 2003). The observed increase in sperm count following administration of honey in the present study is consistent with these previous reports. However, critical observation of the data of Mohamed *et al.* (2012) may suggest a contradictory finding as they reported that 200mg/kg of honey obtained from Malaysia did not significantly increase the sperm count in rat, whereas it did in our own study with honey sourced from Nigeria.

Previous studies using honey that originated from Malaysia or Palestine also reported no significant effects of honey on reproductive hormones including testosterone (T), luteinizing hormone (LH) and follicle-stimulating hormone (FSH) in rat (Mohamed *et al.*, 2012) and monkey (Gill-Sharma *et al.*, 2003). The present study however showed that honey from Nigeria increased T but decreased LH and FSH in rat. The content and composition of honeys vary with different floral sources as well as climatic and environmental conditions (Perez *et al.*, 2007; Gheldof *et al.*, 2002). The variations between the present study and those previous studies may be a result of the difference in the source, and of climatic and environmental conditions of the honey used for the studies. In addition, we used wistar (albino) rats in this study; Mohamed *et al.* (2012) used Sprague-Dawley rats while Gill-Sharma *et al.* (2003) used bonnet monkey. We speculate that species difference could be another cause of the discrepancy in the findings.

The significant effect of honey on the reproductive hormones in this study is of great interest. It has been suggested that honey from Palestine enhances epididymal sperm count by possibly affecting the key enzymes in spermatogenesis and sperm maturation such as increases sorbitol dehydrogenase activity and reduces lactate dehydrogenase activity (Abdul-

Ghani *et al.*, 2008). However, we were able to establish another fact that honey increases sperm count by increasing the testicular production of testosterone, which was evident from the elevated plasma testosterone following treatment with all doses of honey. Expectedly, the increased plasma testosterone caused reduction in plasma LH and FSH through the previously well-established negative feedback mechanism (Kellis and Vickery, 1984; Prasad *et al.*, 1996; Meeuwen *et al.*, 2007; Koehler *et al.*, 2009).

To support our hypothesis that the sperm count boosting potential of honey is testosterone dependent, we studied the effect of exogenous administration of different doses of testosterone. The observed reduction in sperm count, testosterone and gonadotropins following exogenous administration of testosterone in this study is comparable with previous reports (Awoniyi *et al.*, 1989; 1992; Airkin *et al.*, 1989; Sun *et al.*, 1989; Sharpe *et al.*, 1988; Robaire *et al.*, 1979). Moreover, testosterone withdrawal has also been shown to cause spermatogenic cell degeneration (Kerr *et al.*, 1993). Though we did not measure the testicular testosterone and gonadotropins levels, we speculate that reduction in testicular testosterone level following its exogenous administration could have led to reduction in spermatogenesis, and subsequently low sperm count.

A study on the chemical and physical characterization of honey sourced from different sources within Nigeria showed that it is quite rich in minerals such as K, Ca, Ti, Cr, Mn, Fe, Ni, Cu, Zn, Se, Br and Rb, as well as carboxylic acids, aldehydes, alkynes, nitrites, alkenes and ethers (Adebiji *et al.*, 2004). Recently, Nurul Syazana *et al.* (2013) and Odeh *et al.* (2007) respectively identified 35 and 30 volatile compounds in honey collected from Malaysia and Pakistan. These compounds include acids, aldehydes, alcohol, ketones, terpenes, hydrocarbons, furans, etc. However, the specific component(s) of the honey used in this study that exerted effects different from those sampled from Malaysia or Palestine (Gill-Sharma *et al.*, 2003; Abdul-Ghani *et al.*, 2008; Mohamed *et al.*, 2012) is a subject of interest and further investigation. Better knowledge of this will shed more light into the specific testosterone-dependent mechanism by which honey increases sperm count.

However, it is noteworthy that chrysin (5,7-dihydroxyflavone), a natural flavonoid has been reported to be present at high levels in honey, propolis and many plant extracts (Wang and Morris, 2007). The study of Ciftci *et al.* (2012) clearly showed that treatment with chrysin significantly increased serum testosterone levels in rats. Similarly, Jana *et al.* (2008) found that testosterone production was dramatically enhanced in primary cultures utilising Leydig cells isolated from mouse testis when the cells were treated with chrysin. Other investigators have also drawn attention that aromatase inhibition by chrysin could block the conversion of androgens into oestrogens with a consequent increase in testosterone (Jeong, 1999; Le Bail, 1998; Kellis and Vickery, 1984). However, whether or not chrysin or any other agent (s) capable of boosting testosterone level is present in the honey samples used in this study is not known. Further phytochemical analysis of honey to reveal the active ingredient (s) that is/are capable of boosting testosterone will be worthwhile.

## REFERENCES

- Abdul-Ghani AS, Dabdoub N, Muhammad R, Abdul-Ghani R and Qazzaz M (2008) Effect of Palestinian honey on spermatogenesis in rats. *Journal of Medicinal Food* 11: 799–802.
- Adebiji FM, Akpan I, Obiajunwa EI and Olaniyi HB (2004) Chemical/Physical Characterization of Nigerian Honey. *Pakistan Journal of Nutrition* 3: 278–281.
- Adesunkanmi K and Oyelami OA (1994) The pattern and outcome of burn injuries at Wesley Guild Hospital, Ilesha, Nigeria: A review of 156 cases. *Journal of Tropical Medicine and Hygiene* 97: 108–112.
- Al Somai N, Coley KE, Molan PC and Hancock BM (1994) Susceptibility of *Helicobacter pylori* to the Antibacterial Activity of Manuka Honey. *Journal of Royal Society of Medicine* 87: 9–12.
- Al-Waili NS (2003) Effects of daily consumption of honey solution on hematological indices and blood levels of minerals and enzymes in normal individuals. *Journal of Medicinal Food* 6:135–140.
- Awoniyi CA, Santulli R, Sprando RL, Ewing LL and Zirkin BR (1989) Restoration of advanced spermatogenic cells in the experimentally regressed rat testis: quantitative relationship of testosterone concentration within the testis. *Endocrinology* 124:1217–1223.
- Awoniyi CA, Zirkin BR, Chandrashekar V and Schlaff WD (1992) Exogenously administered testosterone maintains spermatogenesis quantitatively in adult rats actively immunized against gonadotropin-releasing hormone. *Endocrinology* 130: 3283–3288.
- Bogdanov S (1984) Characterization of antibacterial substances in honey. *Lebensmittel-Wissenschaft und Technologie* 17: 74–76.
- Bogdanov S (1989) Determination of pinocembrin in honey by using HPLC. *Journal of Apicultural Research* 28: 55–57.
- Cheesebrough M (2000) Examination of semen. Microbiological tests. In: District laboratory practice in tropical countries, part 2, Cambridge University Press, United Kingdom. pp.130–132.
- Chirife J, Herszage L, Joseph A and Koh ES (1983) *In vitro* study of bacterial growth inhibition in concentrated sugar solutions: microbiological basis for the use of sugar in treating infected wounds. *Antimicrobial Agents and Chemotherapy* 23: 766–773.
- Chow J (2002) Probiotics and prebiotic: A brief overview. *Journal of Renal Nutrition* 12: 76–86.
- Christensen AC (1975) Leydig Cell: In: Handbook of Physiology, P. O. Greep, E. B. Astwoods (eds.). American Physiological Society, Washington D.C.
- Ciftci O, Ozdemir I, Aydin M, Beytur A (2012) Beneficial effects of chrysin on the reproductive system of adult male rats. *Andrologia* 44: 181–186.
- Cook NC, Sammon S (1996) Flavonoids: chemistry, metabolism, cardioprotective effects, and dietary sources. *The Journal of Nutritional Biochemistry* 7: 66–76.

- Estevinho L, Pereira AP, Moreira L, Dias LG and Pereira E (2008) Antioxidant and antimicrobial effects of phenolic compounds extracts of Northeast Portugal honey. *Food Chemistry and Toxicology* 46:3774–3779.
- Gethin G and Cowman S (2005) Case series of use of Manuka honey in leg ulceration. *International Wound Journal* 2: 10–15.
- Gheldof N, Wang XH and Engeseth NJ (2002) Identification and quantification of antioxidant components of honeys from various floral sources. *Journal of Agricultural and Food Chemistry* 50: 5870–5877.
- Gill-Sharma MK, D'Souza S, Parte P, Balasinor N, Choudhuri J, Majramkar DD, Aleem M and Juneja HS (2003) Effect of oral tamoxifen on semen characteristics and serum hormone profile in male bonnet monkeys. *Contraception* 67: 409–413.
- Irish J, Carter D and Blair S (2006) Honey prevents biofilm formation in microbial pathogens. Oral presentation. Proceeding of the 1st international conference on the medicinal uses of honey. USM Kelantan.
- Jana K, Yin X, Schiffer RB, Chen JJ, Pandey AK, Stocco DM, Grammas P and Wang X (2008) Chrysin, a natural flavonoid enhances steroidogenesis and steroidogenic acute regulatory protein gene expression in mouse Leydig cells. *Journal of Endocrinology* 197: 315–323.
- Jeong HJ (1999) Inhibition of aromatase activity by flavonoids. *Archive of Pharmaceutical Research* 22: 309–312.
- Kamran AM, Ahmed MM, Shabana S, Auranzeb M, Nazimuddi A and Saeed S (2006) Medicinal properties of honey from northern Pakistan. Oral presentation. Proceeding of the 1<sup>st</sup> international conference on the medicinal uses of honey. USM Kelantan.
- Kellis Jr. JT and Vickery LE (1984) Inhibition of human estrogen synthetase (aromatase) by flavones. *Science* 225: 1032–1034.
- Kerr JB, Millar M, Maddocks S and Sharpe RM (1993) Stage-dependent changes in spermatogenesis and sertoli cells in relation to the onset of spermatogenic failure following withdrawal of testosterone. *Anatomical Record* 235: 547–559.
- Knutson RA, Merbit LA, Creekmore MA and Snipes HG (1981) Use of Sugar and Povidone-iodine to Enhance Wound Healing: Five Years Experience. *Southern Medical Journal* 74: 1329–1335.
- Koehler K, Parr MK, Geyer H, Mester J and Schanzer W (2009) Serum testosterone and urinary excretion of steroid hormone metabolites after administration of high-dose zinc supplement. *European Journal of Clinical Nutrition* 63: 65–70.
- Le Bail JC (1998) Aromatase and 17 beta-hydroxysteroid dehydrogenase inhibition by flavonoids. *Cancer Letter* 133: 101–106.
- Meeuwen JA, Korthagen N, de Jong PC, Piersma AH and Van den Bergh M (2007) (Anti) estrogenic effects of phytochemicals on human primary mammary fibroblasts, MCF-7 cells and their co-culture. *Toxicological and Applied Pharmacology* 221: 372–383.
- Michalkiewicz A, Biesaga M and Pyszynska K (2008) Solid-phase extraction procedure for determination of phenolic acids and some flavonols in honey. *Journal of Chromatography A* 1187:18–24.
- Mohamed M, Sirajudeen KNS, Swamy M, Yaacob NS and Sulaiman SA (2010) Studies on the antioxidant properties of Tualang honey of Malaysia. *African Journal of Traditional Complementary and Alternative Medicine* 7:59–63.
- Mohamed M, Sulaiman SA, Jaafar H and Sirajudeen KNS (2012) Effect of different doses of Malaysian honey on reproductive parameters in adult male rats. *Andrologia* 44: 182–186.
- Molan PC (1998) Honey as a Dressing for Wounds, Burns, and Ulcers: A Brief Review of Clinical Reports and Experimental Studies Primary Intention 6 (4). *Spectrometry Journal of AOAC International* 89: 586-593.
- Mooradan AD, Morley JE and Koreman SG (1987) Biological actions of androgens. *Endocrine Review* 8: 1–28.
- Nurul Syazana MS, Gan SH and Halim AS (2013) Analysis of volatile compounds of Malaysian taulag (Koompassia excelsa) honey using gas chromatography mass spectrometry. *African Journal of Traditional Complementary and Alternative Medicine* 10: 180–188.
- Odeh I, Abu-Lafi S, Dewik H, Al-Najjar I, Imam A, Valery MO and Hanus DL (2007) A variety of volatile compounds as markers in Palestinian honey from Thymus capitatus, Thymelaea hirsuta, and Tolpis virgata. *Food Chemistry* 101: 1393–1397.
- Perez E, Rodriguez-Malaver AJ and Vit P (2006) Antioxidant capacity of Venezuelan honey in Wistar rat homogenates. *Journal of Medicinal Food* 9:510–516.
- Perez RA, Iglesias MT, Pueyo E, Gonzalez M and de Lorenzo C (2007) Amino acid composition and antioxidant capacity of Spanish honeys. *Journal of Agricultural and Food Chemistry* 55:360–365.
- Prakash A, Medhi B, Avti PK, Saikia UN, Pandhi P and Khanduja KL (2008) Effect of different doses of Manuka honey in experimentally induced inflammatory bowel disease in rats. *Phytotherapy Research* 22:1511–1519.
- Prasad AS, Mantzoros CS, Beck BW, Hess JW and Brewer GJ (1996) Zinc status and serum testosterone levels of healthy adults. *Nutrition* 12: 344–348.
- Rahal F, Mimica IM, Pereira V and Athié E (1984) Sugar in the Treatment of Infected Surgical Wounds. *International Surgery* 69: 308.
- Raji Y, Salman TM and Akinsomisoye OS (2005) Antispermatic activity of Morinda lucida extract in male rats. *Asian Journal of Andrology* 7: 405–410.
- Robaire B, Ewing LL, Irby DC and Desjardins C (1979) Interactions of testosterone and estradiol-17P on the reproductive tract of the male rat. *Biology of Reproduction* 21:455–463.
- Sabatier S, Amiot MJ, Aubert S, Tacchini M and Gonnet M (1989) Importance des flavonoids dans les miels de tournesol. *Bulletine of Technology in Apiculture* 64: 171–178.

Sharpe RM, Donachie K and Cooper I (1988) Re-evaluation of the intratesticular level of testosterone required for quantitative maintenance of spermatogenesis in the rat. *Journal of Endocrinology* 117:19–26.

Sikka SC (2008) Wang R. Endocrine disruptors and estrogenic effects on male reproductive axis. *Asian Journal of Andrology* 10: 134–145.

Sun YT, Irby DC, Robertson DM and de Kretser DM (1989) The effects of exogenously administered testosterone on spermatogenesis in intact and hypophysectomized rats. *Endocrinology* 125:1000–1010.

Timm M, Bartelt S and Hansen EW (2008) Immunomodulatory effects of honey cannot be distinguished from endotoxin. *Cytokine* 42:113–120.

Villanueva VR, Barbier M, Gonnet M and Lavie P (1970) Les flavonoids de la propolis. Isolement d'une nouvelle substance bacteriostatique: la pinocembrin (dihydroxy-5, 7-lavanone). *Annales de l'Institut Pasteur* 118: 84–87.

Wang X and Morris ME (2007) Effects of the flavonoid chrysin on nitrofurantoin pharmacokinetics in rats: Potential involvement of ABCG2. *Drug Metabolism and Disposition*. 35:268–274.

White JW (1979) Composition of honey. In: *Honey: A Comprehensive Survey*. E. Crane (ed.), Heinemann, London. pp. 157–192.

Yao L, Jiang Y, D'Arcy B, Singanusong R, Datta N, Caffin N and Raymont K (2004) Quantitative high-performance liquid chromatography analyses of flavonoids in Australian Eucalyptus honeys. *Journal of Agricultural and Food Chemistry* 52: 210–214.

Zumla A and Lulat A (1989) Honey—a remedy rediscovered. *Journal of Royal Society of Medicine* 82: 384–385.