

## Studies on the Nutritional and Phytochemical Properties of *Persea americana* Seed

Nwaogu, L. A., Alisi, C. S. and Ojiako. O. A.

Department of Biochemistry, School of Science, Federal University of Technology, Owerri, Nigeria.

**Corresponding author:** Nwaogu, L. A. Department of Biochemistry, School of Science, Federal University of Technology, Owerri, Nigeria. Email: nwogulinus@yahoo.com

### Abstract

*The phytochemical, nutrient, antinutrient and mineral element composition of seed extract of Persea americana were investigated. Phytochemical analyses of the seed extract revealed the presence of alkaloids, saponins, tannins, flavonoids, cyanogenic glycosides but no anthracine glycosides. Proximate analysis gave 12.90 ± 1.57% moisture; 18.53 ± 0.26% crude fat; 18.55±1.26% crude protein; 47.35±24% carbohydrate; 2.6±0.63% ash; 3.17±0.17% crude fibre. The seed also was a rich source of mineral elements; containing calcium 12.30±0.08 mg; iron 0.307± 0.13mg; magnesium 21. 12±3.86 mg; phosphorus 46.00±1.72 mg; potassium 103.8±0.22 mg; sodium 0.302± 0.02 mg; and zinc 0.087±0.01mg per 100gm dry weight respectively. Quantitative analyses of the anti nutritional components revealed oxalate, phytate, saponins, tannins, flavonoids and cyanogenic glycoside contents of 27.25±4.10, 5.44±0.03, 33.23±1.73, 56.11±0.21, 1.90 ± 0.08 and 24.01± 0.25 mg/100gm dry weight respectively.*

**Keywords:** *Persea americana* seed, Phytochemical screening, Proximate analysis, Antinutritional components, Mineral content

### Introduction

*Persea americana* otherwise known as avocado pear has not attracted much attention especially the seed, despite the several claims credited to the plant by herbalists who used the leaves and seed for herbal preparations. *P. americana* is an evergreen tree, up to 20 metres high with dense, compact and whorled branches. It has been used in ethnomedicinal preparations in some African countries. The fruit skin which "is antibiotic in nature is employed as a vermifuge and also as a remedy for dysentery. The leaves are chewed as a remedy for diarrhoea. Boiled leaves of *Persea americana* are applied on the forehead to relieve neuralgia (Obioha, 1998). The seed when cut into pellets, roasted and ground are given to stop diarrhoea and dysentery. Oil extracted from the seed heals skin eruptions (Ezenwafor, 1992). Edible flesh of *Persea americana* serves as a very high energy source for diabetics due to its low sugar content. The seed could also be suitable for compounding poultry feed (Obioha, 1998).

Dietetically, the edible fleshy part of *Persea americana* is one of the most nutritious of all salad fruits with a fleshy, smooth, buttery pulp. In North America, it is primarily served as a salad vegetable and is merely halved and garnished with seasoning, lime juice, vinegar and other dressings. Often, the halves are stuffed with shrimps, crabs or other sea foods (Vermon, 1983).

Plants have been known to be useful in the formulation of feeds as feed supplements. The aim has been to improve the quality of some animal feeds. Soybean, maize and wheat for example are used in the formulation of standard poultry feed. This study is intended to assess the nutritional and toxicological compositions of *P. americana* seed with the view to evaluating its possible utilization in improving the quality of conventional commercial feeds. The importance of this effort is obvious

considering the high cost of animal feeds and the abundance and apparent zero economic value of this plant resource.

### Materials and Methods

#### Collection and preparation of plant materials:

The seeds of *Persea americana* were bought from Ihiagwa Market in Owerri West LGA of Imo State, Nigeria. The seeds were identified at the Department of Plant Science and Biotechnology of the Imo State University, Owerri, Nigeria. Clean healthy seeds were cut into pellets and sun-dried to constant weight before powdering the dried pellets using a domestic blender. Portions of the powdered sample were then used for the various analyses.

**Phytochemical analysis:** Quantitative phytochemical screening for the presence of alkaloids, saponins, flavonoids, tannins, cyanogenic glycosides, oxalate and phytate were carried out according to the methods described by Harbome (1973) and Trease and Evans (1983).

**Proximate analysis:** The proximate analysis of the seed extract for crude protein, crude fibre and fat contents were determined using the methods described by Pearson (1976). Crude protein determination was done using Kjeldhal's method, while crude fibre determination was done using Wende's method. Fat content was determined using continuous solvent extraction method. Total ash content was determined by furnace incineration using the method of James (1995). Moisture and carbohydrate contents were determined using the methods described by AOAC (2000).

**Elemental analysis:** The minerals: calcium, sodium, potassium, phosphorus, magnesium, iron

and zinc were performed using the Buck scientific atomic absorption/emission spectrophotometer 200A.

**Determination of antinutrients:** Phytate and oxalate contents were determined by the spectrophotometric method of Hang and Lantzsch (1983) while the cyanide content was estimated by the alkaline titration method (AOAC, 2000).

## Results and Discussion

Table 1 shows the proximate composition of the powder seed which revealed the following: protein content  $18.55 \pm 1.26\%$ , carbohydrate  $47.35 \pm 3.24\%$ , fats  $18.53 \pm 0.26\%$ , moisture content  $12.90 \pm 1.57\%$ , ash content  $2.67 \pm 0.63\%$  and crude fibre  $3.17 \pm 0.17\%$  respectively. Results also showed a calorific value of  $430.14+10.42$  kilocalories per gram dried seed powdered.

**Table 1: Proximate composition of *Persea americana* seed extract**

Nutrient composition	Mean Composition*
Crude protein	$18.55 \pm 1.26$
Carbohydrate	$47.35 \pm 3.24$
Fats	$18.53 \pm 0.26$
Ash	$2.67 \pm 0.63$
Moisture	$12.90 \pm 1.57$
Crude fibre	$3.17 \pm 0.17$

\*Values are means of three determinations  $\pm$  S.D

The seed powder although low in crude protein when compared to some leguminous seeds as reported by Boulter, (1997) could be a potential source of protein for livestock. The carbohydrate content of the seed powder is low compared to that reported by Ijeh *et al* (2004) from *Mucuna sloane* seed. The fat content of the seed powder is however higher than that determined from *Mucuna sloane* seed. The moisture content of the seed powder was found to be low indicating that the seed probably has a good keeping quality. This would agree with the reports of Ijeh *et al* (1998) and Ijeh *et al* (2004). The ash content of the powdered seed agrees with the report of Samson (1989). The crude fibre value from the results corroborates the results of Enechi (2001) who put the value as  $4.01 \pm 0.03\%$ .

**Table 2: Phytochemical composition of *Persea americana* seed extract**

Phytochemical constituents	Composition (mg/100g dry weight)*
Saponins	$33.33 \pm 1.73$
Tannins	$56.11 \pm 4.10$
Oxalates	$27.49 \pm 4.10$
Phytates	$5.44 \pm 0.05$
Cyanogenic glycosides	$24.01 \pm 0.25$
Flavonoids	$1.90 \pm 0.08$

\* Values are means of three Determinations  $\pm$  S.D

Quantitative phytochemical analyses (Table 2) revealed the presence of saponins, flavonoids and tannins. Saponins are known to have hypocholesterolemic activities (Price *et al*, 1989). This means that *Persea americana* seeds could

have some protection against coronary heart disease. Also, flavonoids which have been associated with free radical scavenging property as well as an anticarcinogenic potential (Trease and Evans, 1983) had a fairly low concentration ( $1.9\text{mg}/100\text{gm}$  dry weight) in the seed. This means that the powder can be useful in the management of physiological conditions associated with oxidative stress e.g. diabetes and hypertension.

The high concentration of the antinutritional factors phytate, oxalate and cyanogenic glycosides makes the seed to appear potentially toxic. The seed extract contains some micronutrients that are capable of complexing with these antinutritional factors and so make them unavailable. Calcium, for instance, complexes with oxalate to form calcium oxalate. Calcium also complexes with phytate and phosphorus to form calcium phytate and calcium phosphate (hydroxyapatite). Cyanogenic glycosides which are also known to be toxic are also found to be present in the extract. Although the levels of these antinutritional factors are fairly high, it is unlikely that they will pose any health problem as their values are drastically reduced by various processing methods including heating (Munro and Bassir, 1969; Enechi and Odonwodo, 2003).

Table 3 shows the mineral composition of the seed extract of *Persea americana* and reveals that it is a rich source of calcium, magnesium, potassium and phosphorus. It also contains small amounts of zinc, iron and sodium. These minerals play vital roles in diverse metabolic activities (Enechi, 2001). However, their bioavailability should be ascertained since a number of antinutritional factors which are capable of complexing with these useful macro and micronutrients (Enechi *et al*, 1996) thus reducing their bioavailability have also been found in the seed.

**Table 3: Mineral composition of *Persea americana* seed extract**

Mineral Element	Composition (mg/100gm dry weight)*
Calcium	$12.30 \pm 0.08$
Magnesium	$21.12 \pm .86$
Phosphorus	$46.00 \pm 7.20$
Potassium	$103.80 \pm 0.22$
Iron	$0.31 \pm 0.13$
Sodium	$0.30 \pm 0.20$
Zinc	$0.087 \pm 0.10$

\*Values are means of three determinations  $\pm$  S.D.

**Conclusion:** The study has shown that the seed extract of *Persea americana* is probably a good source of carbohydrate, protein and fat as well as some mineral elements including calcium, phosphorus, potassium and magnesium. It therefore means that if the seed is properly processed to reduce the antinutritive components, it can be incorporated into livestock feeds. It could therefore be of nutritional benefit to livestock in addition to its potential medicinal benefits.

The fact that this seed is currently discarded after taking the pulp of the fruit further underscores the relevance of our work and emphasises the need for more interest by researchers in diverse disciplines in this potentially useful but wasted resource.

## References

- AOAC (2000) Official Methods of Analysis of the Association of Analytical Chemists.
- Boulter, D. (1997) Quality Problems in Plant Proteins with Special Attention paid to the protein of Legumes In: *Protein quality from leguminous crops. Proceedings of Seminar in the European Economic Community Programme* organized by INRS Nov. 3,
- Edem, D. V; Amugo, C. I. and Eka, O. D. (1990) Chemical Composition of Yam Bean (*Splzenostylis stenocmpa*). *Tropical Science*, 30: 59 - 63.
- Enechi, O. C. (2001) *Basic Biochemistry of Food Nutrients*. (1<sup>st</sup> ed) Immaculate Publications Ltd Enugu, Nigeria.
- Enechi, O. C; Nwodo, O. F. C and Ogan, A. U. (1996) The Occurrence of Goitrogenic Flavoglycosides in the Roots of *Cissus populnea*. *Plant Prod. Res. Comm.*, 1: 27 - 31.
- Enechi, O. C. and Odonwodo I. (2003). An Assessment of the Phytochemical and Nutrient Composition of the Pulverized Root of *Cissus quadrangularis*. *Bio-Research*, 1: 63- 68.
- Ezenwafor, T. C. (1998) Determination of Active Ingredients 111 *Persea amencmw* seed. Chidimess Ltd Awka, Nigeria. pp 21 - 23.
- Hang, W. and Lantzsch, H.J. (1983) Comparative Methods for the Rapid Determination of Phytate in Cereal Product. *Journal of Science, Food and Agriculture*, 34(12): 1423 - 1426
- Harborne, I. B. (1973) *Phytochemical Methods. A Guide to Modern Techniques of Plant Analysis*. Chapman and Hall, New York. pp 88 - 185.
- Ijeh, I. I; Nwaugu, V. O. and Obidoa, O. (1998) Comparative Studies on the Nutritive Phytochemical and Antimicrobial Properties of two Varieties of *Vernol1ia amygdalil1a*. *Plant Prod. Res. Comm.* 1: 71 - 75
- Ijeh, I. I., Dnaegbo, S. O and Anaga, A.O. (2004) Studies on some Nutrition and Toxicological Properties of *Mucuna slogne* seed. *Bio-Research*, 2: 24 - 28
- James, C. S. (1995) *Analytical Chemistry of Foods*. Chapman and Hall, New York.
- Munro, A. and Bassir, O. (1969) Oxalate in Nigerian Vegetables. *Applied Chemistry*, 12:14-18
- Obioha, F. C. (1992). Breeds and breeding In: A guide to poultry production in the tropics. Alena Publisher, Enugu, Nigeria.
- Osborne D. R and Voogts. P (1978). *Calculation of Calorific value in the Analysis of Nutrients in Foods*. Academic Press, London.
- Pearson, D. (1976). *The Chemical Analysis of Foods*. Churchill Livingstone, Edinburgh.
- Price, K. R; Johnson, L. I. and Feriwick M. (1989). The chemical and biological significance of saponins in foods and feeding stuffs. *CRC Critical Review in Food Science and Nutrition*. 26: 127 - 135.
- Samson, A. A. (1980). *The State of Medicinal Plant Research in Nigeria*. Ibadan University Press. pp 222 - 223
- Trease G. E. and Evans, W. C. (1983). *Textbook of Pharmacognosy*. (12<sup>th</sup> ed) Balliere Tindall and Company Publishers, London.
- Vermon, U. V. (1983) *Tropical Seed Nutrition. American Biochemist*, 45: 17 - 19.