

LABORATORY ELEMENTS FOR ACCURATE DETECTION OF IRON DEFICIENCY IN RESOURCE-LIMITED SETTINGS OF AFRICA.

ATEGBO S.¹; MINTO'O S.¹; KOKO J.¹; DOUKAGA A.²; MOUSSAVOU A.¹; NGOU MILAMA E.³

(Manuscript N° E133. Received 08/12/2008. Accepted in revised form 26/03/2009) *Clin Mother Child Health* 2009; Vol 6, N° 1:1045-1046

¹Department of Paediatric, Faculty of Medicine and Health Sciences, BP. 4009, Libreville.

²Department of Haematology, Libreville Hospital Center, BP. 2228 Libreville.

³Biochemistry Department, Faculty of Medicine and Health Sciences, BP. 4009 Libreville

Correspondances: Dr. ATEGBO S., B.P. 9257 Libreville – Gabon. E mail: sategbo@yahoo.fr

Fax: (241) 73.71.57

I- INTRODUCTION

Iron deficiency is a frequent nutritional problem that predominantly affects children and pregnant women. The World Health Organisation (WHO) criteria for the diagnosis of iron deficiency is a serum ferritin < 12 µg/L, and/or a transferrin saturation < 10% [1]. Unfortunately, these assays remain quite inaccessible in many African countries where iron deficiency remains a public health problem. The objective of this study was to find a simple method of evaluating iron deficiency with a strong predictive value.

II- METHODOLOGY

Our study was approved by the Gabonese Ministry of Public Health. The prevalence of iron deficiency in children aged 6 months to 5 years was determined in Libreville using a random sample of 275 children, in a population estimated at 80,000, who appeared to be healthy in kinder garden and public health center. A blood sample of 3 mL was taken from the children in the morning prior to eating, for a complete blood count (CBC). Using micro method assay, by Coulter STKS of Beckman Coulter Gen. S. System™, the following was measured: haemoglobin (Hb; g/dL), mean corpuscular volume (MCV; fl), mean corpuscular haemoglobin (MCH; pg), mean corpuscular haemoglobin concentration (MCHC; %), and anisocytosis indicator (Rdw; %). The serum ferritin (µg/L) per immuno-enzymatic assay was taken with IM-X™ (Abbott), the serum iron (Fe; µg/dL) with the Alcyon 300/300i™ (Abbott), and the total iron binding capacity (TIBC; µg/dL), then the calculation of the percent transferrin saturation = (Fe/TIBC) × 100. On the basis of serum ferritin < 12µg/L, and by using Bayes theorem, we were able to evaluate positive predictive value (PPV) as well as the specificity and sensitivity of erythrocyte parameters analysed in our series.

III- RESULTS

Table I gives the values of sensitivity, specificity and PPV for the various erythrocyte parameters. The association by pairs of erythrocyte parameters, as evaluated using Pearson's coefficient, showed a strong correlation between microcytosis and anemia ($r = 0,90$; $p < 0,001$) as well as between a decrease of MCH and a rate of $Rdw \geq 15\%$ ($r = 0,67$; $p < 0,001$). No other significant correlations were seen between anemia and $Rdw \geq 15\%$, or between anemia and $MCH < 25$ pg.

Table I- Sensitivity, specificity, and positive predictive value (PPV) of erythrocyte parameters.

	Sensitivity (%)	Specificity (%)	PPV (%)
MCH < 25 pg	87.0	62.0	79.0
MCHC < 30%	30.0	90.0	81.0
RDW $\geq 15\%$	90.0	34.0	66.0
MCV < 70 fl	39.0	88.0	80.0
MCH < 25 pg and RDW $\geq 15\%$	60.0	88.0	93.0
Hb < 11 g/dL and MCV < 70 fl	20.0	90.0	96.0

Hb: Haemoglobin; MCH: Mean Corpuscular Haemoglobin; MCHC: Mean Corpuscular Haemoglobin Concentration; MCV: Mean Corpuscular Volume; RDW: Red cell Distribution Width;

IV- DISCUSSION

The MCH and the RDW could be used in combination to evaluate iron deficiency of an individual, in the presence of a normal haemoglobin rate. The weak sensitivity of the CMV in our study raises the issue of its effectiveness as the only source of early detection of iron deficiency in the tropics. These results differ from that of OSKI et al, who were quite confident in this test for the early detection of iron deficiency [2]. The 90% and 87% sensitivity of the $RDW \geq 15\%$ and $MCH < 25$ pg, respectively, are similar to the results of Das Gupta

et al. in India [3] who found a 96% sensitivity for the RDW $\geq 15\%$ and a 69% sensitivity for the MCH $< 25\text{pg}$. VISWANATH et al, have reported 82% to 100% sensitivity for the RDW $\geq 15\%$ [4]. These two parameters that are present on a simple CBC, would allow for the early discovery of iron deficiency. Therefore KNIGHT et al, affirm that the correlation between RDW $\geq 15\%$ and MCH $< 25\text{pg}$ found again in their series, would allow for their use to detect iron deficiency up to 90% [5]. Kim et al. have found a positive prediction value of 98% for the association of CMV $< 70\text{ fl}$ and RDW $\geq 15\%$ with a specificity of CMV $< 70\text{ fl}$ up to 97.8% for children [6].

V- CONCLUSION

Although the ferritin test remains the method of reference for evaluating the iron deficiency status of an individual, it is not practical to generalize its use in poor African countries, because of the high cost and technical exigencies of its execution. In clinical practice, the evaluation of iron deficiency without interaction from an inflammatory pathology, prevents the occurrence of the pernicious effects of iron deficiency, and guarantees an overall and optimal coverage of this any deficiencies. Our study has permitted us to address these two concerns by reading a simple blood count ■

REFERENCES:

1. World Health Organization (WHO). Iron deficiency anemia, assessment, prevention and control. Department of Nutrition for Health and Development. World Health Organization, 1211 Geneva, Switzerland. 2001; 12-45.
2. Oski FA, Honig AS, Helu BM, Hoanitz. Iron deficiency in infancy and childhood. N Eng J Med 1993; 329:190-93.
3. Das Gupta A, Hedge C, Mistri R. Red cell distribution width as a measure of severity of iron deficiency anemia. Ind J Med Res 1994; 100:177-83
4. Viswanath D, Hegde R, Murthy V, Nagashree S, Shah R. Red cell distribution width in the diagnosis of iron deficiency anemia. Indian J Pediatr. 2001; 68:1117-9.
5. Knight GJ, Heese Hde V, Dempster WS, Kirsten G. Diagnosis of iron deficiency: mean corpuscular haemoglobin (CMH) as a predictor of iron deficiency in infants. Pediatr Res 1983; 17:168-70.
6. Kim SK, Cheong WS, Jun HS, Choi JW, Son BX. Red blood cell indices and iron status according to feeding practices in infants and young children. Acta Pediatr 1996; 85:139- 44.