

EFFECTS OF TRANEXAMIC ACID ON DEATH, VASCULAR OCCLUSIVE EVENTS, AND BLOOD TRANSFUSION IN TRAUMA PATIENTS WITH SIGNIFICANT HAEMORRHAGE (CRASH-2): A RANDOMISED, PLACEBO-CONTROLLED TRIAL

V.M. Mutiso*, MBChB(UON), MMed(Surg) (UON), Fellow (arthroscopy and arthroplasty) (UK), Fellow AO-International (Ger), FCS (ecsa), Consultant Orthopaedic and Trauma Surgeon and Lecturer Department of Surgery, College of Health Sciences, University of Nairobi, **D. Ojuka**, MMed(Surg), Consultant General Surgeon and Lecturer Department of Surgery, Collage of Health Sciences, University of Nairobi and **B. Okanga**, MMed(Surg), Surgeon presently pursuing neurosurgery at the University of Nairobi Medical School at Kenyatta National Hospital.

*National Coordinator for Kenya on behalf of CRASH2 trial collaborators. This manuscript has been reprinted with authority from The Lancet.

Correspondence to: Dr. V.M. Mutiso, P.O. Box 19681-00200, Nairobi, Kenya. Email: mutiso@uonbi.co.ke

ABSTRACT

Background: Tranexamic acid can reduce bleeding in patients undergoing elective surgery. We assessed the effects of early administration of a short course of tranexamic acid on death, vascular occlusive events, and the receipt of blood transfusion in trauma patients.

Methods: This randomised controlled trial was undertaken in 274 hospitals in 40 countries. 20 211 adult trauma patients with, or at risk of, significant bleeding were randomly assigned within 8 h of injury to either tranexamic acid (loading dose 1 g over 10 min then infusion of 1 g over 8 h) or matching placebo. Randomisation was balanced by centre, with an allocation sequence based on a block size of eight, generated with a computer random number generator. Both participants and study staff (site investigators and trial coordinating centre staff) were masked to treatment allocation. The primary outcome was death in hospital within 4 weeks of injury, and was described with the following categories: bleeding, vascular occlusion (myocardial infarction, stroke and pulmonary embolism), multiorgan failure, head injury, and other. All analyses were by intention to treat. This study is registered as ISRCTN86750102, Clinicaltrials.gov NCT00375258, and South African Clinical Trial Register DOH-27-0607-1919.

Findings 10 096 patients were allocated to tranexamic acid and 10 115 to placebo, of whom 10 060 and 10 067, respectively, were analysed. All-cause mortality was significantly reduced with tranexamic acid (1463 [14.5%] tranexamic acid group vs 1613 [16.0%] placebo group; relative risk 0.91, 95% CI 0.85-0.97; $p=0.0035$). The risk of death due to bleeding was significantly reduced (489 [4.9%] vs 574 [5.7%]; relative risk 0.85, 95% CI 0.76-0.96; $p=0.0077$).

Interpretation Tranexamic acid safely reduced the risk of death in bleeding trauma patients in this study. On the basis of these results, tranexamic acid should be considered for use in bleeding trauma patients.

Funding UK NIHR Health Technology Assessment programme, Pfizer, BUPA Foundation, and JP Moulton Charitable Foundation.

INTRODUCTION

Injuries are major causes of death worldwide (1,2). Every year, more than a million people die as a result of road traffic injuries around the world. Road traffic injuries are the ninth leading cause of death globally, and such

injuries are predicted to become the third leading cause of death and disability by 2020. About 1.6 million people die as a result of intentional acts of interpersonal, collective, or self-directed violence every year. More than 90% of trauma deaths occur in low-income and middle-income countries (2). Haemorrhage is responsible for

about a third of in-hospital trauma deaths and can also contribute to deaths from multiorgan failure (3).

The haemostatic system helps to maintain circulation after severe vascular injury, whether traumatic or surgical in origin (4). Major surgery and trauma trigger similar haemostatic responses, and in both situations severe blood loss presents an extreme challenge to the coagulation system. Part of the response to surgery and trauma is stimulation of clot breakdown (fibrinolysis), which might become pathological (hyperfibrinolysis) in some cases (4). Antifibrinolytic agents reduce blood loss in patients with both normal and exaggerated fibrinolytic responses to surgery, and do so without apparently increasing the risk of postoperative complications (5).

Tranexamic acid is a synthetic derivative of the amino acid lysine that inhibits fibrinolysis by blocking the lysine binding sites on plasminogen (6). A systematic review of the randomised trials of tranexamic acid in patients undergoing elective surgery identified 53 studies including 3836 participants (5). Tranexamic acid reduced the need for blood transfusion by a third (relative risk [RR] 0.61, 95% CI 0.54–0.70), with no significant reduction in mortality (0.61, 0.32–1.12) (5). Because the haemostatic responses to surgery and trauma are similar, (4) tranexamic acid might reduce mortality due to bleeding in trauma patients. However, up until now there have been no randomised trials of this drug in such patients (7). We assessed the effects of the early administration of a short course of tranexamic acid on death, vascular occlusive events, and the receipt of blood transfusion in trauma patients with or at risk of significant haemorrhage.

METHODS

Study design and patients

CRASH-2 (Clinical Randomisation of an Antifibrinolytic in Significant Haemorrhage 2) (www.crash2.lshtm.ac.uk) is a large placebo controlled trial of the effects of early administration of a short course of tranexamic acid on death, vascular occlusive events, and the receipt of blood transfusion. The trial was undertaken in 274 hospitals in 40 countries. The first patient was enrolled in May, 2005. The study aims, methods, and protocol have been reported previously. The trial protocol was peer-reviewed and published on *The Lancet* website in 2005 (www.thelancet.com/protocol-reviews/05PRT-1).

Adult trauma patients with significant haemorrhage (systolic blood pressure <90 mm Hg or

heart rate >110 beats per min, or both), or who were considered to be at risk of significant haemorrhage, and who were within 8 h of injury, were eligible for the trial. Patients were included if the responsible doctor was substantially uncertain about whether or not to treat with tranexamic acid (ie, entry was governed by the uncertainty principle) (8). Patients for whom the responsible doctor considered that there was a clear indication for tranexamic acid were not randomly assigned. Similarly, patients for whom there was considered to be a clear contraindication to tranexamic acid treatment were not randomly assigned. However, when the responsible doctor was substantially uncertain as to whether or not to treat with this agent, these patients were eligible for randomisation.

Consent procedures at participating hospitals were established by local regulation and the appropriate ethics committees. Informed consent was obtained from patients if physical and mental capacity allowed. If patients could not give consent, proxy consent was obtained from a relative or representative. If a proxy was unavailable, then if permitted by local regulation, consent was deferred or waived. When consent was deferred or given by a proxy, the patient was informed about the trial as soon as possible and consent obtained for use of the data collected if needed.

Randomisation and masking

After eligibility had been confirmed and the locally approved consent procedures had been completed, patients were randomly assigned. Randomisation was balanced by centre, with an allocation sequence based on a block size of eight, generated with a computer random number generator. In hospitals in which telephone randomisation was not practicable we used a local pack system that selected the lowest numbered treatment pack from a box containing eight numbered packs. Apart from the pack number, the treatment packs were identical. The pack number was recorded on the entry form which was sent to the international trial coordinating centre in London, UK. Hospitals with reliable telephone access used the University of Oxford Clinical Trial Service Unit (CTSU) telephone randomisation service. The randomisation service used a minimisation algorithm balancing for sex, age, time since injury, type of injury (blunt or penetrating), Glasgow Coma Score, systolic blood pressure, respiratory rate, central capillary refill time, and country, taking into account what packs were available at that hospital. Once the treatment pack number was

recorded, the patient was included in the trial whether or not the treatment pack was opened or the allocated treatment started. Both participants and study staff (site investigators and trial coordinating centre staff) were masked to treatment allocation.

Tranexamic acid and placebo ampoules were indistinguishable. Tranexamic acid was manufactured by Pharmacia (Pfizer, Sandwich, UK) and placebo by St Mary's Pharmaceutical Unit, Cardiff, UK. The treatment packs were prepared by an independent clinical trial supply company (Bilcare, Crickhowell, UK). Correct blinding and coding of ampoules was assured by independent random testing of each batch by high performance liquid chromatography to confirm the contents. Emergency unblinding was available by telephoning CTSU.

Procedures

Patients were randomly allocated to receive a loading dose of 1 g of tranexamic acid infused over 10 min, followed by an intravenous infusion of 1 g over 8 h, or matching placebo (0.9% saline). Every patient was assigned a uniquely numbered treatment pack, which contained four ampoules of either tranexamic acid 500 mg or placebo, one 100 mL bag of 0.9% saline (for use with the loading dose), a syringe and needle, stickers with the trial details and randomisation number (for attaching to infusion bags, data forms, and patient medical records), and instructions. Each box contained information leaflets for patients and their representatives, consent forms, and data collection forms. The stickers, instructions, leaflets, and forms were in local languages.

Outcome measures and prespecified subgroup analyses

The primary outcome was death in hospital within 4 weeks of injury. Cause of death was described by the following categories: bleeding, vascular occlusion (myocardial infarction, stroke, and pulmonary embolism), multiorgan failure, head injury, and other. Secondary outcomes were vascular occlusive events (myocardial infarction, stroke, pulmonary embolism, and deep vein thrombosis), surgical intervention (neurosurgery, thoracic, abdominal, and pelvic surgery), receipt of blood transfusion, and units of blood products transfused. Dependency was measured at hospital discharge, or on day 28 if still in hospital, with the 5-point Modified Oxford Handicap Scale. The scale was dichotomised into dead or dependent (dead, fully dependent requiring attention day and night, or dependent but not needing

constant attention) or independent (some restriction in lifestyle but independent, minor symptoms, or no symptoms) (9). Data for the use of recombinant Factor VIIa and for gastrointestinal bleeding as a complication were also collected. Because the expected complications of the trial treatment were collected on the outcome form, only adverse events that were serious, unexpected, and suspected to be related to the study treatment were reported separately. Outcomes were recorded if they occurred while the patient was still in hospital for up to 28 days after randomisation. Data were sent to the coordinating centre either electronically (by encrypted electronic data forms which could be sent by email or uploaded to a secure server) or by fax, and were entered onto a central database at the trial coordinating centre in London. We monitored the quality of the trial data using a combination of centralised statistical data checking and site visits at which patient outcome forms were compared with clinical case notes (10).

We planned to report the effects of treatment on the primary outcome subdivided by four baseline characteristics:

- (1) estimated hours since injury (<1, 1–3, 3–8 h);
- (2) systolic blood pressure (≤ 75 , 76–89, ≥ 90 mm Hg);
- (3) Glasgow Coma Score (severe 3–8, moderate 9–12, mild 13–15); and
- (4) type of injury (penetrating only or blunt, which included blunt and penetrating).

Statistical analyses

The statistical analysis plan was sent to all ethics committees and regulatory agencies before unblinding. Because the risk of death might be around 20%, and even a 2% survival difference (corresponding to an RR of death with tranexamic acid of 0.9) would be important, a trial of 20 000 patients was planned, which would then have an 85% chance of achieving a two-sided p value of less than 0.01 and a 95% chance of a two-sided p value of less than 0.05. All analyses were undertaken on an intention-to-treat basis. For each binary outcome, we calculated RRs and 95% CIs, and two-sided p values for statistical significance. The RR gives the number of times more likely ($RR > 1$) or less likely ($RR < 1$) an event is to happen in the tranexamic acid group compared with the placebo group. For analysis of the prespecified subgroups (primary outcome only) we calculated RRs with 99% CIs with two-sided p values. Heterogeneity in treatment effects across subgroups was assessed with 2 tests. We prespecified that unless there was strong evidence ($p < 0.001$) against homogeneity of effects, the overall RR would be considered the most reliable

guide to the approximate RRs in all subgroups. Means and SDs were estimated for count outcomes, and we calculated two-sided p values of the difference in means of logarithms. A complete case analysis, including only cases for which the relevant outcome data were available, was undertaken. There was no imputation for missing data. During the study, unblinded interim analyses were supplied by an independent statistician to the Data Monitoring and Ethics Committee.

This study is registered as ISRCTN86750102, Clinicaltrials.gov NCT00375258, and South African Clinical Trial Register DOH-27-0607-1919.

Role of the funding source

Funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The Writing Committee had full access to all data in the study and had final responsibility for the decision to submit for publication.

RESULTS

Figure 1 shows the trial profile. 20 211 patients were randomly assigned to tranexamic acid or placebo (figure 1), of whom 20 116 were randomly assigned through the local pack system and 95 through telephone randomisation. The data from four patients were removed from the trial because their consent was withdrawn after randomisation. Five patients enrolled in the study were later found to be younger than 16 years. Age was unknown for four patients. 23 patients were enrolled more than 8 h after their injury. Time of injury was not known for 11 patients. Nine patients had haemorrhage from non-traumatic conditions. Three patients were given a pack that differed from that allocated. The planned consent procedures were not fully followed in 34 patients. The relevant ethics committees were informed and approval for use of data was obtained. All the patients, apart from the four in whom consent was withdrawn, were included in the analysis.

Figure 1: Trial profile




Figure 1
Mortality by day from Randomisation




Table 1
Baseline data or participants

	Tranexamic acid [n=10,093]	Placebo [n=10,114]
Sex		
Men	8,439 (83.6%)	8,496 (84%)
Women	1,654 (16.4%)	1,617 (16%)
Not known	0	1 (0.01%)
Age (years)		
Mean (SD)	34.6(14.1)	34.5 (14.4)
<25 [#]	2,783 (27.6%)	2,855 (28.2%)
25-34	3,012 (29.8%)	3,081 (30.5%)
35-44	1,975 (19.6%)	1,841 (18.2%)
>44	2,321 (23.0%)	2,335 (23.1%)
Not known	2 (0.02%)	2 (0.02%)
Time since injury (h)		
Mean (SD)	2.8 (2.2)	2.9 (2.6)
<1	3,756 (37.2%)	3,722 (36.8%)
> 1 to ≤ 3	3,045 (30.2%)	3,006 (29.7%)
>3*	3,287 (32.6%)	3,380 (33.4%)
Not known	5 (0.05%)	6 (0.06%)
Type of injury		
Blunt [†]	6,812 (67.5%)	6,843 (67.7%)
Penetrating	3,281 (32.5%)	3,271 (32.3%)
Systolic blood pressure (mm Hg)		
≤ 75	1,566 (15.5%)	1,608 (15.9%)
76-89	1,615 (16.0%)	1,697 (16.8%)
>89	6,901 (68.4%)	6,791 (67.1%)
Not known	11 (0.11%)	18 (0.18%)

Respiratory rate (per min)		
<10	160 (1.6%)	149 (1.5%)
10-29	8,355 (82.8%)	8,436 (83.1%)
>29	1,491 (14.8%)	1,429 (14.1%)
Not known	87 (0.86%)	100 (0.99%)
Central capillary refill time (s)		
2 or less	3,432 (34.0%)	3,406 (33.7%)
3 to 4	4,665 (46.2%)	4,722 (46.7%)
>4	1,699 (16.8%)	1,672 (16.5%)
Not known	297 (2.9%)	314 (3.1%)
Heart rate (beats per min)		
<77	875 (8.7%)	871 (8.6%)
77-91	1,727 (17.1%)	1,770 (17.5%)
92-107	2,556 (25.3%)	2,546 (25.2%)
>107	4,872 (48.3%)	4,853 (48.0%)
Not known	63 (0.62%)	74 (0.73%)
Glasgow Coma Score (total)		
Severe [3-8]	1,799 (17.8%)	1,839 (18.2%)
Moderate [9-12]	1,353 (13.4%)	1,351 (13.3%)
Mild [13-15]	6,934 (68.7%)	6,908 (68.3%)
Not known	7 (0.07%)	16 (0.16%)
Any protocol violation	39 (0.4%)	39 (0.4%)

Data are number (% of group total), unless otherwise indicated. #Includes five patients aged younger than 16 years. *Includes 23 patients randomly assigned more than 8 h after injury. †includes patients with both blunt and penetrating and those with only blunt injuries.

Treatment groups were balanced with respect to all baseline patient characteristics (table 1; the web appendix p 1 shows baseline data of patients with follow-up). Primary outcome data were available for 20 127 (99.6%) randomised patients, 10 060 allocated to tranexamic acid and 10 067 placebo, of whom 19 944 (99.1%) patients were known to have completed the loading dose and 18 965 (94.2%) the 8 h maintenance dose. 3076 (15.3%) patients died, of whom 1086 (35.3%) died on the day of randomisation (figure 2). There were 1063 deaths due to bleeding, of which 637 (59.9%) were on the day of randomisation.

All-cause mortality was significantly reduced with tranexamic acid (table 2). The RR of death with tranexamic acid was 0.91 (95% CI 0.85–0.97, $p=0.0035$; table 2). The risk of death due to bleeding was significantly reduced (table 2). This effect was also apparent for deaths due to bleeding on the day of randomisation (282 [2.8%] tranexamic acid group vs 355 [3.5%] placebo group; RR 0.80, 95% CI 0.68–0.93, $p=0.0036$). There were 33 (0.3%) deaths in the tranexamic acid group versus 48 (0.5%) in the placebo group from vascular occlusion (table 2), including seven versus 22 deaths from myocardial

infarction, eight versus five from stroke, and 18 versus 21 from pulmonary embolism, respectively. Deaths from multiorgan failure, from head injury, or due to other causes did not differ significantly in the tranexamic acid group versus the placebo group (table 2).

Vascular occlusive events (fatal or non-fatal) did not differ significantly, with 168 (1.7%) patients with one or more vascular occlusive events (myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis) in patients allocated to tranexamic acid versus 201 (2.0%) in those allocated to placebo (table 3).

Blood product transfusions were given to 5067 (50.4%) patients allocated to tranexamic acid versus 5160 (51.3%) allocated to placebo (table 3). Those allocated to tranexamic acid and transfused received a mean of 6.06 (SD 9.98) blood units, compared with a mean of 6.29 (10.31) for placebo. 4814 (47.9%) patients in the tranexamic acid group received one or more surgical intervention (neurosurgery, or chest, abdominal, or pelvic surgery) versus 4836 (48.0%) in the placebo group (table 3). Only 17 patients received treatment

Table 2
Death by cause

Cause of death	Tranexamic acid (n = 10.060)	Placebo [n = 10.067]	RR (95% CI)	P value (two-sided)
Any cause of death	463 (14-5%)	1.613 (16-0%)	0.91 (0.85 - 0.97)	0.0035
Bleeding	489 (4-9%)	574 (5-7%)	0.85 (0.76 - 0.96)	0.0077
Vascular occlusion*	33 (0-3%)	48 (0-5%)	0.69 (0.44 - 1.07)	0.096
Multiorgan failure	209 (2.1 %)	233 (23%)	0.90 (0.75 - 1.08)	0.25
Head injury	603 (6,0%)	621 (62%)	0.97 (0.87 - 1.08)	0.60
Other causes	129 (1,3%)	137 (14%)	0.94 (0.74 - 1.20)	0.63

Data are number (%). unless otherwise indicated. RR=relative risk.

*Includes myocardial infarction. Stroke and pulmonary embolism.

Table 3
Vascular occlusive events need for transfusion and surgery, and level of dependency

	Tranexamic acid (n=10060)	Placebo (n=10067)	RR (95% CI)	P value
Vascular occlusive events*				
Any vascular occlusive event	168 (1.7%)	201 (2.0%)	0.84 (0.68 - 1.02)	0.084
Myocardial infarction	35 (0.4%)	55 (0.6%)	0.64 (0.42 - 0.97)	0.035
Stroke	57 (0.6%)	66 (0.7%)	0.86 (0.61 - 1.23)	0.42
Pulmonary embolism	72 (0.7%)	71 (0.7%)	1.00 (0.73 - 1.40)	0.93
Deep vein thrombosis	40 (0.4%)	41 (0.4%)	0.98 (0.63 - 1.51)	0.91
Need for transfusion and surgery				
Blood product transfused	5.067 (50.4%)	5.160 (51.3%)	0.98 (0.96 - 1.01)	0.21
Any surgery	4.814 (47.9%)	4.836 (48.0%)	1.00 (0.97 - 1.02)	0.79
Neurosurgery	1.040 (10.3%)	1.059 (10.5%)	0.98 (0.91 - 1.07)	0.67
Chest surgery	1.518 (15.1%)	1.525 (15.2%)	1.00 (0.93 - 1.06)	0.91
Abdominal surgery	2,487 (24.7%)	2,555 (25.4%)	0.97 (0.92 - 1.00)	0.28
Pelvic surgery	683 (6.8%)	648 (6.4%)	1.05 (0.95 - 1.17)	0.31
Median (IQR) units of blood product transfused*	3 (2-6)	3 (2-6)	..	0.591
Dependency				
No symptoms	1,483 (17.3%)	1,334 (15.8%)	1.09 (1.02 - 1.17)	0.0086
Minor symptoms	3,054 (35.5%)	3,061 (36.2%)	0.98 (0.94 - 1.02)	0.39
Some restriction	2,016 (23.5%)	2,069 (24.5%)	0.96 (0.91 - 1.01)	0.13
Dependent (not requiring constant attention)	1,294 (15.1%)	1,273 (15.1%)	1.00 (0.93 - 1.07)	0.98
Fully dependent	696 (8.1%)	676 (8.0%)	1.01 (0.92 - 1.12)	0.79
Alive (disability status not known)	54 (0.6%)	41 (0.5%)
Dead	1,463 (14.5%)	1,613 (16.0%)	0.91 (0.85 - 0.97)	0.0035

Data are number (%). unless otherwise indicated. Counts are for numbers of patients with at least one such event. RR=relative risk includes both fatal and non-fatal events. *Transfused patients only. #Analysis used logarithmic transformation of mean units of blood products transfused

with recombinant Factor VIIa (13 in the tranexamic acid group vs four in the placebo group). 132 patients in each group had gastrointestinal bleeding ($p=0.99$).

Of patients allocated tranexamic acid, 3453 (34.3%) were classified as dead or dependent at discharge or 28 days compared with 3562 (35.4%) of those allocated to placebo (RR 0.97, 95% CI 0.93–1.00; $p=0.12$). 1483 (14.7%) patients in the tranexamic acid group had no symptoms at discharge or day 28 versus 1334 (13.3%) in the placebo group (table 3). 1846 (9.2%) patients were still in hospital at 28 days (958 vs 888).

We had prespecified that unless there was strong evidence ($p<0.001$) against homogeneity of effects, the overall RR would be regarded as the most reliable guide as to the approximate RRs in all subgroups. We recorded no such evidence of heterogeneity for any of the prespecified subgroup analyses: systolic blood pressure (heterogeneity $p=0.51$); Glasgow Coma Score at randomisation ($p=0.50$); type of injury ($p=0.37$); or time from injury to randomisation ($p=0.11$). For the

last of these analyses, because of digit preference (the tendency when reporting figures to round to specific digits) the number of patients in the early category (<1 h) was low and the subgroup estimate was imprecise. We therefore (post hoc) defined the early category as those treated less than or equal to 1 h from injury (figure 3).

No emergency unblinding was needed, and there were no adverse events regarded as serious, unexpected, or suspected to be related to the study treatment.

DISCUSSION

The results show that the early administration of tranexamic acid to trauma patients with, or at risk of, significant bleeding reduces the risk of death from haemorrhage with no apparent increase in fatal or nonfatal vascular occlusive events. All-cause mortality was significantly reduced with tranexamic acid.




Figure 3: All-cause mortality by subgroups
GCS=Glasgow Coma Score.*95% CI.

The trial inclusion criteria were clinical and did not depend on the results of laboratory tests. Patients were enrolled if they were judged to have on-going significant haemorrhage, as evidenced by hypotension or tachycardia, or if they were considered to be at risk of significant haemorrhage—eg, patients with compensated haemorrhage and stable vital signs, or those in whom bleeding might have stopped but who might recommence bleeding following volume resuscitation. The use of clinical inclusion criteria is appropriate in the context of traumatic bleeding in which a range of clinical signs need to be assessed when establishing the presence or absence of haemorrhage, while taking into account remedial measures such as fluid resuscitation. The clinical inclusion criteria, and the large numbers of patients studied in a range of different health-care settings, help these results to be generalised widely.

Our study had strengths and limitations. The randomisation methods ensured that participating clinicians did not have foreknowledge of treatment allocation. Baseline prognostic factors were well balanced. All analyses were on an intention-to-treat basis and, because almost all randomised patients were followed up, there was no need to use imputation methods for missing data (11). The primary endpoint was all-cause mortality, and the observed reduction in mortality with tranexamic acid was both statistically significant and clinically important. The diagnosis of traumatic haemorrhage can be difficult, and some of the included patients might not have been bleeding at the time of randomisation. This misdiagnosis would have reduced the power of the trial to show an effect of tranexamic acid on mortality from bleeding. Nevertheless, we recorded a significant reduction in death due to bleeding.

Although we recorded no increased risk of non-fatal vascular occlusive events with tranexamic acid, the precision of the estimates was low and we cannot exclude the possibility of some increase in risk. In the context of outcome assessment in clinical trials, estimates of the RR are unbiased even when the sensitivity of diagnosis is imperfect, provided that there are few false positives (high specificity) (12). Therefore, we sought high specificity in the diagnosis of non-fatal vascular occlusive events and stipulated that occlusive events should be recorded only when there was clear clinical evidence. As a result, we might have under-reported the frequency of these events. However, our estimates of the RR of non-fatal occlusive events should be unbiased (12).

One weakness of this trial is that it provides limited insight into how tranexamic acid reduces the risk of death in bleeding trauma patients. Early coagulation abnormalities are frequent in severely injured trauma patients and are associated with substantially increased mortality (13). Recent research showing that hyperfibrinolysis is a common feature of these abnormalities raises the possibility that antifibrinolytic agents such as tranexamic acid might operate via this mechanism (13). Furthermore, intravenous tranexamic acid administration has an early (within 4 h) antifibrinolytic effect (14). However, although this mechanism is plausible, because we did not measure fibrinolytic activity in this trial we cannot conclude that this agent acts by reducing fibrinolysis, rather than another mechanism. Further studies are needed into the mechanism of action of tranexamic acid in bleeding trauma patients. Measurement of blood loss is difficult in trauma patients. Much of the bleeding occurs at the scene of the injury and the bleeding that occurs in hospital is often concealed and difficult to quantify, such as, for example, bleeding into the chest, abdomen, pelvis, and soft tissues. However, we did not find any substantial reduction in the receipt of a blood transfusion or the amount of blood transfused in trauma patients. This finding could be an indication of the difficulty of accurate estimation of blood loss in trauma patients when assessing the need for transfusion. Another possible explanation is that after the loading dose, tranexamic acid was infused over 8 h, whereas decisions about transfusion are made soon after admission. Finally, fewer deaths occurred in patients allocated to tranexamic acid than to placebo, and the patients who survived as a result of tranexamic acid administration would have had a greater opportunity to receive a blood transfusion (competing risks).

The tranexamic acid loading dose was given within 8 h of injury, followed by a maintenance infusion over 8 h. We chose the early administration of a short course of tranexamic acid because most deaths from bleeding occur on the day of the injury and we postulated that the drug would act by reducing bleeding. Generally, after the first day, the risk of death from haemorrhage is reduced but the risk of vascular occlusive events might remain. We therefore selected a regimen that would allow for the effect of tranexamic acid on the early risk of haemorrhage without extending into the period when the risk of vascular occlusive events might be increased by this treatment. The absence of any increase in vascular occlusion with tranexamic acid, whether fatal or non-fatal, provides

reassurance that this regimen is safe. Although the effect of this drug on all-cause mortality did not vary substantially according to the time from injury, there was some suggestion that early treatment might be more effective. However, even if this were not the case, the fact that most deaths from haemorrhage occur in the first few hours after injury implies that every effort should be made to treat patients as soon as possible (15–17).

The dose of tranexamic acid used in this trial was based on studies of this drug in surgical patients in which loading doses range from 2.5 mg/kg to 100 mg/kg, and maintenance doses from 0.25 mg/kg/h to 4 mg/kg/h, delivered over 1–12 h (5). Findings from studies of the effect of different doses of tranexamic acid on blood loss and blood transfusion showed no significant difference between high and low doses. Studies in cardiac surgery have noted that a 10 mg/kg loading dose of tranexamic acid followed by an infusion of 1 mg/kg/h produces plasma concentrations sufficient to inhibit fibrinolysis, and that a larger dose does not provide any additional haemostatic benefit (18,19). In emergency situations, the administration of a fixed dose is practicable since determining the weight of a seriously injured patient can be difficult. We therefore selected a fixed dose within the range shown to inhibit fibrinolysis and provide haemostatic benefit that would be efficacious for larger patients (>100 kg) but also safe in smaller patients (<50 kg), to the extent that the dose per kg that smaller patients would receive has been used in surgical trials without adverse effects. The possibility that a higher dose of tranexamic acid would have a greater treatment effect remains open to debate and warrants further study.

The knowledge that tranexamic acid reduces the risk of death from traumatic bleeding raises the possibility that it might also be effective in other situations in which bleeding can be life threatening or disabling. Traumatic brain injury is commonly accompanied by intracranial bleeding, which can develop or worsen after hospital admission. Traumatic intracranial haemorrhage is associated with an increased risk of death and disability, and irrespective of location, haemorrhage size is strongly correlated with outcome (20–22). If tranexamic acid reduced intracranial bleeding after isolated traumatic brain injury, then patient outcomes might be improved. Studies that assess the effect of tranexamic acid on the extent of intracranial bleeding are needed.

Tranexamic acid might also have a role in bleeding conditions apart from traumatic injury. Post-

partum haemorrhage is a leading cause of maternal mortality, accounting for about 100 000 maternal deaths every year (23). Although evidence suggests that this drug reduces postpartum bleeding, the quality of the existing trials is poor and none has been large enough to assess the effect of tranexamic acid on endpoints that are important to women (24). A large trial is being undertaken to assess the effect of tranexamic acid on the risk of death and hysterectomy in women with postpartum haemorrhage (25).

In conclusion, tranexamic acid could be given in a wide range of health-care settings, and safely reduced the risk of death in bleeding trauma patients in our study. The option to use tranexamic acid should be available to doctors treating trauma patients in all countries, and this drug should be considered for inclusion on the WHO List of Essential Medicines. On the basis of these results, tranexamic acid should be considered for use in bleeding trauma patients.

CRASH-2 trial coordination

Writing Committee—Haleema Shakur (chair), Ian Roberts (chief investigator), Raúl Bautista (Mexico), José Caballero (Peru), Tim Coats (UK), Yashbir Dewan (India), Hesham El-Sayed (Egypt), Tamar Gogichaishvili (Georgia), Sanjay Gupta (India), Jorge Herrera (Colombia), Beverley Hunt (UK), Pius Iribhogbe (Nigeria), Mario Izurieta (Ecuador), Hussein Khamis (Egypt), Edward Komolafe (Nigeria), María-Acelia Marrero (Cuba), Jorge Mejía-Mantilla (Colombia), Jaime Miranda (Peru), Carlos Morales (Colombia), Oluwole Olaomi (Nigeria), Fatos Olldash (Albania), Pablo Perel (UK), Richard Peto (UK), PV Ramana (India), R R Ravi (India), Surakrant Yutthakasemsunt (Thailand).

National coordinators—Jonathan Dakubo (Ghana), Tamar Gogichaishvili (Georgia), Nyoman Golden (Indonesia), Mario Izurieta (Ecuador), Hussein Khamis (Egypt), Edward Komolafe (Nigeria), Jorge Loría-Castellanos (Mexico), Jorge Mejía-Mantilla (Colombia), Jaime Miranda (Peru), Ángeles Muñoz (Spain), Vincent Mutiso (Kenya), Patrick Okwen (Cameroon), Zulma Ortiz (Argentina), María Pascual, CENCEC (Cuba), R Ravi (India), April Roslani (Malaysia), Stefan Trenkler (Slovakia), Annalisa Volpi (Italy), Surakrant Yutthakasemsunt (Thailand).

Trial Coordinating Centre Team— Ian Roberts (clinical coordinator, chief investigator), Haleema Shakur (trial manager), Pablo Perel (regional coordinator), Lin Barnetson (data manager), Maria Ramos (trial administrator), Lisa Cook (assistant trial manager, regional coordinator from 2007), Taemi Kawahara

(assistant trial manager, regional coordinator from 2007), Eni Balogun (regional coordinator from 2006), Matthew Berle (trial assistant from 2007), Collette Barrow (assistant administrator from 2008), Tony Brady (programmer to 2006), Chris Rubery (data assistant from 2009), Jackie Wayte (UK nurse co-ordinator from 2008), Cynthia To (data assistant 2007–09).

Steering Committee—Ian Franklin (chair), Brigitte Chaudhry, Tim Coats, Charles Deakin, Steve Goodacre, Beverley Hunt, David Meddings, Richard Peto, Ian Roberts, Peter Sandercock.

Management Group—Ian Roberts (chair), Haleema Shakur, Tim Coats, Phil Edwards, Beverley Hunt, Maria Ramos.

Data Monitoring and Ethics Committee—Rory Collins (chair), Adrian Grant, John Myburgh, Alex Baxter (independent statistician).

CRASH-2 trial collaborators by country

Albania (115)—National Trauma Centre Hospital: Fatos Olldash, Mihal Kerçi, Tefik Zhurda, Klotilda Ruçi; Spitali Civil Durres: Arben Banushi. *Argentina (51)*—Hospital Ángel Cruz Padilla: Mario Sardón Traverso, Juan Jiménez; Hospital Regional Rio Grande: Jorge Balbi; Hospital “4 de Junio” Dr Ramon Carrillo: Christian Dellerá; Hospital Castro Rendón: Silvana Svampa; Hospital San Martín de La Plata: Gustavo Quintana; Hospital Municipal de Agudos “Dr Leonidas Lucero”: Gustavo Piñero; Hospital Interzonal General de Agudos “Dr Oscar Alende”: Jorge Teves. *Australia (17)*—Nepean Hospital: Ian Seppelt; Sir Charles Gairdner Hospital: David Mountain; John Hunter Hospital: Zsolt Balogh. *Bangladesh (12)*—United Hospital Limited: Maniruz Zaman. *Belgium (51)*—Sint-Vincentius Hospital: Patrick Druwé, Robert Rutsaert; Centre Hospitalier Regional de Namur: Guy Mazairac. *Cameroon (124)*—Tombel District Hospital: Fogang Pascal, Zognou Yvette, Djeuchon Chancellin; St Theresa’s Catholic Medical Centre: Patrick Okwen; Bamenda Provincial Hospital: Jules Djokam-Liapoe; Bali District Hospital: Ernest Jangwa; Bafut District Hospital: Lawrence Mbuagbaw; Fundong District Hospital: Ninying Fointama; St John of God Medical Centre: Nguemo Pascal. *Canada (2)*—Hamilton General Hospital: Frank Baillie. *China (51)*—Renji Hospital: Ji-yao Jiang, Guo-yi Gao, Yin-hui Bao. *Colombia (2940)*—Hospital Universitario San Vicente de Paúl, Universidad de Antioquia: Carlos Morales, Juan Sierra, Santiago Naranjo, Camilo Correa, Carolina Gómez; Hospital Universitario San Jose de Popayan: Jorge Herrera, Liliana Caicedo, Alexei Rojas, Henry Pastas, Hugo Miranda; Hospital Pablo Tobon Uribe: Alfredo Constaín, Mayla Perdomo, Diego Muñoz,

Álvaro Duarte, Edwin Vásquez; Hospital San Andrés de Tumaco: Camilo Ortiz, Bernardo Ayala, Hernán Delgado, Gloria Benavides, Lorena Rosero; Fundación Clínica Valle del Lili: Jorge Mejía-Mantilla, Ana Varela, Maríaisabel Calle, José Castillo, Alberto García; Clínica las Americas: Juan Ciro, Clara Villa, Roberto Panesso; Hospital General de Medellín: Luz Flórez, Argemiro Gallego; Hospital San Felix ESE: Fabián Puentes-Manosalva, Leonor Medina, Kelly Márquez; Hospital Universitario del Caribe: Adalgiza Reyes Romero, Ricardo Hernández, Julio Martínez; Hospital Universitario San Jorge: Wilson Gualteros; Hospital San Rafael Tunja: Zulma Urbina, Julio Velandia; Clínica La Estancia SA: Federico Benítez, Adolfo Trochez; Fundación Hospital San José de Buga: Andrés Villarreal, Pamela Pabón; Hospital Civil de Ipiales: Hernán Delgado; Hospital Universitario Departamental Nariño: Héctor López; Hospital Universitario del Valle: Laureano Quintero; Hospital Universitario de Neiva: Andrés Rubiano; Hospital Manuel Uribe Ángel: Juan Tamayo. *Cuba (575)*—Hospital Clínico-Quirúrgico Docente “Saturnino Lora”: Marjoris Piñera, Zadis Navarro, Deborah Rondón, Bárbara Bujan; Hospital General Universitario “Carlos Manuel de Céspedes”: Leonel Palacios, Daymis Martínez, Yalisa Hernández, Yaimara Fernández; Hospital Provincial Docente “Manuel Ascunce Domenech”: Eugenio Casola; Hospital Universitario “Arnaldo Milión Castro”: Rodolfo Delgado, Carlos Herrera, Migdacelys Arbolaéz, Mario Domínguez; Hospital Universitario “Dr Gustavo Aldereguía Lima”: Marcos Iraola, Omar Rojas, Alba Enseñat; Hospital Abel Santamaría Cuadrado: Irene Pastrana, Daniel Rodríguez, Sergio Álvarez de la Campa; Hospital Miguel Enríquez: Thorvald Fortún; Hospital General Calixto García: Martha Larrea; Hospital Antonio Luaces Iraola: Lensky Aragón; Hospital Provincial Docente VI Lenin: Aida Madrazo. *Czech Republic (17)*—Research Institute for Special Surgery and Trauma: Petr Svoboda. *Ecuador (1198)*—Hospital Luis Vernaza: Mario Izurieta, Alberto Daccach, Mónica Altamirano, Antonio Ortega, Bolívar Cárdenas, Luis González; Hospital José Carrasco Arteaga: Marcelo Ochoa, Fernando Ortega, Fausto Quichimbo, Jenny Guiñanzaca; Hospital de Niños Dr Roberto Gilbert Elizalde: Ines Zavala, Sayra Segura; Hospital Naval Guayaquil: Johnny Jerez; Hospital Alcivar: Daniel Acosta; Hospital “Dr Rafael Rodríguez Zambrano”: Fabián Yáñez; Clínica De Especialidades Medicas “San Gregorio”: Rubén Camacho. *Egypt (2234)*—Mataria Teaching Hospital: Hussein Khamis, Hossam Shafei, Ali Kheidr, Hani Nasr, Moetaz Mosaad, Safwat Rizk; Suez Canal University: Hesham El Sayed, Taha Moati, Emad Hokkam; Aswan Teaching Hospital: Mamdouh Amin, Hany Lowis, Medhat Fawzy, Nabil Bedir, Mohamed

Aldars. *El Salvador (84)*—Hospital Nacional Rosales: Virginia Rodríguez, Juan Tobar, Jorge Alvarenga. *Georgia (1783)*—Tbilisi State University Clinical Hospital 'I Javakhishvili': Budu Shalamberidze, Elza Demuria, Nikoloz Rtveliashvili, Gocha Chutkerashvili, David Dotiashvili; Tbilisi First Hospital, University Clinic, Neurosurgery Center: Tamar Gogichaishvili, George Ingorokva, David Kazaishvili, Besik Melikidze, Natia Iashvili; Tbilisi City Hospital #1: Gia Tomadze, Manana Chkhikvadze, Leri Khurtsidze, Zviad Lomidze, Diana Dzagania; Tbilisi State Medical University ER Department: Nikoloz Kvachadze, Giorgi Gotsadze, Vakhtang Kaloiani; Institute of Critical Care Medicine: Nino Kajaia. *Ghana (136)*—Korle Bu Teaching Hospital: Jonathan Dakubo, Simon Naaeder, Priscilla Sowah; Nyinahin Government Hospital: Adamu Yusuf, Alhaji Ishak; Sogakope District Hospital: Paul Selasi-Sefenu; Methodist Hospital Wenchi: Ballu Sibiri; Effia Nkwanta Regional Hospital: Sampson Sarpong-Peprah; Saint Theresa's Hospital: Theodore Boro. *India (4768)*—Medical Trust Hospital Kochi: Kanjithanda Bopaiah, Kishore Shetty, Raja Subbiah, Lukman Mulla, Anand Doshi; Christian Medical College Ludhiana: Yashbir Dewan, Sarvpreet Grewal, Pradipta Tripathy, Jacob Mathew, Bharat Gupta; Aditya Neuroscience Centre: Anil Lal, Majulie Choudhury; Sri Sai Hospital: Sanjay Gupta, Smita Gupta, Arun Chug; Care Hospital: Venkataramana Pamidimukkala, Palaniappan Jagannath, Mohan Maharaj, Ramaraju Vommi, Naresh Gudipati; North Bengal Neuro Research Centre: W H Chhang; Sheth VS General Hospital and NHL Municipal College: Pankaj Patel, Nilay Suthar, Deepa Banker, Jyotish Patel; LTM Medical College and General Hospital: Satish Dharap, Ranjeet Kamble, Shraddha Patkar, Sushil Lohiya; Government Medical College and Associated Hospitals Jammu: Rakesh Saraf, Dinesh Kumar, Satish Parihar, Rahul Gupta; MKCG Medical College: Rasananda Mangual, Alagumuthu, Don Kooper, Chinmaya Mohapatra; Christian Medical College Hospital Vellore: Suresh David, Wesley Rajaleelan, Appas; KLE Hospital and Medical Research Centre: Ashok Pang, Vivek Saraf, Santhosh Chikareddy; NKP Salve Institute of Medical Sciences and Lata Mangeshkar Hospital: Sushil Mankar, Anil Golhar, Rahul Sakhare, Nilesh Wagh; Sanjivani Diagnostics and Hospital: Anil Lal, Dhiman Hazarika; Parkar Hospital: Pratyush Chaudhuri; Jeevan Jyoti Hospital and Research Centre: Prakash Ketan; Mansarovar Hospital: Govindbhai Purohit, Yogesh Purohit, Mandakini Pandya; Postgraduate Institute of Medical Science Rohtak: Rakesh Gupta, Shashi Kiran, Saurab Walia; Goyal Hospital Jalna: Sonam Goyal, Sidhant Goyal, Satish Goyal; Government Medical College Chandigarh: Sanjay Gupta, Ashok Attri, Rajeev

Sharma; Oberai Hospital: Ashok Oberai, Mahesh Oberai, Supriya Oberoi; Rajeev Gandhi Memorial Hospital and Research Centre: Gajendra Kant Tripathi; Calicut Medical College Hospital: Vijayan Peettakkandy, Premkumar Karuthillath, Pavithran Vadakammuriyil; Krishnamai Medical and Research Foundation's NIKOP Hospital: Jalindar Pol, Sunita Pol, Manisha Saste; St Stephen's Hospital: Subrat Raul, Shashi Tiwari, Neileino Nelly; Government Rajaji Hospital: M Chidambaram; Medical College Trivandrum: Viswanathan Kollengode, Sam Thampan; Sanjeevani Hospital: Sunder Rajan, Sushrut Rajan; Kamineni Hospital: Subodh Raju, Renuka Sharma; Sri Sakthi Hospital: Subbiah Venkatesh Babu, Chellappa Sumathi; Bhattacharya Orthopaedic and Related Research Centre: Protyush Chatterjee, Alok Agarwal; Sushrut Hospital: Hemant Magar, Meera Magar; All India Institute of Medical Sciences: Manmohan Singh, Deepak Gupta; GM Hospital (P): LTD: Anil Lal, Kamal Haloi; Government Medical College and Superspeciality Hospital Nagpur: Varsha Sagdeo, Pramod Giri; Government Medical College New Civil Hospital: Nimesh Verma, Ravi Jariwala, Ashish Goti; Chikitsa Hospital: Aman Prabhu-Gaonkar, Sagar Utagi; Apollo Health City: Mahesh Joshi, Ruchit Agrawal; Apex Neurotrauma and Superspeciality Hospital: Gopal Sharma, Gurvinder Saini; Neuro Center Gola Ghat: Vinod Tewari; NSCB Medical College: Yad Yadav, Vijay Parihar; BGS Global Hospital: Neelam Venkataramana, Shailesh Rao; Chettinad Hospital and Research Institute: Narayana Reddy, SG Chander; Sir Sayajirao General Hospital and Medical College Baroda: Virsing Hathila; Goyal Hospital and Research Centre Jodhpur: Vithal Das; Krishna Surgical Hospital and Trauma Care Centre: Kantibhai Agaja; Nizam's Institute of Medical Sciences: Aniruddh Purohit; Niramay Hospital: Akilesh Lahari; Apex Hospital Bhopal: Rajesh Bhagchandani; Dr Jeyasekharan Medical Trust: Bala Vidyasagar; Himalayan Institute of Medical Sciences: P K Sachan; Apollo Gleneagles Hospitals: Tanmoy Das; Civil Hospital Gandhinagar: Sharad Vyas; Sukhdev Raj Soini Hospital: Sujoy Bhattacharjee; Sancheti Institute for Orthopaedics and Rehabilitation: Parag Sancheti; St James Hospital: T Manoj; Al Shifa Hospital: Mubarak Moideen; Anant Institute of Medical Sciences: Kailash Pansey; Vinayaka Mission Hospital: V P Chandrasekaran; Gauhati Medical College and Hospital: Kabul Saikia; Krishna Hospital and Medical Research Centre: Hoshedar Tata; Ruby Hall Clinic: Sanjay Vhora; Shreejee Hospital: Aniket Shah; Nazareth Hospital: Gordon Rangad; Ganga Hospital: S Rajasekaran; Vadamalayan Hospitals: S T Shankarlal; Devadoss Multispeciality Hospital: Sathish Devadoss; KIOT Hospital: M Saleem; Baby Memorial

Hospital: Haroon Pillay; Bethany Hospital: Zulfiqer Hazarika; Suretech Hospital and Research Centre: Parikshit Deshmukh; Surya Hospital: S P Murugappan; Apollo Clinic Varanasi: Amit Jaiswal; Fortis Escorts Hospital: Deepak Vangani; Gokul Hospital and Trauma Centre: Prakash Modha; International Hospital Assam: Chawngrolien Chonzik; Lifeline Multispeciality Hospital: Megha Praveen; Meenakshi Mission Hospital and Research Centre: Vijaya Sethurayar; MOSC Medical College Hospital: Sojan Ipe; MS Ramaiah Memorial Hospital: Naresh Shetty; Saykhedkar Hospital and Research Centre: Aniket Shah; Shanti Mukand Hospital: R P Gupta; Shri KM Memorial Jain Heart and General Hospital: Vinod Jain; Usha Hospital: Ketan Shah. *Indonesia (706)*—Soebandi Hospital Jember: Moch Dwikoryanto; Sanglah General Hospital: Nyoman Golden, Kuning Atmadjaya, Ketut Wiargitha, Ketut Sudiasa, Gede Suwedagatha; Saiful Anwar General Hospital: Farhad Bal'arif; Dr Soetomo General Hospital: Vicky Budipramana, Tabrani, Agung Lemuel; Cipto Mangunkusumo Hospital: Susilo Chandra; Muhammadiyah Lamongan Hospital: Faisol Ama. *Iran (134)*—Nemazi Hospital: Ehsan Sherafatkazemzadeh, Ehsan Moradi, Alireza Sheikhi; Erfan Hospital: Ali Ziaee, Ahmad Fanaei; Loqman Medical Center: Esmaeil Hajinasrollah; Imam Hosain Hospital: Afshin Amini. *Iraq (392)*—Diwanayah College of Medicine: Bassim Mohammad, Najah Hadi. *Italy (57)*—Spedali Civili di Brescia: Giovanna Perone, Elena de Peri; Azienda Ospedaliera Di Parma: Annalisa Volpi. *Jamaica (11)*—University Hospital of the West Indies: Jean Johnson. *Japan (9)*—Fukuoka University Hospital: Masayoshi Abe. *Kenya (31)*—Kenyatta National Hospital: Vincent Mutiso, Benjamin Okanga; Kapenguria District Hospital: Daniel Ojuka. *Malaysia (216)*—Hospital University Science Malaysia: Baharudin Abdullah, Hishamuddin Rahman, Yazid Noh; Sungai Buloh Hospital: Sabariah Jamaluddin, Hasnah Dawal; University of Malaya Medical Centre: April Roslani, Chee-Wei Law, P Devashanti; Hospital Tengku Ampuan Rahimah: Yusof Wahab, Shanta Velaiutham; Ampang Hospital: Ridzuan Dato. *Mexico (479)*—Hospital General Regional 25: Jorge Loría, Erandy Montes, Eduardo Gómez, Víctor Cazales, Paúl Bautista; Hospital Gustavo Roviroso: Raúl Bautista, David Ahumada, Edwin Hernández, Germahin Velásquez; Hospital General de Uruapan "Dr Pedro Daniel Martínez": Patricia Ortega, Gabriel Lira, Francisco Estrada; Hospital General Ecatepec Las Américas: Jorge Martínez; Hospital General La Perla: Juan Martínez; Hospital General de Ecatepec "Dr José María Rodríguez": Julio Casasola. *Nigeria (2053)*—National Hospital Abuja: Oluwole Olaomi, Yari Abubakar, Kanati Apollo, Olawale Badejo, Ogemdi Ihekire; University of Benin Teaching Hospital: Pius Iribhogbe, Olugbenga Oludiran, Emmanuel Obeta, Collins Okojie, Ernest Udefiagbon; Obafemi Awolowo University Teaching Hospitals: Edward Komolafe, Patrick Olaleye, Tochukwu Uzochukwu, Uvie Onakpoya; Irrua Specialist Teaching Hospital: Andrew Dongo, Osas Uhumwagho, Ehi Eighemerio, Eghosa Morgan; Olabisi Onabanjo University Teaching Hospital: Lateef Thanni; University College Hospital Ibadan: Adefemi Afolabi, Titilade Akinola, Adeyinka Ademola, Olusola Akute; Ahmadu Bello University Teaching Hospital: Lawal Khalid, Lawal Abubakar, Muhammad Aminu, Mike Ogirima; Baptist Medical Centre: Ambrose Attansey, Durodola Michael, Olaniran Aremu; University of Ilorin Teaching Hospital: Odebode Olugbenga, Udoffa Ukpong, Yusuf Salman; Enugu State University Teaching Hospital: Nene Obianyo, Chinenye Ani, Roderick Ezeadawi; LAUTECH Teaching Hospital: Oluwadiya Kehinde, Agodirin Olaide; Federal Medical Centre Makurdi: Andrea Jogo, Terna Bitto; Nnamdi Azikiwe University Teaching Hospital: Stanley Anyanwu, Okechukwu Mbonu; Lagos State University Teaching Hospital: Mobolaji Oludara, Michael Somoye; Usmanu Danfodiyo University Teaching Hospital: Bello Shehu, Nasir Ismail; National Orthopaedic Hospital Enugu: Amechi Katchy; University of Calabar Teaching Hospital: Rowland Ndoma-Egba, Ngim Grace-Inah; University of Abuja Teaching Hospital: Zumnan Songden, Abdulrahman Abdulaheem; University of Uyo Teaching Hospital: Akpan Otu, Timothy Nottidge; Federal Medical Centre, Yenagoa: Domingo Inyang, David Idiapho; Seventh Day Adventist Hospital: Herb Giebel; Federal Medical Centre Birnin-Kebbi: Ramatu Hassan; Abia State University Teaching Hospital: Adeyinka Adisa; Wesley Guild Hospital: Akinbolaji Akinkuolie; Federal Medical Centre, Umuahia: Kalu Okam; University of Maiduguri Teaching Hospital: Abubakar Musa; National Orthopaedic Hospital, Igbobi: Ignatius Falope; University of Nigeria Teaching Hospital Enugu: John Eze. *Peru (452)*—Hospital Regional Docente de Trujillo: José Caballero, Wenceslao Azabache, Oscar Salirrosas; Hospital Nacional Hipolito Unanue: Alonso Soto, Elfi Torres, Gloria Ramírez, Mónica Pérez; Clínica Santa Ana: Cesar Malca; Hospital La Caleta: Juan Velez; Hospital Nacional Sergio E Bernales: Raul Yopez; Hospital de Apoyo de Sullana: Hernan Yupanqui; Hospital IV Essalud Huancayo: Pedro Lagos; Hospital Nacional Arzobispo Loayza: Diana Rodriguez; Hospital Municipal Los Olivos: Jorge Flores; Hospital Jose Cayetano Heredia: Anselmo Moya; Hospital Nacional

Carlos Alberto Seguin Escobedo: Alejandro Barrionuevo; Hospital Nacional Dos De Mayo: Marco Gonzales-Portillo; Hospital Nacional Cayetano Heredia: Edgar Nunez. *Saudi Arabia* (70)—King Khalid University Hospital: Abdelazeem Eldawlatly, Mohammed Al Naami, Bilal Delvi; King Khalid National Guard Hospital: Walid Alyafi. *Serbia* (1)—Klinicki Centar Srbije: Branko Djurovic. *Singapore* (2)—National Neuroscience Institute: Ivan Ng. *Slovakia* (38)—FNsP Ružinov: Aktham Yaghi; NsP Poprad: Anton Laincz; NsP JA Reiman Hospital: Stefan Trenkler; Faculty Hospital F D Roosevelt: Jozef Valky. *South Africa* (76)—Dr George Mukhari Hospital: Mphako Modiba, Peter Legodi, Thomas Rangaka; George Provincial Hospital: Lee Wallis. *Spain* (23)—Hospital Universitario Virgen del Rocío: Ángeles Muñoz; Hospital Ramón y Cajal de Madrid: Ana Serrano; Hospital Universitario Germans Trias i Pujol: Maite Misis; Hospital Torrecardenas: Martin Rubi; Hospital Universitario Virgen de la Victoria: Victoria de la Torre. *Sri Lanka* (103)—National Hospital of Sri Lanka: Ranjith Ellawala, Samitha Wijeratna, Lukshri Gunaratna, Crishantha Wijayanayaka. *Tanzania* (64)—Muhimbili Orthopaedic Institute: Kitugi Nungu, Billy Haonga, Grenda Mtapu. *Thailand* (903)—Khon Kaen Regional Hospital: Surakrant Yutthakasemsunt, Warawut Kittiwattanagul, Parnumas Piyavechvirat, Tawatcahi Impool, Santipong Thummaraj; Pattani Hospital: Rusta Salaeh; Suratthani Hospital: Sakchai Tangchitvittaya; Bhumibol Adulyadej Hospital: Kamol Wattanakrai, Chatchai Soonthornthum, Teerasak Jiravongbunrod; Lampang Hospital: Surasak Meephant; Rayong Hospital: Pusit Subsompon; Roi-Et Hospital: Phaiboon Pensuwan; Phrae Hospital: Wichanrat Chamnongwit. *Tunisia* (36)—Hospital Habib Thameur: Zouheir Jerbi, Abderraouef Cherif. *UK* (135)—University Hospital of North Staffordshire: Mark Nash; Royal London Hospital: Tim Harris; Leicester Royal Infirmary: Jay Banerjee; Nottingham University Hospitals NHS Trust: Ramzi Freij; Frenchay Hospital: Jason Kendall; Countess of Chester Hospital: Stephen Moore; Hull Royal Infirmary: William Townend; Royal Sussex County Hospital: Rowland Cottingham; Derby Hospitals NHS Trust: Dan Becker; Bedford Hospital NHS Trust: Stuart Lloyd; Royal Liverpool University Hospital: Peter Burdett-Smith; Colchester General Hospital: Kazim Mirza; Royal Lancaster Infirmary: Andrew Webster; Worthing Hospital: Suzanne Brady, Amanda Grocutt; Darent Valley Hospital: John Thurston; Hope Hospital: Fiona Lecky; Northern General Hospital: Steve Goodacre. *Zambia* (62)—University Teaching Hospital, Lusaka: Yakub Mulla, Dennis Sakala; Nchanga North General Hospital: Charles Chengo.

CONFLICTS OF INTEREST

Members of the Writing Committee declare that they have no conflicts of interest.

ACKNOWLEDGMENTS

Telephone randomisation system, unblinding, and adverse event reporting service was provided by the CTSU, University of Oxford, Oxford, UK. Randomisation codes were generated by Alan Young of CTSU. The London School of Hygiene and Tropical Medicine supported the core trial coordinating staff during the first year of the trial set-up. Funding for the run-in stage was provided by J P Moulton Charitable Foundation and the BUPA Foundation. A grant-in-aid for purchasing the tranexamic acid and placebo was provided by Pfizer. The main phase of this trial was funded by the UK NIHR Health Technology Assessment programme and will be published in full in the Health Technology Assessment journal series. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Department of Health.

The research was funded by the UK NIHR Health Technology Assessment Programme.

Pfizer Inc provided funding for the purchase of the tranexamic acid and the placebo.

1. Tranexamic acid is to be added to the WHO Model List of Essential Medicines following the recent review by the 18th Expert Committee on the Selection and Use of Essential Medicines. Their meeting report states: Addition of tranexamic acid injection for the treatment of adult patients with trauma and significant risk of ongoing haemorrhage. On the basis of the results of a very large trial of the use of tranexamic acid specifically for trauma patients - including those who have been in road traffic accidents, the Committee concluded that there is sufficient evidence to support the proposal that listing tranexamic acid may contribute to a reduction in this cause of death. See http://www.who.int/selection_medicines/complete_UNEDITED_TRS_18th.pdf further information.
2. The CRASH-2 Collaborators won the British Medical Journal Research Paper of the Paper of the Year 2011 Award.

REFERENCES

- Peden, M., McGee, K. and Sharma, G. The injury chart book: a graphical overview of the global burden of injuries. Geneva: World Health Organization, 2002.
- Gosselin, R. A, Spiegel, D. A, Coughlin, R., Zirkled, L. G. Injuries: the neglected burden in developing countries. *Bull World Health Organ* 2009; **87**: 246.
- Sauaia, A., Moore, F. A., Moore, E. E., *et al.* Epidemiology of trauma deaths: a reassessment. *J Trauma* 1995; **38**: 185–93.
- Lawson, J. H. and Murphy, M. P. Challenges for providing effective hemostasis in surgery and trauma. *Sem Hematol* 2004; **41**: 55–64.
- Henry, D. A., Carless, P. A., Moxey, A. J., *et al.* Anti-fibrinolytic use for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2007; **4**: CD001886.
- Okamoto, S., Hijikata-Okunomiya, A., Wanaka, K., Okada, Y. and Okamoto, U. Enzyme controlling medicines: introduction. *Semin Thromb Hemost* 1997; **23**: 493–501.
- Coats, T., Roberts, I. and Shakur, H. Antifibrinolytic drugs for acute traumatic injury. *Cochrane Database Syst Rev* 2004; **4**: CD004896.
- Baigent, C., Peto, R., Gray, R., Parish, S. and Collins, R. Large-scale randomized evidence: trials and meta-analyses of trials. In: Warrell DA, Cox TM, Firth JD, eds. Oxford Textbook of Medicine (5th edn). Oxford: Oxford University Press, 2010: 31–45.
- Perel, P., Edwards, P., Shakur, H. and Roberts, I. Use of the Oxford Handicap Scale at hospital discharge to predict Glasgow Outcome Scale at 6 months in patients with traumatic brain injury. *BMC Med Res Methodol* 2008, **8**: 72.
- Duley, L., Antman, K., Arena, J., *et al.* Specific barriers to the conduct of randomized trials. *Clin Trials* 2008; **5**: 40–48.
- Sterne, J., White, I., Carlin, J., *et al.* Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 2009; **338**: b2393.
- Rogers, A. and MacMahon, S. Systematic underestimation of treatment effects as a result of diagnostic test inaccuracy: implications for the interpretation and design of thromboprophylaxis trials. *Thromb Haemost* 1995; **73**: 167–71.
- Brohi, K., Cohen, M. J., Ganter, M. T., *et al.* Acute coagulopathy of trauma: hypo-perfusion induces systemic anticoagulation and hyper-fibrinolysis. *J Trauma* 2008; **64**: 1211–17.
- Ekbäck, G., Axelsson, K., Rytberg, L., *et al.* Tranexamic acid reduces blood loss in total hip replacement surgery. *Anesth Analg* 2000; **91**: 1124–30.
- Kauvar, D. S., Lefering, R. and Wade, C. E. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma* 2006; **60**: S3–11.
- Mock, C. N., Jurkovich, G. J., nii-Amon-Kotei, D., Arreola-Risa, C. and Maier, R. V. Trauma mortality patterns in three nations at different economic levels: implications for global trauma system development. *J Trauma* 1998; **44**: 804–12.
- Wyatt, J., Beard, D., Gray, A., Busuttill, A. and Robertson, C. The time of death after trauma. *BMJ* 1995; **310**: 1502.
- Fiechtner, B. K., Nuttall, G. A., Johnson, M. E., *et al.* Plasma tranexamic acid concentrations during cardiopulmonary bypass. *Anesth Analg* 2001; **92**: 1131–36.
- Horrow, J. C., Van Riper, D. F., Strong, M. D., Grunewald, K. E. and Parmet, J. L. The dose-response relationship of tranexamic acid. *Anesthesiology* 1995; **82**: 383–92.
- MRC CRASH Trial Collaborators. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ* 2008; **12**: 12.
- Maas, A. I., Steyerberg, E. W., Butcher, I., *et al.* Prognostic value of computerized tomography scan characteristics in traumatic brain injury: results from the IMPACT study. *J Neurotrauma* 2007; **24**: 303–14.
- Perel, P., Roberts, I., Bouamra, O., Woodford, M., Mooney, J. and Lecky, F. Intracranial bleeding in patients with traumatic brain injury: a prognostic study. *BMC Emerg Med* 2009, **9**: 15.
- Hogan, M. C., Foreman, K. J., Naghavi, M., *et al.* Maternal mortality for 181 countries, 1980–2008: a systematic analysis of progress towards Millennium Development Goal 5. *Lancet* 2010; **375**: 1609–23.
- Ferrer, P., Roberts, I., Sydenham, E., Blackhall, K. and Shakur, H. Anti-fibrinolytic agents in postpartum haemorrhage: a systematic review. *BMC Pregnancy Childbirth* 2009; **9**: 29.
- Shakur, H., Elbourne, D., Gülmezoglu, M., *et al.* The WOMAN Trial (World Maternal Antifibrinolytic Trial): tranexamic acid for the treatment of postpartum haemorrhage: an international randomised, double blind placebo controlled trial. *Trials* 2010; **11**: 40.