

Vertical Axillary Incision for Treatment of Breast Outer Quadrants Tumors

Taha Abd El-wahab Baiomy, Nawel El Sayed Hussein Abd El Hamid Matar,

Aisha Ahmad Ashtaiwe*, Joseph Rizk Israel

Onco-Surgery Unit, Department of Surgery, Faculty of Medicine, Zagazig University, Egypt

*Corresponding Author: Aisha Ahmad Ashtaiwe, Mobile : 00218910477032, Email: aishaahmad8519@gmail.com

ABSTRACT

Background: Breast cancer is the second most common cancer among women, with an estimated 3.5 million survivors as of 2015. There is continuous annual improvement of the overall survival of breast cancer patients, with 5-year overall survival estimates increasing from 84.6% to 90.9% over the previous two decades.

Objective: The aim of the current work was to improve aesthetic satisfaction and oncological outcome of patients with breast outer quadrant tumors.

Patients and Methods: This interventional study included a total of 18 cases with early breast cancer at upper and lower-outer quadrants of breast, attending at Onco-Surgery Unit, General Surgery Department, Faculty of Medicine, Zagazig University. This study was conducted between December 2020 to May 2021. All patients were subjected to demographic data taking, complete clinical examination, and investigations including imaging, biopsies, and histopathology.

Results: The Histopathological type of tumor, all of them had Invasive ductal carcinoma. Most of the tumors were (ER) and (PR) Positive with HER2/neo Negative. The mean duration of operation was 86.94 ± 10.45 (range 70–105) minute and the mean blood loss was 286 ± 12.34 (range 250-315) milliliter and all patients had no intraoperative complication.

Conclusions: It could be concluded that the vertical axillary incision is a safe and effective approach for the management of tumors of the outer quadrant of the breast; and cosmetically appropriate maneuver for these tumors.

Keywords: Breast Conserving Surgery (BCS), Breast cancer, Quadrant tumors.

INTRODUCTION

Breast cancer is the most common cancer among women. There is continuous annual improvement of the overall survival of breast cancer patients, with 5-year overall survival estimates increasing from 84.6% to 90.9% over the previous two decades⁽¹⁾.

The outer quadrant of the breast is the most common for breast cancer quadrant and axillary extension of the breast⁽²⁾.

Breast Conserving Surgery (BCS) is nowadays the standard surgical approach for the treatment of most cases of early breast cancer. During the last decades, the trend of breast conserving surgery has shifted from quadrantectomy, to large excision, to lumpectomy with the intent to minimize the excision volume in order to reduce breast shape deformities, asymmetry, and scar length⁽³⁾.

The possibility of carrying out this conservative treatment will depend, on one hand, the tumor volume compared to that of the breast (tumor volume / breast volume ratio), and on the other hand, tumor location⁽⁴⁾.

The possible distortion of breast shape and the following poor cosmetic result led surgeons to develop different surgical techniques in order to overcome the aesthetic discomfort for the patient⁽⁵⁾.

We performed this study to evaluate vertical axillary incision as an option for improving aesthetic satisfaction and oncological outcome in the management of breast outer upper and lower quadrant tumors.

PATIENTS AND METHODS

This interventional study included a total of 18 cases with early breast cancer at upper and lower-outer quadrants of breast, attending at Onco-Surgery Unit, General Surgery Department, Faculty of Medicine, Zagazig University. This study was conducted between December 2020 to May 2021.

Ethical Consideration:

Written informed consent was obtained from all participants and the study was approved by the Research Ethical Committee, Faculty of Medicine, Zagazig University. The work has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for studies involving humans.

Inclusion criteria:

Patients with early breast cancer at upper and lower-outer quadrants of breast provided that tumor pathology is invasive ductal carcinoma, less than 5 cm in the biggest diameter and the skin overlying it is not involved.

Exclusion criteria:

Patients not eligible to breast conservative treatment. Patients presenting with advanced malignant that is (More than 5 cm in diameter and the skin overlying it is involved). Patients refuse to have breast conservative treatment.



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY-SA) license (<http://creativecommons.org/licenses/by/4.0/>)

Preoperative investigations:

All patients were subjected to demographic data taking, complete clinical examination, investigations included imaging, biopsies, and histopathology. Bilateral breast mammo- ultrasound was performed. The breast was assessed in the four main quadrants, namely outer upper, outer inner, lower outer, and lower inner quadrants and the retro areolar space. Any lesions identified during the examination were marked as a breast 'o clock position for future follow up sessions. US-Guided Core needle biopsy was performed for all patients.

For histopathological exam and assessment of PR, ER, HER2/ neo in specimen for all patients.

Routine Pre-operative Labs; complete blood count (CBC), PT, PTT and INR, random blood sugar, Liver and Kidney function tests and cardiac investigation if needed.

Operative:

Surgical technique

The design of the incision was done either at the night of the surgery or in the operating room before induction of an anesthesia. The patient was instructed to sit and her arm is abducted 90 degree to expose the axillary hollow part. A vertical line was drawn just behind anterior axillary fold starting at the intersection of long head of biceps and pectoralis major then extended downward for 7-10 cm according to the site of tumor. Then the patient was placed in supine position with arms abducted to 110 degrees and a pillow was placed under the patient shoulder & back to elevate it for more exposure. The incision was done according to the mark through skin & subcutaneous, then dissection was advanced medially through the plane between the breast tissue and the subcutaneous fat. The upper flap was undermined until reaching beyond the tumor with the enough safety margin needed. Then resection of the tumor with safety margin all around was done and continued laterally towards the axilla. Then axillary lymph node dissection proceeded as the classic operation. After retrieval of specimen, hemostasis was ensured, suction drain was inserted in tumor bed & axilla. Closure of the wound.

Postoperative follow up

Early post-operative follow up during two weeks after ensuring the general patient stability post-operatively; The Patients were examined for wound infection, seroma or nerve injury. The safety of post-operative tumor margin in histopathological examination of the specimen.

Late Post-operative Follow up during three month. Assessment of both patient & surgeon satisfaction regarding the Cosmetic outcome. Any limitation in shoulder mobility.

Statistic analysis

The collected data were coded, processed and analyzed using the SPSS (Statistical Package for Social Sciences) version 22 for Windows® (IBM SPSS Inc, Chicago, IL, USA). Data were tested for normal distribution using the Shapiro Walk test. Qualitative data were represented as frequencies and relative percentages. Chi square test (χ^2) to calculate difference between two or more groups of qualitative variables. Quantitative data were expressed as mean \pm SD (Standard deviation). Independent samples t-test was used to compare between two independent groups of normally distributed variables (parametric data). P value < 0.05 was considered significant.

RESULTS

Table (1): Patient personal data.

Item	Frequency (N=18)
Mean age (year)	54.7 \pm 8.93
Occupation	
Employee	5
House wife	13
Special habits	
No habits	18
Parity	
Nulli-para	3
Para 1	3
Multi-para	12
Breast feeding	
Yes	13
No	5
Oral contraception use	
Yes	4
No	14
Menstrual state	
Pre-menopausal	5
Post-menopausal	13

Table 1 shows that the mean age of patients in the study was 54.7 \pm 8.93 years with most of them (72.2 %) were housewives. That most of the patients were Multipara having breast fed their babies, most of them didn't have contraceptive pills and at the study, most of them were post-menopause.

Table (2): Pathological data.

Item	Frequency (N=18)
Laterality	
LT	12
RT	6
Quadrant site of tumour	
UOQ	14
LOQ	4
Size of tumour	
Mean	2.56 ± 0.71 cm
Range	1.4 : 3.8 cm
LN status	
N0	11
N1	7
Presence of skin changes	
No changes	18
PR receptor status	
Positive	12
ER receptor status	
Positive	13
Her2/neo receptor status	
Positive	3

Table 2 shows that the mean size of tumor was 2.56±0.71 with most of them were in the upper outer quadrant on the left site and most of them had no palpable lymph node. that the Histopathological type of tumor all of them were Invasive ductal carcinoma. Most of the tumors in our study were (ER) and (PR) Positive with HER2/neo Negative.

Table (3): Operative data.

Item	Frequency (N=18)
Duration of the operation (min)	86.94 ± 10.45
Amount of blood loss (ml)	286 ± 12.34
Intra-operative complications	
No	18

The only recorded complication in our study is seroma in 4 cases table 5.

Table (4): Post-operative follow up data.

Item	Frequency (N=18)
Safety of resection	
Free safety margin	18
Complications	
Seroma	4
Wound infection	0
Wound dehescence	0
Patient satisfaction regarding cosmeses	
Excellent	15
Good	3
Surgeon satisfaction regarding cosmeses	
Excellent	18
Good	0

All the tumors were resected with free surgical margin. that the most of the patient's evaluation for cosmetic outcome by patients showed complete

satisfaction (83.3%), while all of the patients evaluation by surgeons was excellent (100%) table 4.

DISCUSSION

In the current study the mean age of patients was 54.7±8.93 years. This came in agreement with **Bordoni et al.**⁽⁶⁾ who reported the patients' mean age was 57.5 years. In contrast **Haytham et al.**⁽⁷⁾ reported younger mean age 46.3 ± 7.1. Also **Elnahas et al.**⁽⁸⁾ revealed that the mean age of patients of vertical mammoplasty technique was 45 years. While **Foissac et al.**⁽⁴⁾ reported older mean age of study participants with mean age 62.34 years. All these findings are consistent with the fact that breast cancer as most of other malignancies affects female at 5th & 6th decades with recent tendency to affect younger ones at 4th decade of life.

The current study showed that multi-parous represented (66.6%), (72.2%) of the patients practised breastfeeding, (77.7%) didn't receive Oral contraceptive and (72.2%) were post-menopausal. While **Redondo et al.**⁽⁹⁾ reported similar findings, that parity was reported in (82.9%), breastfeeding (65.8%) and (89.6%) of the patients were post-menopausal.

Several case-control and cohort studies have examined the association between parity and breastfeeding and the risk of Triple negative breast cancer. Although an inverse association of parity with risk of ER+ breast cancer has been found by many studies, studies of ER- breast cancer indicate a positive, risk-enhancing association with parity^(10,11).

Fortner et al.⁽¹²⁾ found that breastfeeding was inversely associated with ER- breast cancer, no association was observed for the ER+ subtype. In contrast, parity was associated with lower risk of ER+ breast cancer, but not associated with ER- disease.

In this study, most lumps (77.8%) located at UOQ of breast in agreement with **Haytham et al.**⁽⁷⁾ who investigated the lateral mammary sulcus incision as an oncoplastic approach for laterally located breast cancer the study reported that (83.6%) of cases had their tumors located in the upper outer quadrant. Also, **Acea-Nebril et al.**⁽¹³⁾ reported that UOQ tumor was the most frequent site for breast cancer (58.41%).

In the current study, all lesions were not attached to the skin and the mean size was 1.4-3.8 cm in agreement with **Elnahas et al.**⁽⁸⁾ who reported that the tumor size ranged from 1.5- 3.5 cm. Also, in agreement with our findings **Haytham et al.**⁽⁷⁾ reported that mean size 1-3.7cm.

In the current study the majority of patients (61.1%) were without lymph node involvement (N0), and those with lymph node involvement (N1) represented (38.9%).This were concordant with **Elnahas et al.**⁽⁸⁾ who reported that the lymph nodes (N0) were (73.21%) and involved lymph node (N1) were (26.7%).

Regarding pathological finding in our study, all the patients had invasive ductal carcinoma. Supporting

our results **Foissac et al.** ⁽⁴⁾ reported all cases had ductal carcinoma with a majority of grade I (50%) and II (26.6%) infiltrating ductal carcinoma. Furthermore **Chen et al.** ⁽¹⁴⁾ reported the Histological types of breast cancer in study participants were 14 out of 16 had invasive ductal carcinoma (87.5%) and one case mucinous breast cancer and one case ductal carcinoma in situ. and this goes with the fact that Ductal Carcinoma is more prevalent than other variants.

In our study, estrogen receptor was expressed among patients (72.2%) and Progesterone receptor was positive in (66.7%) patients, while Human epidermal growth factor receptor/2 was expressed among (16.7%) patients in agreement with **Chen et al.** ⁽¹⁴⁾ who reported that estrogen receptor was Positive in (62.5%) and Progesterone receptor was Positive in (62.5%) while Human epidermal growth factor receptor/2 was expressed only in (12.5%). **Bordoni et al.** ⁽⁶⁾ also reported 64.8% were positive for estrogen and progesterone receptors and 16% were positive for Human epidermal growth factor receptor/2.

In our study the mean duration of operation (per minutes) was 86.94 ± 10.45 minute and the mean Blood loss (per milliliter) was 286 ± 12.34 . None of patients had Intraoperative complication. but in the study of **Acea-Nebril et al.** ⁽¹³⁾ the mean surgical time was 51.67 ± 14.62 minutes which is considerable shorter than ours

In the current study, all patients had tumor-free surgical margins, and no further oncological operations were required. This came in agreement with **Barnea et al.** ⁽¹⁵⁾ who investigated oncoplastic breast reduction technique using a vertical scar superior-medial pedicle pattern for immediate partial breast reconstruction. He reported that all patients had tumor-free surgical margins, and no further oncological operations were required.

In this study, seroma was developed in four (22.2%) patients and was treated by frequent aspiration. **Abdelhamid et al.** ⁽¹⁶⁾ who reported (11.1%). **Elnahas et al.** ⁽⁸⁾ investigated multiple surgical techniques in a study included 56 patients of early-stage breast cancer. They were divided into 4 groups according to the surgical technique; Group IV vertical mammoplasty technique had slightly higher complications but without statistical significance 2 cases complained of wound dehiscence, two cases complained of seroma formation and one case had wound infection.

In **Hyatham et al.** ⁽⁷⁾ study conducted on 122 female patients with breast cancer who were suitable for BCS to assess the lateral mammary sulcus incision as an oncoplastic approach for laterally located breast cancer reported Surgical site infection was the most frequently encountered complication 9.8% and treated medically without the need for surgical intervention. Fat necrosis occurred in 4.9%, diagnosed by radiological study (either ultrasonography or mammography) and confirmed by core needle biopsy. Seroma encountered in 9% and treated by frequent aspiration. Complications

were statistically significant with increased body mass index.

In our work, post-operative limitation in shoulder mobility was not detected in any cases.

In the current study both the surgeons and the patients reported a high degree of satisfaction with the surgical outcome. Cosmetic outcome as reported by patients revealed that, 83.3% of patients reported excellent outcome, 16.7%, of patients reported good outcome, and cosmetic outcome as reported by surgeons revealed that, 100% of surgeon reported excellent outcome.

The results were nearly agreed with **Foissac et al.** ⁽⁴⁾ who reported (86.6%) of patients considered the quality of their scar as excellent with (3.4%) judged their result to be average due to a significant scar enlargement, and (10%) good with considerable satisfaction. None of the patients asked for another surgery for aesthetic reasons. Likewise, no deformation in areolar complex or sequelae of the breast was noted by the examiner throughout the follow-up of patients. The surgeons who made this incision were satisfied with the surgical approach authorized by the incision and the uniqueness of the incision.

The results were also agreed with **Abdelhamid et al.** ⁽¹⁶⁾ who reported (66.6%) of patients showed excellent results, (27.7%) of patients showed good results. One (5.5%) patient showed satisfactory result.

CONCLUSION

The vertical axillary incision is a safe and effective approach for the management of tumors of the outer quadrant of the breast; and cosmetically appropriate maneuver for these tumors.

Financial support and sponsorship: Nil.

Conflict of interest: Nil.

REFERENCES

1. **Howlader N, Noone A, Krapcho M et al. (2019):** SEER Cancer Statistics Review, 1975-2016, National Cancer Institute. Bethesda, MD, /, based on November 2018 SEER data submission, posted to the SEER web site. Available from: https://seer.cancer.gov/csr/1975_2016.
2. **Sisti A, Huayllani M, Boczar D et al. (2020):** Breast cancer in women: A descriptive analysis of the national cancer database. *Acta Bio Medica: Atenei Parmensis*, 91(2): 332-36.
3. **Calabrese C, Casella D, Di Taranto G et al. (2018).** Oncoplastic conservative surgery for breast cancer: long-term outcomes of our first ten years experience. *Eur Rev Med Pharmacol Sci.*, 22(21): 7333-42.
4. **Foissac R, Ihrari T, Cegarra-Escolano M et al. (2014):** Incision en «V» axillaire dans la prise en charge des tumeurs du quadrant supéro-externe et du prolongement axillaire du sein: étude rétrospective et description technique. In *Annales*

de Chirurgie Plastique Esthétique. Elsevier Masson. Pp. 320-326.

5. **Bordoni D, Cadenelli P, Rocco N et al. (2017).** Oncoplastic resection of breast cancers located in the upper-inner quadrants: a safe and effective surgical technique. *Eur J Plast Surg.*, 41(2): 157-164
6. **Bordoni D, Cadenelli P, Ornelli M et al. (2019):** The axillary flap in oncoplastic resection of breast cancers located in the upper-outer quadrants: a new surgical technique. *BMC Surgery*, 18(1): 1-7.
7. **Haytham M, Sharaan M, Gebaly A et al. (2020):** Oncoplastic Surgery for Laterally Located Breast Cancer: A New Approach. *Clin Surg.*, 5: 2790-95.
8. **Elnahas W, Fady T, Hegazy M (2011).** Oncoplastic resection of upper outer quadrant breast cancer. *Egyptian Journal of Surgery*, 30 (2): 321-26.
9. **Redondo C, Gago-Domínguez M, Ponte S et al. (2012).** Breast feeding, parity and breast cancer subtypes in a Spanish cohort. *PloS One*, 7(7): 40543-47.
10. **Phipps A, Chlebowski R, Prentice R et al. (2011):** Reproductive history and oral contraceptive use in relation to risk of triple-negative breast cancer. *J Natl Cancer Inst.*, 103: 470-477.
11. **Palmer J, Boggs D, Wise L et al. (2011):** Parity and lactation in relation to estrogen receptor negative breast cancer in African American women. *Cancer Epidemiol Biomarkers Prev.*, 20: 1883-1891.
12. **Fortner R, Sisti J, Chai B et al. (2019):** Parity, breastfeeding, and breast cancer risk by hormone receptor status and molecular phenotype: results from the Nurses' Health Studies. *Breast Cancer Research*, 21(1): 1-9.
13. **Acea-Nebril B, García-Novoa A, Cereijo-Garea C et al. (2019):** Single-Incision Approach for Breast-Conserving Surgery: Effectiveness, Complications and Quality of Life. *Annals of Surgical Oncology*, 26(8): 2466-2474.
14. **Chen Y, Xu J, Liang Y et al. (2020):** A challenging therapeutic method for breast cancer: Non-lipolytic endoscopic axillary surgery through periareolar incisions. *Oncology Letters*, 19(6): 4088-4092.
15. **Barnea Y, Inbal A, Barsuk D et al. (2014):** Oncoplastic reduction using the vertical scar superior-medial pedicle pattern technique for immediate partial breast reconstruction. *Canadian Journal of Surgery*, 57(4): 134-38.
16. **Abdelhamid M, Alkilany M, Lotfy M (2018):** Lazy lateral technique: an innovative approach for upper outer quadrant breast cancer near the anterior axillary fold. *The Egyptian Journal of Surgery*, 37(1): 1-5.