Letter to the Editor Beuy J. et al. 683

# LETTER TO THE EDITOR

# Rainfall and Development of Zika Virus

# Beuy Joob<sup>1\*</sup>, Viroj Wiwanitkit<sup>1</sup>

#### OPEN ACCESS

Citation: Beuy Joob, Viroj Wiwanitkit. Rainfall and Development of Zika Virus. Ethiop J Health Sci.2017;27(6):683. doi:http://dx.doi.org/10.4314/ejhs.v27i6.

1/

Received: May 4, 2017 Accepted: May 10, 2017 Published: November 1, 2017

Copyright: © 2017 Beuy J. et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: None

Competing Interests: The authors declare that this manuscript was approved by all authors in its formand that no competing interest exists.

### **Affiliation and Correspondence:**

<sup>1</sup>Sanitationa 1 Medical Academic Center, Bangkok Thailand <sup>2</sup>TB/HIV Research Directorate, Ethiopian Public Health Institute, Addis Ababa, Ethiopia

<sup>2</sup>Viisitng professor, Hainan Medical University, China

Dear Editor,

We read the publication on "Rainfall and Development of Zika Virus" with a great interest (1). Tosepu mentioned that "there is a relationship between the incidence of vectorborne diseases with rainfall" (1). In fact, this is not a new finding. The relationship between rainfall and incidence of arbovirus disease such as dengue is well demonstrated (2). For Zika virus an infection, a similar observation can be expected. A recent report from Thailand can also show the expected pattern of the prevalence of Zika virus infection in the areas with high rainfall (3). Nevertheless, it should be noted that there are also other factors that affect the epidemiology of Zika virus infection. The temperature is a good example. For the Indonesian case, Tosepu recently published another similar study confirming the relationship between temperature and Zika virus infection prevalence (4). Finally, the important fact is that Zika virus infection has multimodal transmission. The non-mosquito transmission is possible and this makes a significant distortion in the relationship between either rainfall or temperature and Zika virus infection prevalence.

## REFERENCES

- 1. Tosepu R. The Appropriate Rainfall to Development of Zika Virus: An Indonesian Case. Ethiop J Health Sci. 2017 Jan;27(1):104-105
- 2. Wiwanitkit V. An observation on correlation between rainfall and the prevalence of clinical cases of dengue in Thailand. J Vector Borne Dis. 2006 Jun;43(2):73-6.
- 3. Wiwanitkit S, Wiwanitkit V. Predicted pattern of Zika virus infection distribution with reference to rainfall in Thailand. Asian Pac J Trop Med. 2016 Jul;9(7):719-20. 4. Tosepu R. Temperature and development of Zika virus infection: An Indonesian case. Asian Pac J Trop Med. 2017 Mar;10(3):320.

<sup>\*</sup>Email: beuyjoob@hotmail.com