

**ORIGINAL ARTICLE****SOIL TRANSMITTED HELMINTHS AND ASSOCIATED FACTORS AMONG SCHOOLCHILDREN IN GOVERNMENT AND PRIVATE PRIMARY SCHOOL IN JIMMA TOWN, SOUTHWEST ETHIOPIA**Serkadis Debalke<sup>1</sup>, Amare Worku<sup>2</sup>, Nejat Jahur<sup>1</sup>, Zeleke Mekonnen<sup>1</sup>**ABSTRACT**

**BACKGROUND:** Soil transmitted helminth infections are among the most common human infections. They are distributed throughout the world with high prevalence rates in tropical and sub-tropical countries mainly because of lack of adequate sanitary facilities, inappropriate waste disposal systems, lack of safe water supply, and low socio-economic status.

**METHODS:** A comparative cross sectional study was conducted from December 2011 to June 2012 to determine and assess the prevalence of soil transmitted helminths and their associated factors among government and private primary school children. Stool samples were collected from 369 randomly selected children and examined microscopically for eggs of soil transmitted helminth following McMaster techniques. Soil samples were collected from different parts of the school compound and microscopic examination was performed for eggs of the helminths using sodium nitrate flotation technique.

**RESULTS:** The overall prevalence rate of soil transmitted helminth infections in private and government schools was 20.9% and 53.5% respectively. *T. trichiura* was the most common soil transmitted helminth in both schools while hookworm infections were identified in government school students only. Type of school and sex were significantly associated with soil transmitted helminth. Soil contamination rate of the school compounds was 11.25% with predominant parasites of *A. lumbricoides*.

**CONCLUSION:** Higher prevalence of soil transmitted helminth infection was found among government school students. Thus, more focus, on personal hygiene and sanitary facilities, should be given to children going to government schools.

**KEYWORDS:** Soil transmitted helminths, school children, government and private schools

DOI: <http://dx.doi.org/10.4314/ejhs.v23i3.6>

**INTRODUCTION**

Soil transmitted helminths (STH) have been identified as a serious public health problem, predominantly among poor communities in the developing world (1, 2). Over one billion people in the world are affected by STHs alone (3) particularly putting school age (5-15 years) children at risk (2, 4).

High incidence rate of STH infections occur in the Americas, China, East Asia and Sub-

Saharan Africa (2). Prevalence rate of 15.6% (Thailand (5)), 88.4% (Turkey (6)), 69.9 % Tuvalu (7), 53% (Guinea (8)), 40.2% (China (9)), 92.6% (India (10)), and 43.5% (Ethiopia (11)) were reported from the different parts of the world.

*Ascaris lumbricoides* (*A.lumbricoides*), Hookworm (*Ancylostoma duodenale* and *Necator americanus*) and *Trichuris trichiura* (*T.trichiura*) are among the most common STHs which have

<sup>1</sup>Department of Medical Laboratory Science and Pathology, Jimma University, Jimma, Ethiopia

<sup>2</sup>Addis Continental Institute of Public Health, Addis Ababa, Ethiopia

**Corresponding Author:** Serkadis Debalke, Email: [serkadis.debalke@ju.edu.et](mailto:serkadis.debalke@ju.edu.et)

been reported among several hundred millions of people worldwide (2, 12). The geographical distribution of STHs is influenced by various factors including external environmental conditions like soil (13), absence of sanitary facilities, unsafe waste disposal system, inadequacy and lack of safe water supply, types of toilet (4,11) and human factors including age, sex, socio-economic status and occupation (9,10).

A study by Soriano *et al.* (2001) suggested that all these factors should be assessed as a whole when evaluating the prevalence of intestinal parasites in a given population, taking into account the parasite-host-environment relationship (13). In addition, in developing prevention and control strategies and for the empirical treatment of STH, knowledge of the most likely causative agents, the possible risk factors and status of soil contamination are essential. However, to our knowledge, particularly in Jimma, no published research report is available on the prevalence of STH among children comparing private and government schools where the socio-economic status and other associated factors may vary. On top of this, the geographical difference in the status of contaminated source (eg. soil) was also not well studied. Therefore, this study was done to determine the prevalence of STH and assess the associated factors including status of soil contamination with STHs among government and private primary schools.

## MATERIALS AND METHODS

A comparative cross-sectional survey was conducted from December 2011 to June 2012 on randomly selected two private and two government primary schools in Jimma Town, South-west Ethiopia. All students, 5-15 years, with and without clinical symptoms of gastro intestinal tract (GIT) infection who had not taken anti-helminthic treatment in the last 3 months were included based on their willingness.

The sample size was based on a predicted prevalence of STH of 60% (14) among school children  $\pm$  5% precision and 95% confidence level. To investigate the associated factors, being in government school was assumed as an exposure (15) to detect odds of 4.3 at 95% confidence interval and 80% power. Accounting for a 10% non-response rate and a design effect of 2, the

final estimated sample size was 369. A multistage sampling with proportional to size sampling method was employed considering type of school, students school grade and class section. A total of 278 and 91 children were randomly selected from government and private school respectively.

Data on socio-demographic characteristics, variables related to type of latrine used and hand washing and shoe wearing habits of the children were collected. All the study subjects were supplied with labeled plastic containers with applicator sticks, and instructed to provide a recent stool sample (an interval of less than 4 hours). Then, the stool was processed and examined for STH using McMaster method as described by Charlier *et al.* (16). The laboratory results were recorded on laboratory formats. During sample collection, physical inspection was made on toilet facilities of the school compounds and fingernail status of the students.

Eighty soil samples were collected from different sites of the school's compound including the playgrounds, the area near the water tap, the classroom and the toilet surroundings. The samples were analyzed by sodium nitrate flotation technique as described by Mizgajska-Wiktor (17). Briefly, soil samples from the ground were collected from a depth of 3 centimeters and 40 grams of dried and sifted soil sample was suspended in 5% sodium hydroxide (NaOH) and left for 1 hour to separate eggs from the soil. The sample was centrifuged for 3 minutes at 1500 rotations per minute (rpm). The supernatant was discarded and the sediment was washed with distilled water. After the sediment was washed, it was re-suspended in saturated sodium nitrate (NaNO<sub>3</sub>) solution with specific gravity of 1.30 and centrifuged at 1500 rpm for 3 minutes. The tube was then filled up to the top with the flotation fluid, NaNO<sub>3</sub>. A cover slip was placed on top of the fluid for 30 minutes and it was lifted from the tube, placed on a clean glass slide and examined under the microscope for STH eggs.

Data was entered and analyzed using statistical package for social science (SPSS) version 16. Frequencies, percentages and measures of central tendency were calculated. Bivariate logistic regression of independent variables with STH infection was done to select subset of predictor variables. Multivariate logistic analysis was done to observe and identify

determinant factors associated with STH. All variables with P-value < 0.2 were included in the final model. A P-value of less than 0.05 was considered statistically significant. A total of 366 children were included for the logistic regression.

Ethical clearance was obtained from the College of Public Health and Medical Sciences Research Committee, and Research Project Office of Jimma University. The purpose of the study and the entire voluntary nature of the participation were clearly explained to all students or their parents before obtaining a written informed consent. Oral consent was obtained from the heads of the schools. Stool examination results were revealed to all participants and appropriate drug was given to students infected with STH.

## RESULTS

A total of 369 children responded to the questionnaire and samples were collected from all students. Three samples were discarded due to insufficiency of the amount of sample for examination. Out of 366 students, 75.1% were from government primary schools. The majority of the children were females with male to female ratio of 0.79:1. More than half (53.5%) of the students were within the age range of 5-10 years with a mean of  $10.27 \pm 2.14$ .

The overall prevalence rates of STH infections in private and government schools were 20.9% and 53.5% respectively. High prevalence rate of STH infections was identified among students with the age range of 11-15 years in both private (25%) and government (57.04%) schools. Males' infection rate was 25% among private schools while females' was 59.7% among government school students (Table 1).

Table 1: Distribution of STH infections by Socio demographic feature of the students of Selected Private and Government Primary schools, Jimma, Ethiopia, 2011/12.

Variables	Private School Students			Government school Students		
	Number Examined	Number Infected	Infected rate (%)	Number Examined	Number Infected	Infected rate (%)
<b>Age</b>						
5-10	63	12	19.1	133	66	49.62
11-15	28	7	25	142	81	57.04
<b>Sex</b>						
Male	40	10	25	121	55	45.5
Female	51	9	17.6	154	92	59.7
<b>Grade level</b>						
1-3	47	10	21.3	157	89	56.7
4-6	44	9	20.5	118	58	49.2
<b>Total</b>	91	19		275	147	

Half of the students (49.5%) were infected with some form of intestinal helminthes, out of these (45.4 %) were positive for STH with *T. trichiura*, *A. lumbricoides* and hookworm infections. *A. lumbricoides* and *T. trichiura* infections were identified from both private and government

schools while hookworm infections were only identified from government schools. *T. trichiura* was the most common STH in both private (59.1%) and government (46.4%) schools (Table 2).

Table 2: Distribution of STH identified from Private and Government Primary school students, Jimma, Ethiopia, 2011/12.

STH Type	Private Schools Infected No (%)	Government Schools Infected No (%)	Both schools Infected No (%)
<i>T. trichiura</i>	13 (59.1)	98 (46.4)	111 (47.6)
<i>A. lumbricoides</i>	9 (40.9)	83 (39.3)	92 (39.5)
Hookworm	0 (0)	30 (14.3)	30 (12.9)
<b>Total</b>	<b>22(100)</b>	<b>211(100)</b>	<b>233(100)</b>

Single, double and triple infections were found with infection rates of 64.5%, 30.7% and 4.8% respectively. Double infections were the most common incidents among students of government

primary schools (32.7%) compared to those in students of private schools (15.79%). Triple infections (5.4%) were only found in students of government schools (Table 3).

Table 3: Types of STH infections in students from private and government primary schools, Jimma, Ethiopia, 2011/12.

<i>Type of Infection</i>	<i>Private Schools No (%)</i>	<i>Government Schools No (%)</i>	<i>Total No(%)</i>
Single Infection	16 (84.2%)	91 (61.9%)	107 (64.5%)
Double Infection	3 (15.8%)	48 (32.7%)	51 (30.7%)
Triple Infection	0 (0%)	8 (5.4%)	8 (4.8%)
<b>Total</b>	<b>19 (100%)</b>	<b>147 (100%)</b>	<b>166(100%)</b>

Of the total multiple infections, (62.7%) of the students had double infection predominantly with

a combination of *A. lumbricoides* plus *T.trichiura* (Table 4).

Table 4: Multiple infections in selected Private and Government Primary schools, Jimma, Ethiopia, 2011/12.

<i>Parasites Combination</i>	<i>Infection No (%)</i>
<i>A. lumbricoides</i> + <i>T.trichiura</i>	37 (62.7%)
<i>A. lumbricoides</i> + Hookworm	7 (11.9%)
<i>T.trichiura</i> + Hookworm	7 (11.9%)
<i>A. lumbricoides</i> + Hookworm + <i>T.trichiura</i>	8 (13.5%)
<b>Total</b>	<b>59 (100%)</b>

In all schools, functional pit latrine facilities were available. However, the ratio of the number of students sharing the toilet to the number of toilet facilities varied from school to school. In the government schools, the ratio of the number of toilet facilities to the number of students was 1:127 while it was 1:39 in private schools. Among government school students, 62.9% used open field or wood type latrine at home and 37.1% had pit latrine. None of them had water system latrine at home. Forty three (47.3%) of the private school students had pit latrine at home and 31.9% use water system latrine.

A total of 80 (40 from private and the rest 40 from government schools) soil samples were collected from different sites of the school compound including playgrounds, water tap, class

room and toilet. Soil contamination rate of the school compounds with STH was 11.25%. The majority of STHs identified were from government schools (77.8%) and the rest (22.2%) were from private school compounds. Egg of *A. lumbricoides* and *T.trichiura* parasites were found with the infection rate of 77.8% and 22.2% respectively.

Binary logistic analysis showed age, school type, type of latrine at home and wearing shoe were independently associated with STH. The multivariate logistic regression model revealed male sex (AOR, 0.637, 95%CI, 0.40 - 0.99) and enrollment in government school (AOR, 2.63, 95% CI, 1.30 - 5.30) were significantly associated (P<0.05) with STH (Table 5).

Table 5: Bivariate and multivariate logistic regression analysis of factors associated with STH infections among the students of Selected Private and Government Primary schools, Jimma, Ethiopia, 2011/12.

<i>Variable</i>	<i>STH (%)</i>	<i>NoSTH(%)</i>	<i>OR (95% CI)</i>	<i>P-value</i>	<i>AOR (95% CI)</i>	<i>P-value</i>
<b>Age</b>						
5-10	78(39.8)	118(60.2)	1		1	
11-15	88(51.8)	82(48.2)	1.624(1.072,2.459)	0.022**	1.472(0.935,2.317)	0.095
<b>Sex</b>						
Female	101(49.3)	104(50.7)	1		1	
Male	65(40.4)	96(59.6)	0.697(0.459,1.058)	0.090	0.637(0.408,0.995)	0.047**
<b>School Type</b>						
Private	19 (20.9)	72(79.1)	1		1	
Government	147(53.5)	128(46.5)	4.352(2.490,7.606)	0.000**	2.626(1.300,5.304)	0.007**
<b>Home Latrine</b>						
Cement (Pit)	56(38.6)	89(61.4)	1		1	
Wood	105(54.7)	87(45.3)	1.918(1.237,2.975)	0.000**	1.484(0.909,2.424)	0.115
Sink (Flush)	5(17.2)	24(82.8)	0.331(0.119,0.918)	0.004**	0.768(0.242,2.438)	0.655
<b>Water source</b>						
Pipe	150(45.6)	179(54.4)	1		1	
Others	16(43.2)	21(56.8)	0.909(0.458,1.805)	0.786	-	-
<b>Finger nails</b>						
Trimmed	78(41.7)	109(58.3)	1		1	
Untrimmed	88(49.2)	91(50.8)	1.351(0.894,2.042)	0.153	0.960(0.608,1.513)	0.859
<b>Hand wash BE</b>						
Always	73(41.5)	103(58.5)	1		1	
Sometimes	93(48.9)	97(51.1)	1.353(0.895,2.045)	0.152	1.149(0.725,1.823)	0.554
<b>Hand wash AD</b>						
Always	50(40.3)	74(59.7)	1		1	
Sometimes	116(47.9)	126(52.1)	1.363(0.879,2.113)	0.167	0.912(0.545,1.526)	0.726
<b>Wear shoe</b>						
Always	84(37.2)	142(62.8)	1		1	
Sometimes	82(58.6)	58(41.4)	2.390(1.553,3.678)	0.000**	1.638(0.986,2.721)	0.057

Hand wash BE= Hand washing habit before eating, Hand wash AD= Hand washing habit after defecation  
Sink = Water system/flush

## DISCUSSION

This comparative cross sectional study revealed that an overall high prevalence rate of STH infections was found in government school students (53.5%) as compared to private school students (20.9%). Even though there was a slight difference in the infection rate, this finding was in line with a comparative study conducted in Nigeria where the rate of helminthiasis was higher among government school students (46.6%) than among private school students (16.6%) (15). This might be due to variation between the schools with regard to factors that contribute for the transmission of STH including socioeconomic status, shoe wearing habit, safe water supply at home and school, personal hygiene, environmental sanitation of the school compound, waste disposal

system within the school, type of toilet accessible at home and availability of clean and sufficient number of toilet facilities at school, which have been observed in the present study.

In the present study, *T. trichiura* was the most common STH parasite in both private (59.1%) and government schools (46.4%). On the contrary, a study conducted on helminthiasis among pupils of private and government schools in Nigeria found out that the predominant parasites were *A. lumbricoides* (6.0%) and hookworm (26.0%) respectively (15). The finding was also incomparable with other non-comparative cross sectional studies conducted on STH among school age children in Sri Lanka (18), South-Easter Nigeria (19), Pacific Island (7), Uganda (20), Turkey (6) and other parts of Ethiopia including North West of the country (21),

Adwa (22) and Chilga (23). This might be due to differences in environmental factors like climate, topography (24), surface temperature, altitude, soil type and rainfall which have a great impact on the distribution of STH (25). In addition, recently at the study site and other parts of the world, *T. trichiura* has low drug efficacy to commonly used anthelmintic drugs unlike *A. lumbricoides* and hookworm (26). However, this study showed a similar finding to one study done on STH among primary school children in Thailand (27) where *T. trichiura* was the predominant parasite.

Regarding the associated factors, multivariate analysis showed that sex and school type were the best predictors of STH among primary school children. Accordingly, government school students were 2.6 times more likely to have STH as compared to private school students. Soil sample results from the government school compound support the finding as well. This result is similar with a study carried out in Nigeria where pupils in public primary schools were 2.5 times more infected than private school students (15).

Sex was one of the factors for STH where male students were 0.6 times less likely to have STH as compared to female students. Though a report from Nepal (28) suggest that females are at higher risk for STH, other studies have presented mixed results. A study in Nigeria (15) and Ethiopia, Chilga (23), show that males had an overall higher prevalence rate. This could be explained by personal hygiene related to trimming fingernails where in the present study the majority of the participants who had untrimmed nails were female students.

Though there was no significant association between STH infection and latrine type, an overall high prevalence rate was observed among students who used open space or wood type latrine (54.7%) as compared to pit (38.6%) and water system (17.2%) users. This finding was in agreement with a finding in Nigeria where pupils who used open spaces and bush had higher prevalence (77.7%) than those who used the pit latrine (33.79%) and water system users (11.74%) (15).

In the present study, the soil contamination rate of the school compounds was found to be lower (11.25%) than that of a previous study conducted in Nepal, Kathmandu Valley (28.5%) (28). This might be due to difference in socioeconomic status, health information, waste

disposal system, sample collection season, soil type, as well as climatic and topographic factors of the countries. However, a similar finding was observed in Nepal (28) where the predominant parasite isolated from the soil samples was *A. lumbricoides*.

The prevalence of STH was significantly higher among government school students than their private counterparts. Enrollment in government school and male sex were associated with STH among primary school children. Therefore, improving availability of clean and sufficient number of toilet facilities, safe and adequate water supply, environmental sanitation of the school compound and waste disposal system at school especially within the government school, and health education on personal hygiene are highly recommended. Given the current level of the problem among school children and the immediate environmental context that these children live in, assessment and appropriate intervention should be an integral part of control and prevention strategies and programs. In addition, further study with large sample size is recommended to investigate risk factors among both government and private schools independently.

## ACKNOWLEDGEMENTS

We sincerely acknowledge all of the study participants, teachers, directors of the selected primary schools for their participation and unreserved support. We are also grateful to the contributions of Jimma University, Medical Laboratory Science and Pathology Department, technicians working in the department as well as in the hospital. We are also indebted to Jimma University for providing financial support for this study.

## REFERENCES

1. Mordi RM, Okaka CE. Prevalence of Intestinal Parasites in Edo State. *Int J Health Res.* 2009; 2(3): 253-257.
2. WHO Soil-transmitted helminthes. (Updated) 2010. Available at [http://www.who.int/intestinal\\_worms/en/](http://www.who.int/intestinal_worms/en/). Accessed on January 2011.

3. Crompton DW. How much human helminthiasis is there in the world? *J Parasitol.* 1999; 85: 397-403
4. WHO, Prevention and Control of Schistosomiasis and Soil transmitted Helminthiasis. Report of a WHO Expert Committee. 2002; Technical Report Series, No 912. pp. 63 Geneva, World Health Organization.
5. Anantaphruti MT, Waikagul J, Maipanich W, Nuamtanong S, Pubampen S. Soil-transmitted helminthiasis and health behaviors among schoolchildren and community members in a west-central border area of Thailand. *Southeast Asia J Trop Med Public Health.* 2004; 35(2): 260-6.
6. Ulukanligil M, Seyrek A, Aslan G, Ozbilge H, Atay S. Environmental Pollution with soil-transmitted Helminths in Sanliurfa, Turkey. *Mem Inst Oswaldo Cruz.* 2001; 96(7): 903-909.
7. Speare R, Latasi FF, Nelesone T. Prevalence of soil transmitted nematodes on Nukufetau, a remote Pacific island in Tuvalu. *BMC Infectious Disease.* 2006; 6: 110.
8. Glickman LT, Camara Ao, Glickman NW, McCabe GP. Nematode intestinal parasites of children in rural Guinea, Africa: prevalence and relationship to geophagia. *International Journal of Epidemiology.* 1999; 28: 169-174.
9. Mofid LS, Bickle Q, Jiang J-Y, Du Z-W, Patrick E. Soil-Transmitted Helminthiasis in rural South-West China: Prevalence, intensity and risk factor analysis. *Southeast Asia J Trop Med Public Health.* 2011;42(3): 513-526.
10. Naish S, McCarthy J, Williams GM. Prevalence, intensity and risk factors for soil transmitted helminth infection in a South Indian fishing village. *Acta Tropica.* 2004; 91(2): 177-187.
11. Belyhun Y, Medhin G, Amberbir A. Prevalence and risk factors for soil-transmitted helminth infection in mothers and their infants in Butajira, Ethiopia: a population based study. *BMC Public Health.* 2010; 10:21.
12. Crompton DWT, Savioli L. Helminthiasis for Public Health. USA: CRC Press. 2006.
13. Soriano SV, Barbieri LM, Pierangeli NB. Intestinal parasites and the environment: Frequency of intestinal parasites in children of Neuquen, Patagonia, Argentina. *Rev Latinoam Microbiol.* 2001; 43(2):96-101.
14. Mekonnen Z, Levecke B, Boulet G., Bogers J-P., Vercruyse J: Efficacy of different albendazole and mebendazole regimens against high-intensity *Trichuris trichiura* infections in school children, Jimma, Ethiopia.(under review)
15. Ogwurike BA. Ajayi JA, Ajayi OO. A Comparative Study of Helminthiasis among Pupils of Private and Public Primary Schools in Jos North local Government Area of Plateau State, Nigeria. *Nigerian Annals of Natural Sciences.* 2010; 10 (1): 28 – 47.
16. Charlier J, Geiger SM, Hoa NTV. A Comparison of the Sensitivity and Fecal Egg Counts of the McMaster Egg Counting and Kato-Katz Thick Smear Methods for Soil-Transmitted Helminths. *PLoS Negl Trop Dis.* 2011; 5(6): e1201.
17. Mizgajska-Wiktor H. Recommended method for recovery of *Toxocara* and other geohelminth eggs from soil. *Wiad Parazytol.* 2005; 51(1): 21-22.
18. Gunawardena K, Kumarendran B, Ebenezer R, Gunasingha MS, Pathmeswaran A, de Silva N. Soil-Transmitted Helminth Infections among Plantation Sector Schoolchildren in Sri Lanka: Prevalence after Ten Years of Preventive Chemotherapy. *PLoS Negl Trop Dis.* 2011; 5(9): e1341.
19. Uneke CJ, Eze KO, Oyibo PG, Azu NC, Ali E. Soil-Transmitted Helminth Infection In School Children In South-Easter Nigeria: The Public Health Implication. *Internet Journal of Third world Medicine.* 2007; 4 (1): DOI: 10.5580/3f7
20. Kabatereine NB, Tukahebula EM, Kazibwe F. Soil-Transmitted helminthiasis in Uganda: epidemiology and cost of control. *Trop Med Int Health.* 2005;10(11):1187-1189.
21. Alemu A, Atnafu A, Addis Z. Soil transmitted helminths and *schistosoma mansoni* infections among school children in zarima town, northwest Ethiopia. *BMC Infectious Diseases.* 2011; 11:189.
22. Legesse L, Erko B, Hailu A. Current status of intestinal Schistosomiasis and soil transmitted helminthiasis among primary school children in Adwa Town, Northern Ethiopia. *Ethiop. J. Health Dev.* 2010; 24(3):191-197.

23. Jemaneh L. Soil-Transmitted Helminth Infections and *Schistosoma mansoni* in School Children from Chilga District, Northwest Ethiopia. *Ethiop. J Health sci.* 2001; 11(2): 80-87.
24. Brooker S, Singhasivanon P, Waikaqul J. Mapping soil-transmitted helminths in Southeast Asia and implications for parasite control. *Southeast Asian J. Trop. Med. Public Health.* 2003;34 (1):24-36.
25. Appleton C.C, Gouws E. The distribution of common intestinal nematodes along an altitudinal transect in KwaZulu-Natal, South Africa. *Ann. Trop. Med. Parasitol.* 1996; 90:181-188.
26. Vercruyssen J, Behnke JM, Albonico M. Assessment of the Anthelmintic Efficacy of Albendazole in School Children in Seven Countries Where Soil-Transmitted Helminths Are Endemic. *PLoS Negl Trop Dis.* 2011; 5(3): e948.
27. Anantaphruti MT, Nuamtanong S, Muennoo C, Sanguankiat S, Pubampen S. *Strongyloides Stercoralis* Infection and Chronological Changes of Other Soil-Transmitted Helminthiases in an Endemic Area of Southern Thailand. *South East Asian J Trop Med Public Health.* 2000;31(2): 378-382.
28. Shrestha A, Rai SK, Basnyat SR, Rai CK, Shakya B. Soil transmitted helminthiasis in Kathmandu, Nepal. *Nepal Med coll J.* 2007; 9(3):166-9.