FOOD ANIMAL SUPPLY AND CONSUMPTION PATTERN IN AKWA IBOM STATE,

NIGERIA

O. L. OBASI, E. O. EKANEM, O. J. IFUT and M. E. OGBEBOR

(Received 26 February 2001; Revision accepted 6 June, 2001)

ABSTRACT

A three-year retrospective study (1994-1996) of food animal supply and consumption in Akwa lbom State was carried out, based on records kept at the Veterinary Services Department of the Ministry of Agriculture. This was complemented with visits to appraise operational areas and interviews of the operators.

The animals that provided meat included cattle (47%); Pig (37.5%), Goat (9.2%); Dog (4.8%) and sheep (1.5%). Most of these animals originated from the Northern part of Nigeria, and were transported by road. Poultry, fish and other sea foods abound but were not covered in this report. The supply and slaughter of the various animal species peaked during the periods of Christian and Traditional festivities (December, March, April and May). The calculated animal protein intake per person per day from these sources was 0.57g, equivalent to 3.45g of meat. The calculated monetary value of the meat trade was N1.6 billion, representing 26.23% of the state's budget for the period.

In conclusion, livestock and dog do not appear to provide adequate meat for the inhabitants of the state. There is the need to adequately coordinate food animal production and procurement and also ensure proper abattoir operations.

KEY WORDS - FOOD ANIMAL, MEAT, SUPPLY, CONSUMPTION.

INTRODUCTION

Agriculture is the major employer of labour in Akwa Ibom State. Crop farming, fishing and livestock farming are the important areas. Animal agriculture is the least developed (Abasiattai, 1987). There is therefore a great deal of dependence on outside sources in order to meet the animal protein need of the inhabitants.

The quantity and quality of meat consumed by a people, and the animal species used for food are governed by several factors, among which are social status and culture. Foods of animal origin are essential in man's diet

and are an excellent source of essential amino acids, vitamins and minerals (CAST, 1997).

The food and Agriculture organization's (FAO, 1980) estimate for Nigeria is that cattle is the major meat source (36%), followed closely by poultry (34%), small ruminants (24%) and pigs (6%).

The various ecological zones of Nigeria are not equally endowed with livestock resources. An interdependence is thus created, whereby high livestock producing areas supply areas with a net deficit (Agyemang, 1998). This interdependence is well pronounced because of minimal obstacles and restriction on livestock movement and livestock trade.

This study was undertaken to evaluate

O. L. OBASI, Department of Animal Science, Faculty of Agriculture, University of Uyo, Uyo, Nigeria

E. O. EKANEM, Department of Animal Science, Faculty of Agriculture, University of Uyo, Uyo, Nigeria O. J. IFUT, Department of Animal Science, Faculty of Agriculture, University of Uyo, Uyo, Nigeria

M. E. OGBEBOR, Department of Animal Science, Faculty of Agriculture, University of Uyo, Uyo, Nigeria

the structure of food animal supply and consumption in Akwa Ibom State. It was, in particular designed to determine the volume and the value of food animal and the meat trade.

METHODOLOGY

The study was conducted between December 1997 and August 1998. It was a 3 year retrospective study of food animal supply and consumption in Akwa Ibom State (January 1994 to December, 1996).

Akwa Ibom State is located in the humid forest zone of Nigeria. It is made up of 31 Local Government Areas; with Uyo urban doubling as the State Capital and Uyo Local Government Headquarter. Uyo is also the only metropolitan city of the state. The human population of Akwa Ibom State is 2, 359, 736 (NPC/CBN, 1992).

Information on number and type of livestock transported into the state and the daily slaughter figure were collected from the livestock control post and abattoir units record books of the Department of Veterinary Services,

Ministry of Agriculture, Uyo. Some abattoirs, livestock sales out posts and meat markets were visited and an on the spot assessment of activities recorded. Operators of these areas as well as the butchers operating outside the abattoirs were also interviewed and evaluated. Data collected were analyzed, using descriptive statistics, while Scheffe's method (Morad, 1990) was used to separate means.

RESULTS AND DISCUSSION

The animals used for food were cattle, pig, goat, dog and sheep. These animals mainly originated from the Northern part of Nigeria, principally Kano, Kaduna, Plateau and Gongola (now Taraba and Adamawa) states, and arrive Akwa Ibom State by road. (Poultry, fish and other seafoods and snails abound but were not covered by this study. The meat from these animal species are widely consumed in Nigeria, except the dog which has not recorded a wide and open acceptance, and the pig which is limited by the Islamic faith. Ekumankama and Ekumankama (1998) reported 10% acceptance

TABLE 1: TOTAL NUMBER OF LIVESTOCK TRANSPORT INTO AKWA IBOM STATE (1994 – 1996)

Month	Cattle No (%)	Goat No(%)	Sheep No (%)	Pig (%)	Dog No (%)
January	2605 (8.8)	2888(2.9)	456(10.0)	2525(5.3)	1622(7.2)
February	1995 (6.7)	2496(2.5)	354(7.8)	2310(4.9)	1260(5.6)
March	2926 (9.7)**	4905(5.0)**	546(12.0)**	2342(4.9)	1810(8.1)
April	3484 (11.6)	3810(3.9)	501(11.0)	4805(10.1)	2603(11.6)**
May	2737 (9.1)	2821(2.9)	345(7.6)	5654(11.9)**	1686(7.5)
June	2541 (8.5)	2586(2.6)	256(5.6)	2354(4.9)	1240(5.5)
July	1846 (6.2	2564(2.6)	289(6.3)	2491(5.2)	1856(8.3)
August	1936 (6.5)	2460(2.5)	184(4.0)	2404(5.1)	950(4.2)
September	1807 (6.0)	2650(2.7)	345(7.6)	2642(5.6)	1644(7.3)
October	2571 (8.6)	3121(3.2)	362(8.0)	2528(5.3)	1950(8.6)
November	2429 (8.1)	3560(3.6)	417(9.2)	3129(6.6)	2526(11.3)
December	3052 (10.2)*	64705(65.7)*	497(10.9)*	14448(30)*	3330(14.8)*
TOTAL	29929 (100)	98566(100)	4552(100)	47,628(100)	22,457(100)

Highest value; ** 2nd highest value

Source: Adapted from Livestock Control Post Records, Akwa Ibom State (1994 1996)

TABLE II:	TOTAL NUMBER OF LIVESTOCK SLAUGHTERED IN AKWA IBOM STATE (1994 – 1996)
-----------	--

Month	Cattle/No (%)	Goat No (%)	Sheep No (%)	Pig No (%)	Dog No (%)
January	2247 (7.3)	4112(8.3)	412(7.7)	2000(5.2)	1575(3.6)
February	2377 (7.8)	3677(7.5)	480(8.9)	2762(7.2)	3141(7.2)
March	22311 (7.5)	5507(11.2)**	542(9.8)	2815(7.3)	4860(11.1)
April	3857 (12.6)**	3055(6.2)	673(12.6)**	2832(7.4)	6676(15.2)
May	2152 (7.0)	3109(6.3)	415(7.8)	2735(7.1)	7323(16.7)**
June	2303 (7.5)	3050(6.2)	332(6.2)	3106(8.1)	1764(4.0)
July	1937 (6.3)	2829(5.7)	309(5.8)	2259(5.9)	(577(3.6)
August	1772 (5.8)	2932(5.9)	359(6.7)	1901(4.9)	1865(4.3)
September	1697 (5.5)	2766(5.6)	320(5.9)	3236(8.4)**	1682(3.8)
October	818 (2.7)	2515(5.1)	270(5.0)	2505(6.5)	1741(4.0)
November	1943 (6.4)	3186(6.5)	324(6.1)	2487(6.5)	1524(3.5)
December	7200 (23.5)*	12,581(25.5)*	938(17.5)*	9815(25.5)*	10,094(23.0)*
TOTAL.	30,614(100)	49,319(100)	.356(100)	38,453(100)	43,821(100)

Highest value; ** 2nd highest value

Source: Adapted from abattoir records, Dept. of Vet, Services, Akwa Ibom State (1994 - 1996)

of dog meat in Abia State. Okpuduedu (1994) and Odo et al., (1998) also reported dog meat consumption in Akwa Ibom State and Enugu State respectively.

The total number of animals transported into the state during the period January 1994 to December, 1996 is shown in Table 1. More an mals were transported in December followed by March (cattle, goat, sheep, dog) and May (pigs). The slaughter records also showed consistent highest values in December for all the animal species, followed by March (goat); April (cattle, sheep); May (dog) and September (pig) (Table II). The high supply and slaughter values for December and March; April and May correlates with the period of high demand. These periods fall within the traditional Christian festivals, when also the population of the state is swollen by returnees, coming home to celebrate the season with their kiths and kins. Most'cultural activities and marriage ceremonies are also scheduled within these periods.

In terms of the quantity of meat derived from the slaughtered animals per annum; cattle vielded 991 893 6 kg; pig 807,513 kg; goat

182,480.3 kg; Dog 102,249 kg and sheep 25,228.8 kg representing 47%; 37.5%; 9.2%; 4.8% and 1.5% respectively of the total quantity of the meat derived per annum (Table These values were based on a₂ calculated carcass yield of 54% and 48% respectively for cattle and sheep (Williamson and Payne, 1978); 55.5% for goat and 70% for pig (Thornton, 1974). There was no referenced carcass yield for dog; and being a monogastric had to be approximated to the same value as for Cattle provided the bulk of meat the pig. consumed in the state, while sheep was the least (Table III). In a study of meat supply in Enugu State (1999-1995), Odo et al; (1998) reported an average contribution of 54% for cattle; 21% for small ruminants; 6.44% for pigs and 0.18% for dog. The significant contribution to meat supply by the pig and the dog in our study is note worthy. Meat of both species are more commonly vended. Pork in particular was very cheap (N180/kg); Mutton and goat meat (N200/kg) and Poultry meat (N250 per table weight). The relatively low price of pork coupled with the absence of religious and

cultural taboos might, in a great part explain increased contribution to consumable meat. If meat derived from food animals is assumed to contain 16% protein (Obioha, 1976); 2109, 364.7kg (mean quantity/annum) of meat shall therefore yield 348,041.1kg of protein. adjustment of 40% to cover non reported and non recorded slaughter (Okpuduedu, brings the total value to 487,263,1kg protein For a population of 2,359,736 per annum. persons (NPC/CBN, 1992), this value translates to 0.57g protein per person per day (equivalent to 3.45g meat/caput/day). An adult Nigerian was estimated to require 65g protein per day, out of which 35g should be from animal sources (Olubajo, 1976; FAO, 1980). Ademosun (1976) projected a daily per capita meat consumption of 5.7g equivalent to 0.85g protein for an adult Figerian in 1980. Estimates on the animal protein intake in Nigeria vary between 7g and 8.27g (Obioha, 1970). On the lower margin, the difference of 6.43g protein (7g less 0.57g) should be supplied by the other animal protein sources not covered in this study, for the inhabitants of Akwa Ibom State to meet the national average. In this regard, fish and other seafocds, poultry, rabbit and snail should play a significant role. The various rivers of the state are home to a variety of fishes and other seafoods. Fishing also constitutes the major occupation of the riverine people. Game meat is derived from the forested areas, wild Elephants constitute a danger to natives in Ikot Abasi (Okoh, 1985) and are often hunted for food. A significant contribution is, however, expected from these sources in order to meet the FAO (1980) minimum requirement of 35g animal protein/caput/day. A study in this respect would be helpful to avert grave implication of an endemic protein deficiency.

Annual distribution of livestock supply and slaughter values is shown in Table IV. Cattle and sheep showed no significant difference (P>0.05) in the number transported into the state between the years. The value for goat differed significantly P<0.05) across the years, while the dog showed a significant difference (P<0.05) in 1994. The study also revealed that the number of cattle transported into the state in 1994 was higher than the number slaughtered, while the reverse was the case in 1995 and 1996. The difference was also negative for goat and sheep in 1994 and 1995, while 1996 showed a positive value for

TABLE III: QUANTITY OF MEAT DERIVED FROM THE RESPECTIVE FOOD ANIMALS (1994 – 1996)

	PARAMETER	ANIMAL SPECIES					
		CATTLE	GOAT	SHEEP	PIG	DOG	TOTAL
s	NUMBER OF ANIMAL	30614	49319	5256	38453	43821	1.67463
2.	TOTAL LIVE WEIGHT (kg)	5,510,520 (180 kg)	986380 (20°kg)	157,690 (30°kg)	3,460,770 (90°kg)	438,210 (10°kg)	10,553,5
3.	QUANTITY OF MEAT (kg)	2,975,680.8 (54**%)	547,440.9 (55.5**%)	75,686.4 (48 %)	24,225.39 (70 %)	306,747 (70°kg)	6.328,094
4.	MEAN QUANTITY						
	(per annum) (kg)	991,893.6	182,480.3	25,228.8	807,513	102,249	2,109,364
5.	MEAN QUANITY PROTEIN						
	(per annum) (Assume meat contains 16.5% protein)	163,662.4	30,109.2	4,162.8	133,239.6	16,871.1	348,045,1
6.	% OF ALL ANIMALS	47	9.2	1.5	37.5	4.8	100 "

^{+ =} Average live weight per animal.

^{++ =} Carcass yield.

		1996)			3 1
				and were the second second and the second	
10807	12652	1486	11047	10034	
9389	16671	1488	<u>2361</u>	15988	
+1418	-4015	-2	-1314	-5954	
9646	14357	1350	13913	6603	
10500	18473	2130	11508	19232	
-854	-4 16	-780	+2405	-12629	
9476	71551	1716	22668	5820	
10725	14175	1638	14584	100,8	
-1249	+57,378 +78	+8084	-2781		
	10807 9389 +1418 9646 10500 -854 9476 10725	10807 12652 9389 16671 +1418 -4015 9646 14357 10500 18473 -854 -4116 9476 71551 10725 14175	10807 12652 1486 9389 16671 1488 +1418 -4015 -2 9646 14357 1350 10500 18473 2130 -854 -4116 -780 9476 71551 1716 10725 14175 1638	10807 12652 1486 11047 9389 16671 1488 2361 +1418 -4015 -2 -1314 9646 14357 1350 13913 10500 18473 2130 11508 -854 -4116 -780 +2405 9476 71551 1716 22668 10725 14175 1638 14584	10807 12652 1486 11047 10034 9389 16671 1488 2361 15988 +1418 -4015 -2 -1314 -5954 9646 14357 1350 13913 6603 10500 18473 2130 11508 19232 -854 -4116 -780 +2405 -12629 9476 71551 1716 22668 5820 10725 14175 1638 14584 8,601

both. The value for pig was negative for 1994 and positive for 1995 and 1996. The dog showed consistent negative values. The difference between the number of animals transported into the state and the number slaughtered is not unexpected. Where the supply is greater than slaughter, it could mean that some of the animals were withheld across the period. Similarly where slaughter is greater than supply the difference might be accounted or by the contribution from the locally produced tock, and withdrawals from the holding pens.

The calculated monetary value of the rneat trade for the 3-year period was \$\\1,594,679,713\$ (or an average value of \$\\531,559,904.3\$ per annum). This value was calculated based on the quantity of meat derived from all the animal species (6,328,094,1kg; Table III), adjusted for 40% non reported and non documented slaughter (8,8,859,331.74 kg), and sold for an average price of \$\\180/kg\$. This value represents 26.23% of the states total budget for the period.

CONCLUSION

Food animal production is low in Akwa Ibom State. there is therefore a great deal of dependence on supplies from other parts of Nigeria, principally the Northern state. the total consumable meat and therefore, the quantity of animal protein per caput per day was grossly below the FAO recommended level and the national average. The quantity however was augmented by poultry and other non-livestock meat. The meat trade has an immense potential to enhance the development of the state employment through and generation. At present there is need to adequately coordinate food animal procurement and ensure proper abattoir operations. This would plug loopholes in revenue losses and that accurate data on livestock purchases, slaughter and distribution are kept.

REFERENCES

- Abasiattai, M. B. 1987. The Land the people and culture of Akwa Ibom State and Cross River State. Wusen Press Calabar, pp. 133 143.
- Ademosun, A. 1976. The Livestock subsector in the Third National Development Plan: Nigeria Journal of Animal Production 1(3): 18 – 27.
- Agyemang, K. 1998. Interdependence of Ecological Zones in Sustainable Animal Production in West Africa. In: Animal Agriculture in West Africa: The Sustainability Question. Oduguwa, O. O., Fanimo, O. A. Osinowo, O. A. (eds). Pp. 121 122.
- CAST, 1997. Council for Agricultural Science and Technology. Contributions of Animal Products to Healthful Diets. CAST Task Force Report, 1997.
- Ekumankama Ijeoma, O. and Ekumankama Oji, O. 1998. Household Production of Livestock and Consumption of Animal Protein in Umudike. In: Sustainable Animal Production in West Africa: the Sustainability Question. Oduguwa, O. O.; Fanimo, O. A.; Osinowo, O. A. (eds.) pp. 133 134.
- F.A.O. 1980. Production Year Book Vol. 34. Pp. 159.
- Morad, M. 1990. Design and Analysis of Small Agricultural Experiments. Agricultural Teacher Training Institute, Wageningen. The Netherlands, 42 pp.
- NPC/CBN (1992). National Population Commission and Central Bank of Nigeria Joint Statistical Bulletin. 31st December, 1992.

- Obioha, F. C. (1976). The Contribution of Poultry, Swine and Rabbit Production to the Short-term Solution of Meat Scarcity in Nigeria. Nigeria Journal of Animal Prod. 1(3): 45 48.
- Odo, B. J. Alaku, S. O. Omeje, S. I and Akpa, M. O. 1998.

 The Contribution of Small Ruminants to Meat Supply in Enugu State. in: Animal Agriculture in West Africa: The Sustainability Question. Oduguwa, O. O. Osinowo, O. A. (eds.) pp. 121-122.
- Okoh I. K. 1985. Preservation of Meat in Developing Countries. World Animal Review No. 10, pp. 8 - 14.
- Okpuduedu, F. E. 1994. Livestock Movement and Transportation in Akwa Ibom State of Nigeria: An Economic Appraisal. B. Sc. Thesis, University of Uyo, Uyo.
- Olubajo, F. O. 1976, Production of Meat from Ruminants. Nigerian Journal of Animal Prod. 1(3) 18 27.
- Thornton, H. and Gracey, J. F. 1974. Textbook of Meat Hygiene. 6th ed. Bailliere Tindall, New York, p. 18.
- Wilson, P. N. 1958. The Effect of Plane of Nutrition on the Growth and Development of the African Dwarf Goat
- Part II. Age Changes in the Carcass Composition of the Female Kids. J. Agric Science 51 (4 21).
- Williamson, G. and Payne, W. J. A. 1975. An introduction to Animal Husbandry in the Tropics. 3rd ed. Longman Group Ltd. London pp. 390 and 443.