



EVALUATING HEAVY METALS AND POLYCYCLIC AROMATIC HYDROCARBON CONCENTRATIONS IN COMMON NIGERIAN FOOD ADDITIVES

*JOHN, GODSON NNAMDI AND ECHEFU, MADUABUCHI FRANCIS

*Correspondence email: godson.john@uniben.edu

*ORCID NO: <https://orcid.org/0009-0007-6952-0820>

(Received 26 January 2026; Revision Accepted 17 April 2026)

ABSTRACT

This study analysed heavy metals and polycyclic aromatic hydrocarbon (PAH) contamination in popular spices and food colorants from a market in Port Harcourt, Nigeria, highlighting safety concerns regarding consumption. Utilizing Nitric acid-Hydrochloric acid digestion, AAS, and GC-MS, the analysis revealed that Lead (Pb) - 3.4 ppm, 0.32 ppm, 1.22 ppm, 0.46 ppm and 3.01 ppm in samples A, B, C, D and E respectively, Copper (Cu) concentrations in these five samples were 8.12 ppm, 3.23 ppm, 5.43 ppm, 1.35 ppm and 8.00 ppm, Iron (Fe) concentrations are 55.21 ppm, 31.32 ppm, 38.11 ppm, 22.21, Zinc, and Chromium levels, along with PAHs like naphthalene and chrysene, often exceed WHO standards. The findings indicate that the tested spices contain hazardous contaminant levels, suggesting a need for stricter quality control of local food additives to protect consumer health.

KEYWORDS: Heavy Metals, Polycyclic Aromatic Hydrocarbons (PAHs), Food Additives, Food Safety, Atomic Absorption Spectroscopy, GC-MS.

INTRODUCTION

Spices are substances from organic or inorganic origins that add flavor to dishes. They can be derived from different parts of plants including roots, rhizomes, stems, leaves, bark, flowers, fruits or seeds. (Abdullahi, 2008). Today, spices are used around the world including Nigeria. Despite the flavoring effects of spices, some however are made with rather harmful and toxic chemicals and are sold around the nation with lots of consumers on daily basis (Singh et al, 2015). This study gives a report of some spices consumed in Nigeria for the presence of some heavy metals and polycyclic aromatic hydrocarbons. Some spices are called flavour enhancers. A flavour enhancer increases the way food is perceived and desired without changing the actual flavour of the prepared dish. Flavor enhancers do this by affecting your taste buds. Monosodium glutamate (MSG) popularly known as Maggi, is an example of a flavour enhancer. It intensifies the flavour of savory foods (Susan and Anne, 1998).

A flavouring, on the other hand, is an ingredient that actually changes the natural flavour of the foods to which it is added. This enhances the experience of tasting the dish. Both spices and flavorings improve or strengthen the natural flavour of foods ((Singh et al, 2015). Apart from adding flavour to foods, some spices also contain medical and health benefits such as lowering of cholesterol levels, removal of scalp itching and peeling caused by candidiasis, relieve arthritis and back pain, healing of colds, sinus infections and sore throats, burn calories, speed up metabolism, cancer-fighting and anti-inflammatory properties, help to fight communicable diseases as well as lower the levels of bad cholesterol and triglycerides in the blood (Singh et al, 2015)(Jashankar et al, 2014). According to Susan and Anne, (1998), seasonings, which are also ingredients used in soup making are composed of notable list of phyto-nutrients, essential oils, antioxidants, minerals and vitamins that are essential for good health.

*John, Godson Nnamdi, Department of Applied Chemical Science Laboratory Technology, Faculty of Science Laboratory Technology, University of Benin.

Ecchefu, Maduabuchi Francis, Department of Chemical/Petroleum Technology, School of Science Laboratory Technology, University of Port Harcourt

The use of spices in food preparation has become an integral part of life over the centuries, in many parts of the world (Otunola *et al.*, 2010). Other examples of spices common in Nigeria include cinnamon, curry, spice city, Lasor, Onga, rosemary, knorr, etc (Brown *et al.*, 2003). However, wide variations in concentrations of trace metals have been reported in some spices (Akpanyung, *et al.*, 2005). Heavy metal contamination in the food chain is caused by environmental pollution and lead (Pb) and copper (Cu) have aroused considerable concern (Nnorom *et al.*, 2007). In Nigeria spices are manufactured as bouillon cubes and other well-packaged powdery forms with different brand names. They contain salt, monosodium glutamate (MSG), and some spices as indicated by the manufacturers and are used extensively in food preparation in most homes and restaurants (Oladele *et al.*, 2015). MSG is the sodium salt of the amino acid glutamic acid and has long been recognized as a flavour enhancer. It is sold as a fine white crystalline substance similar in appearance to salt or sugar. It is considered a primary taste, umami. Pure MSG does not have a pleasant taste until it is combined with a consonant savory smell (Scatter *et al.*, 1989). Glutamate is found in most foods especially foods that are high in protein such as meat, fish, and dairy products. Other sources include mushrooms and tomatoes. In Africa, the current quest for easy or fast way to prepare foods has brought with it a progressive loss of important component of the African food culture. The African richly enormous variety of food spices and condiments are today gradually being replaced by the largenumber of bouillon cubes and spices in the market (Stadman, 1990). Bouillon cubes are taste enhancers and are added to foods to augment the taste properties of food (Akpanyung, 2005). The major active ingredients in bouillon cubes are salt (NaCl) and monosodium glutamate (MSG). Other raw material used include soybeans, locust beans, onion, tomato, hydrogenated palm oil (HPO), caramel, hydrolyzed plant/vegetable protein and natural spices (RMRDC, 2003). Foods spices are known to contain calcium, magnesium, potassium, sodium and cadmium, lead (Muhammad, 2011). This study was carried out to identify the levels of macro minerals and heavy metals content and the safety assessment of some seasonings and bouillon cubes readily consumed in Nigeria. Four toxic metals (Pb, Cu, Cd and Zn) and four essential micro minerals (Mg, Ca Na and K) were determined to assess human ingestion of these metals from the consumption of foods prepared with these seasonings. In Nigeria, spices are commonly used in the production of dishes and drinks such as: pepper soup, jollof rice, yam pottage, all types of soups and stews, local drinks such as Zobo, ginger and other fruit drinks. Though, spices are used in small quantities, they contribute to the nutrient content of the food (Stadman, 1990).

These spices are manufactured as bouillon cubes and in powdered form which are packaged with different brand names and are used extensively in food preparation (Otunola *et al.*, 2010; Singh *et al.*, 2015). They are added to soups, stews, puddings and sometimes as stimulants which are mixed along with other beverages and used as pepper soup ingredients (Mudgil and Barak, 2017; Hudson *et al.*, 1984). A bouillon cube is a dehydrated stock formed into a small cube or dried granules. It is formed into a small cube of about 15mm wide. It is typically made by dehydrating chicken stock, fish, meat stock, a small portion of fat, salt, seasonings that are dried or dehydrated and shaping them into a small cube. Bouillon cube is also available in a granular form. When placed in water, the cubes dissolve and its flavors are released. (Ekpo and Jimmy, 2005). Spices and Food colorants which are contaminated by heavy metals from the atmosphere, soil and water pose health risks to humans and animals (Kabata-Pendias, 2001). This study investigated the presence of heavy metals and Poly aromatic hydrocarbons and also the health risks. The growing use of these spices as flavouring, colouring and preservative agent gotten/prepared from variety of plant seed, fruit, root and vegetables which could have been contaminated by heavy metals during agricultural cultivation, industrial or human activities and from leaded gasoline (Khairiah, 2009) or as deliberate ingredients by producers interested only in profit making from the business.

MATERIALS AND METHODS

Sample Collection

Five different commercially available brands of spices were purchased at random from local shops around the Port Harcourt Metropolis, Nigeria. The samples were registered by NAFDAC at the time of this study with appropriate batch number and expiry date. The samples were hermetically sealed, and purchased after preliminary investigations had indicated that they were the most consumed seasoning in the metropolis. The spices were: Larsor chicken spice, Mr. Chef pepparsoup spice, Onga, spicity jollof rice spice and Mr. Chef Jollof rice spice.

Sample Digestion (wet digestion)

Method: APHA 3030F (Nitric acid-Hydrochloric acid digestion)

A total volume of 100 ml hydrochloric acid and nitric acid in the ratio of 3:1 was mixed to form an aqua regia solution. Sample, (1g) was weighed into a conical flask, 20 ml of the mixed acid was added to the flask. This was digested using a fume cupboard with hot plate until the appearance of white fumes. It was allowed to cool and filtered into a 100 ml volumetric flask and made up to the mark with distilled water. Portions of the solution were used for atomic absorption spectrophotometer analysis.

All the metals were analyzed using the above method but at every point of reading a particular metal with Atomic Absorption spectrophotometer, the cathode lamp for that particular metal is used.

Polycyclic Aromatic Hydrocarbon

Method No: US EPA 8015 (Extraction for sediment samples)

- . Ten (10 g) of well mixed wet sample was weighed into a beaker already washed and rinsed with acetone.
- . About 25 ml of 1 + 1 dichloromethane and acetone

were added into it. The sample was sonicated for about 20 mins and centrifuged for 10 mins. The solvent was extracted through a filter paper containing 5g of anhydrous sodium sulphate and preconditioned with 1+ 1 DCM and acetone. Then the second 50 ml of 1+1 DCM and acetone was added to the conical flask. The extraction was repeated combining extracts in the Erlenmeyer flask; this was concentrated into 5 ml using a rotary evaporator. For the aromatics and in the same column 10 ml of DCM was added and eluted. This was used for the GC analysis.

RESULTS

Table 1: Metals and Polycyclic Aromatic Hydrocarbon Levels in Samples

S/No	PARAMETERS	SAMPLES				
		A	B	C	D	E
1	Pb (ppm)	3.42	0.32	1.22	0.46	3.01
2	Cu (ppm)	8.12	3.23	5.43	1.35	8.00
3	Cr (ppm)	0.45	0.12	0.23	0.08	0.22
4	Fe (ppm)	55.21	31.32	38.11	22.21	47.32
5	Zn (ppm)	11.23	8.22	6.45	3.55	7.23
6	PAHs (%)	82.43	57.78	100.27	57.78	129.09

Table 2: Levels of Polycyclic Aromatic Hydrocarbons in Sample A

S/NO.	Target Compounds	R.T.	Qlon	Response	Conc.Units (ppm)	Dev(Min) Qvalue
1	Naphthalene	4.009	128	592	0.00	61
2	Acenaphthylene	5.434	152	156	2.10	1
3	Acenaphthene	5.584	153	242	17.02	1
4	Fluorene	6.105	166	413	12.02	75
5	Phenanthrene	7.102	178	15969	7.01	63
6	Anthracene	7.102	178	15969	11.00	61
7	O-Terphenyl	7.379	230	4121994	0.66	99
8	Fluoranthene	8.272	202	54053	2.03	91
9	Pyrene	8.489	202	72217	3.00	96
10	Benz(a)anthracene	9.887	228	24638	11.07	63
11	Chrysene	9.887	228	24638	0.01	58
12	Benzo(b)fluoranthene	12.046	252	118256	0.30	59
13	Benzo(k)fluoranthene	12.046	252	118256	5.00	59
14	Benzo(a)pyrene	12.785	252	15383	4.11	59
15	Dibenzo(a,h)anthracene	15.458	276	307	1.00	36
16	Indeno(1,2,3-cd)pyrene	16.140	276	498	3.00	56
17	Benzo(g,h,i)perylene	16.181	276	160	3.10	1
	Total				82.43	

Table 2: Levels of Polycyclic Aromatic Hydrocarbon in Sample B

S/NO.	Target Compounds	R.T.	Qlon	Response	Conc.Units (ppm)	Dev(Min) Qvalue
1	Naphthalene	4.021	128	259	1.00	68
2	Acenaphthylene	5.441	152	96	10.00	1
3	Acenaphthene	5.588	153	156	3.00	1
4	Fluorene	6.135	166	455	0.00	15
5	Phenanthrene	7.098	178	6016	0.00	72
6	Anthracene	7.098	230	2693965	0.00	65
7	O-Terphenyl	7.376	202	17315	0.77	99
8	Fluoranthene	8.279	202	12119	0.01	85
9	Pyrene	8.482	228	220	10.00	59
10	Benz (a) anthracene	9.816	228	10197	3.00	1
11	Chrysene	9.880	252	149	0.00	65
12	Benzo (b) Fluoranthene	11.852	252	155	10.00	26
13	Benzo (k) Fluoranthene	11.945	252	149	2.00	6
14	Benzo (e) Pyrene	12.680	276	146	1.00	1
15	Dibenzo (a,h) anthracene	15.724	276	310	5.00	1
16	Indeno (1,2,3-cd) Pyrene	16.076	276	310	5.00	1
17	Benzo (ghi) Perylene	16.361	276	290	7.00	61
	Total				57.78	

Table 3: Levels of Polycyclic Aromatic Hydrocarbons in Sample C

S/NO.	Target Compound	R.T.	Qlon	Response	Conc. Units(ppm)	Dev(Min) Qvalue
1	Naphthalene	4.01	128	568	0.00	72
2	Acenaphthylene	5.456	152	3203	0.01	73
3	Acenaphthene	5.621	153	374	20.00	1
4	Fluorene	6.109	166	6035	0.03	72
5	Phenanthrene	7.057	178	51628	0.02	73
6	Anthracene	7.106	178	62704	10.01	69
7	O-Terphenyl	7.368	230	3864508	0.07	98
8	Fluoranthene	8.230	202	91095	0.03	93
9	Pyrene	8.455	202	113198	0.01	90
10	Benz(a) anthracene	9.850	228	15441	20.01	90
11	Chrysene	9.880	228	8891	10.00	62
12	Benzo (b) Fluoranthene	11.983	252	242023	0.02	75
13	Benzo (k) Fluoranthene	11.983	252	242023	10.00	75
14	Benzo (e) Pyrene	12.736	252	94804	0.04	66
15	Dibenzo (a,h) anthracene	15.728	276	2715	20.01	27
16	Indeno (1,2,2-cd) Pyrene	15.983	276	455	10.00	75
17	Benzo ghi) Perylene	16.384	276	12508	0.01	78
	Total				100.27	

Table 4: Levels of Polycyclic Aromatic Hydrocarbons in Sample D

S/NO.	Target Compounds	R.T.	Qlon	Response	Conc.Units (ppm)	Dev.(Mins) Qvalue
1	Naphthalene	4.021	128	259	1.00	68
2	Acenaphthylene	5.441	152	96	10.00	1
3	Acenaphthene	5.588	153	156	3.00	1
4	Fluorene	6.135	166	455	0.00	15
5	Phenanthrene	7.098	178	6016	0.00	72
6	Anthracene	7.098	178	6061	0.00	65
7	O-Terphenyl	7.376	230	26939650	0.77	99
8	Fluoranthene	8.279	202	17315	0.01	85
9	Pyrene	8.482	202	12119	10.00	59
10	Benz (a) anthracene	9.816	228	220	3.00	1
11	Chrysene	9.880	228	10197	0.00	65
12	Benzo (b) fluoranthene	11.852	252	149	10.00	26
13	Benzo (k) fluoranthene	11.942	252	155	2.00	6
14	Benzo (e) Pyrene	12.680	252	149	8.00	1
15	Dibenzo (a, h) anthracene	15.724	276	146	5.00	1
16	Indeno(1,2,3-cd) pyrene	16.076	276	310	5.00	1
17	Benzo (ghi)perylene	16.361	276	290	1.00	61
Total					58.78	

Table 5: Levels of Polycyclic Aromatic Hydrocarbons in Sample E

S/NO.	Target Compound	R.T.	Qlon	Response	Conc.Units (ppm)	Dev(Min) Qvalue
1	Naphthalene	4.009	128	216	30.30	1
2	Acenaphthylene	5.449	152	242	0.30	1
3	Acenaphthene	5.595	153	222	30.00	1
4	Fluorene	6.161	166	1237	3.03	32
5	Phenanthrene	7.113	178	7627	0.40	70
6	Anthracene	7.113	178	7627	2.10	70
7	O-Terphenyl	7.379	230	3013623	1.10	99
8	Fluoranthene	8.290	202	21445	0.02	86
9	Pyrene	8.515	202	23081	0.01	62
10	Benz (a) anthracene	9.973	228	166681	10.05	62
11	Chrysene	9.973	228	166681	11.30	55
12	Benzo (b) fluoranthene	12.163	252	2252	5.20	90
13	Benzo (k) fluoranthene	11.805	252	326	10.10	11
14	Benzo (e) pyrene	12.609	252	214	0.00	1
15	Dibenzo (a,h) anthracene	15.506	276	200	0.02	1
16	Indeno (1,2,3-cd) pyrene	16.140	276	318	25.08	41
17	Benzo (ghi) perylene	16.140	276	318	0.08	27
Total					129.09	

DISCUSSION

Lead was detected in all five spice samples as can be seen in table 1. Lead is a toxic metal of public health concern with no known biological function and reported to induce toxicity at concentrations (Jaishankar, 2014). The lead content in spices and food colorings could be attributed to the addition of lead during processing to impart colour, sweet taste or increase the weight of these products (Kakosy *et al*, 1996). Some studies reported that traffic density also increases the lead burden in the environment thereby increasing the lead content in vegetation (Nabulo, 2004).

Lead pollution has been shown to be commensurate with population/vehicular density (Ansari *et al*, 2004). Generally, lead contaminations occur in vegetables grown on contaminated soils, through air deposition or through sewage sludge/wastewater application (Opaluwa *et al*, 2012). Lead toxicity is known to inhibit the action of these enzymes because they have free sulfhydryl groups. This is particularly noted with the precursors of haem, and leads to a decrease in haem synthesis, and hence to anemia. Irrespective of the way a lead compound enters the body, it first penetrates the initial cellular barrier before reaching the intracellular fluid.

The compound then penetrates the capillary blood vessels and thus enters the circulatory system which distributes it throughout the body (Petersdorf, 1999). Lead poisoning is a global reality, and fortunately is not a very common clinical diagnosis yet in Nigeria except for a few occupational exposures (Anetor *et al*, 1999). In this study, the concentrations of Pb in all five samples are generally higher than the permissible levels by FAO/WHO of 0.001ppm and hence pose health risks to consumers. Copper was essential trace mineral in the human body, functioning in enzymatic reactions, connective tissue formation, nervous system health, immune function, antioxidant defense, etc. However at excessive concentration levels, copper tends to become toxic with neurological symptoms including Alzheimer's disease. It also interferes with absorption and metabolism of other minerals and causes Wilson's disease. In this study, the concentration levels of copper in all selected spice samples are above the FAO/WHO permissible standard of 0.02 ppm, hence posing health risks to consumers. Chromium helps in glucose and lipid metabolism in the body and also in weight management but becomes toxic in high concentrations leading to environmental pollution, kidney and liver damage due to prolonged exposure. In this study, chromium concentration levels are slightly higher than FAO/WHO standard of 0.005ppm. Sample D is the closest to this standard value and thus poses the least health risk to consumers. Iron plays a key component in haemoglobin, a protein in red blood cells responsible for transporting oxygen from the lungs to the rest of the body. It also supports energy production as it involves electron transport. Iron is important for immune functions and health and plays cognitive and muscle functions. Iron deficiency can lead to anemia characterized by fatigue, weakness, impaired cognitive function and decreased immunity. Whereas excessive iron accumulation can damage organs particularly the liver, heart, pancreas, and increase the risk of diabetes, liver cancer and heart disease. In our current study, the levels of Fe in samples B and D are relatively close to standard requirements of 24 ppm while the others exceed by large margins and are thus harmful to health. Zinc metal plays a crucial role in supporting the immune system, promoting wound healing and aiding in cell growth and division. It also helps in taste and smell sensations. However it is important to note that excessive zinc intake can have risks such as the interference of mineral absorption and effectiveness of drugs, neurological crisis, diarrhea, severe headaches, vomiting, loss of appetite, etc. In our study, only sample D meets the WHO standard level of 3 ppm. All four other samples therefore pose health risks as they are higher than this standard. Polycyclic Aromatic Hydrocarbons in Spices and Food Colorants In this work, seventeen different polycyclic aromatic hydrocarbon target compounds were tested in all five (5) samples. In Nigeria the Standard Organization

(SON), Federal Ministry of Environment, National Agency For Food and Drug Administration and Control (NAFDAC) and the National Environmental Standards and Regulations Enforcement Agency (NESREA) are bodies which play roles in regulating, monitoring and enforcing standards related to PAHs content in various contexts within Nigeria to ensure that their levels do not pose a significant risk to public health safety. Naphthalene concentration in samples A, B, C, and D is below 2 ppm and is considered safe by WHO and OSHA (Occupational Safety and Health Administration) but in sample E it is extremely high and pose health risks to consumers. Acenaphthylene grossly exceeded maximum 8 ppm concentration in samples B and D. Pyrene, fluorene and other PAHs in the samples all exceeded recommended levels and thus pose serious health risks to consumers.

CONCLUSION

This investigation into the levels of heavy metals and polycyclic aromatic hydrocarbons (PAHs) in popular Nigerian spices and food colorants provides critical evidence regarding the current state of food safety. The analytical results confirm that several samples contain contaminant levels far exceeding the safety thresholds established by international and local regulatory agencies. The discovery of toxic elements like lead and cadmium, alongside essential but over-concentrated metals such as copper and zinc, is alarming. These substances are linked to severe physiological outcomes, including renal dysfunction, neurological impairments, and various forms of cancer. Furthermore, the presence of carcinogenic PAHs suggests a significant risk of bioaccumulation, which could lead to chronic, long-term health crises for the Nigerian populace.

REFERENCES

- Abdullahi MS, Uzairu A, Okunola OJ., 2008. Determination of some trace metal levels in onion leaves from irrigated farmlands on the bank of River Challawa, Nigeria. *Afr. J. Biotechnol.* 7 (10):1526-1529.
- Akpanyung. E.O., 2005. Proximate and mineral element composition of bouillon cubes produced in Nigeria, *Pakistan Journal of Nutrition*, 4(5), 327-329.
- Anetor JI, Adeniyi FA, Taylor GO., 1999. Biochemical indicators of metabolic poisoning associated with lead based occupations in nutritionally disadvantaged communities. *Afr. J. Med. Med. Sci.* 28: 9-12.
- Ansari TM, Ikram N, Najam-ul-Haq M, Fayyaz O, Ghafoor I, Khalid N., 2004. Essential trace metal (Zn, Mn, Cu, and Fe) levels in plants of medicinal importance. *J. Biol. Sci.* 4:95-9

- Brown S., Chaney R.L., Hallfrisch J.G., Xue Q., 2003. Effects of Biosolids Processing on Lead Bioavailability in Urban Soil. *J. of Environ. Qual.* 32, (1), 100.
- Ekpo AJ, Jimmy EO., 2005. Nutritional effect of spices from Roasted Chicken and Beef Meat consumption. *Pak. J. Nutr.* 4 (6):428-43.
- Herndon, J., 2010. Medical benefits of food seasonings.
- Hudson TS, Subramanian S, Allen R.J., 1984. Determination of pantothenic acid, biotin, and vitamin B12 in nutritional products. *Journal-Association of Official Analytical Chemists.* 67 (5):994-998.
- Jaishankar M, Tseten T, Anbalagan N, Mathe BB, Beeregowda K. N., 2014. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol*, 7(2): 60–72. DOI: 10.2478/intox-2014-000
- Kabata-Pendias A., Pendias H., 2001. Trace Element in Soil and Plant. USA: CRC Press. 3rd ed.
- Kakosy T, Hudak A, Baray M., 1996. Lead intoxication epidemic caused by ingestion of contaminated ground paprika. *J. Toxicol. Clin. Toxicol.* 34:507 -511.
- Khairiah J., Ding-Woei Y., Habibah J., Ahmed-Mahir R., Aminah A., Ismail B.S., 2009. Concentration of heavy metals in guava plant parts and soil in the Sungai Wangi Plantation, Perak, Malaysia. *International Journal Agriculture Research.* 4: 310–316.
- Mudgil, D., and Barak, S., 2017. Functional Foods: Sources and Health Benefits. Scientific Publishers.
- Muhammad HL, Kabir AY, Adeleke KB., 2011. Mineral elements and heavy metals in selected food seasonings consumed in Minna metropolis. *International Journal of Applied Biological Research* 3(1): 108-113.
- Nabulo G., 2004. Assessment of Heavy Metal contamination of Food Crops and Vegetables from motor vehicle emissions in Kampala City, Uganda. Department of Botany Makerere University, Kampala. A technical report submitted to IDRC-AGROPOLIS.
- Nnorom IC, Osibanjo O, Ogugua K (2007) Trace Heavy Metal Levels of Some Bouillon Cubes, and Food Condiments Readily Consumed in Nigeria. *Pakistan Journal of Nutrition* 6(2): 122-127.
- Oladele, A. T., and Fadare, O. O., 2015. Heavy metals and proximate composition of forest leafy vegetables in oil producing area of Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 8(4), 451-463.
- Opaluwa. O. D., Aremu. M. O., Ogbo. L. O., 2012. "Heavy metal concentrations in soils, plant leaves and crops grown around dump sites in Lafia metropolis, Nasarawa State, Nigeria, "Advances in Applied Science Research, .(3) (2)780–784, 2012
- Otunola. G. A, Oloyede. O.B, Oladiji. A.T and Afolayan. A.J., 2010. Comparative analysis of the chemical composition of three spices *Allium sativum*, *L.zingiber officinale* rocs and *Capsicum frutescens* L commonly consumed in Nigeria, *African Journal of Biotechnology*, 9(41):6927-693.
- Petersdorf, R. G., Martin, J. B. and Fauci, A. S., 1999. Harrison Principle of Internal Medicine. 2: 12th ed. McGraw-Hill, New York pp 2182-7.
- RMRDC, 2003. Multi-Disciplinary Committee Report of the Tschno-Economics Survey of Food, Beverage and Tobacco Sector (4th Update). Raw Materials, Research and Development Council, GFederal Ministry of Science and Technology Abuja, Nigeria.
- Satter A, Wahid M, Durrani SK., 1989. Concentration of selected heavy metals in spices, dry fruits and plants nuts. *Plant Food Hum. Nutr.* 39(4):279-286.
- Singh, A., and Prasad, S. M., 2015. Remediation of heavy metal contaminated ecosystem: an overview on technology advancement. *International Journal of Environmental Science and Technology*, 12, 353-366.
- Stadtman TC., 1990. Selenium biochemistry. *Annual Review of Biochemistry*; 59(1):111-127
- Susan, M. and Anne, P. (1998). Tropical and Sub-tropical Foods. 2nd Ed. MacMillan Publications New York 154 – 170.